

# Marie Anne Pringle

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3045762/publications.pdf>

Version: 2024-02-01

8  
papers

307  
citations

1307594  
7  
h-index

1588992  
8  
g-index

10  
all docs

10  
docs citations

10  
times ranked

533  
citing authors

#	ARTICLE	IF	CITATIONS
1	ERdj5 Is the ER Reductase that Catalyzes the Removal of Non-Native Disulfides and Correct Folding of the LDL Receptor. <i>Molecular Cell</i> , 2013, 50, 793-804.	9.7	116
2	Cytosolic thioredoxin reductase 1 is required for correct disulfide formation in the <scp>ER</scp>. <i>EMBO Journal</i> , 2017, 36, 693-702.	7.8	65
3	<scp>ER</scp> p18 regulates activation of <scp>ATF</scp> 6 $\hat{1}$ during unfolded protein response. <i>EMBO Journal</i> , 2019, 38, e100990.	7.8	44
4	Folding of a single domain protein entering the endoplasmic reticulum precedes disulfide formation. <i>Journal of Biological Chemistry</i> , 2017, 292, 6978-6986.	3.4	34
5	The membrane topology of vitamin K epoxide reductase is conserved between human isoforms and the bacterial enzyme. <i>Biochemical Journal</i> , 2016, 473, 851-858.	3.7	15
6	The mammalian cytosolic thioredoxin reductase pathway acts via a membrane protein to reduce ER-localised proteins. <i>Journal of Cell Science</i> , 2020, 133, .	2.0	15
7	Activation of the UPR sensor ATF6 $\hat{1}$ is regulated by its redox-dependent dimerization and ER retention by ERp18. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2022, 119, e2122657119.	7.1	15
8	A cytosolic reductase pathway is required for efficient N-glycosylation of an STT3B-dependent acceptor site. <i>Journal of Cell Science</i> , 2021, 134, .	2.0	3