Joseph M Connors

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3047392/publications.pdf Version: 2024-02-01



LOSEDH M CONNORS

| # | Article | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Circos: An information aesthetic for comparative genomics. Genome Research, 2009, 19, 1639-1645. | 2.4 | 9,003 |
| 2 | Revised Response Criteria for Malignant Lymphoma. Journal of Clinical Oncology, 2007, 25, 579-586. | 0.8 | 4,061 |
| 3 | Confirmation of the molecular classification of diffuse large B-cell lymphoma by immunohistochemistry using a tissue microarray. Blood, 2004, 103, 275-282. | 0.6 | 3,574 |
| 4 | The Use of Molecular Profiling to Predict Survival after Chemotherapy for Diffuse Large-B-Cell Lymphoma. New England Journal of Medicine, 2002, 346, 1937-1947. | 13.9 | 3,474 |
| 5 | Report of an International Workshop to Standardize Response Criteria for Non-Hodgkin's Lymphomas. Journal of Clinical Oncology, 1999, 17, 1244-1244. | 0.8 | 3,209 |
| 6 | Somatic mutations altering EZH2 (Tyr641) in follicular and diffuse large B-cell lymphomas of germinal-center origin. Nature Genetics, 2010, 42, 181-185. | 9.4 | 1,504 |
| 7 | Genetics and Pathogenesis of Diffuse Large B-Cell Lymphoma. New England Journal of Medicine, 2018, 378, 1396-1407. | 13.9 | 1,443 |
| 8 | Frequent mutation of histone-modifying genes in non-Hodgkin lymphoma. Nature, 2011, 476, 298-303. | 13.7 | 1,428 |
| 9 | Chronic active B-cell-receptor signalling in diffuse large B-cell lymphoma. Nature, 2010, 463, 88-92. | 13.7 | 1,402 |
| 10 | Results of a Pivotal Phase II Study of Brentuximab Vedotin for Patients With Relapsed or Refractory Hodgkin's Lymphoma. Journal of Clinical Oncology, 2012, 30, 2183-2189. | 0.8 | 1,332 |
| 11 | Prediction of Survival in Follicular Lymphoma Based on Molecular Features of Tumor-Infiltrating Immune Cells. New England Journal of Medicine, 2004, 351, 2159-2169. | 13.9 | 1,293 |
| 12 | Oncogenically active MYD88 mutations in human lymphoma. Nature, 2011, 470, 115-119. | 13.7 | 1,292 |
| 13 | The revised International Prognostic Index (R-IPI) is a better predictor of outcome than the standard IPI for patients with diffuse large B-cell lymphoma treated with R-CHOP. Blood, 2007, 109, 1857-1861. | 0.6 | 1,193 |
| 14 | Tumor-Associated Macrophages and Survival in Classic Hodgkin's Lymphoma. New England Journal of Medicine, 2010, 362, 875-885. | 13.9 | 1,141 |
| 15 | Molecular Diagnosis of Primary Mediastinal B Cell Lymphoma Identifies a Clinically Favorable Subgroup of Diffuse Large B Cell Lymphoma Related to Hodgkin Lymphoma. Journal of Experimental Medicine, 2003, 198, 851-862. | 4.2 | 1,002 |
| 16 | Brentuximab Vedotin (SGN-35) in Patients With Relapsed or Refractory Systemic Anaplastic Large-Cell Lymphoma: Results of a Phase II Study. Journal of Clinical Oncology, 2012, 30, 2190-2196. | 0.8 | 890 |
| 17 | Introduction of Combined CHOP Plus Rituximab Therapy Dramatically Improved Outcome of Diffuse Large B-Cell Lymphoma in British Columbia. Journal of Clinical Oncology, 2005, 23, 5027-5033. | 0.8 | 874 |
| 18 | Molecular subtypes of diffuse large B-cell lymphoma arise by distinct genetic pathways. Proceedings of the National Academy of Sciences of the United States of America, 2008, 105, 13520-13525. | 3.3 | 868 |

| # | Article | IF | CITATIONS |
|----|--|------|-----------|
| 19 | The proliferation gene expression signature is a quantitative integrator of oncogenic events that predicts survival in mantle cell lymphoma. Cancer Cell, 2003, 3, 185-197. | 7.7 | 848 |
| 20 | Molecular Diagnosis of Burkitt's Lymphoma. New England Journal of Medicine, 2006, 354, 2431-2442. | 13.9 | 824 |
| 21 | Concurrent Expression of MYC and BCL2 in Diffuse Large B-Cell Lymphoma Treated With Rituximab Plus Cyclophosphamide, Doxorubicin, Vincristine, and Prednisone. Journal of Clinical Oncology, 2012, 30, 3452-3459. | 0.8 | 824 |
| 22 | Oncogenic <i>CARD11</i> Mutations in Human Diffuse Large B Cell Lymphoma. Science, 2008, 319, 1676-1679. | 6.0 | 784 |
| 23 | ALKâ^' anaplastic large-cell lymphoma is clinically and immunophenotypically different from both ALK+ ALCL and peripheral T-cell lymphoma, not otherwise specified: report from the International Peripheral T-Cell Lymphoma Project. Blood, 2008, 111, 5496-5504. | 0.6 | 784 |
| 24 | Burkitt lymphoma pathogenesis and therapeutic targets from structural and functional genomics. Nature, 2012, 490, 116-120. | 13.7 | 759 |
| 25 | An enhanced International Prognostic Index (NCCN-IPI) for patients with diffuse large B-cell lymphoma treated in the rituximab era. Blood, 2014, 123, 837-842. | 0.6 | 693 |
| 26 | MYC gene rearrangements are associated with a poor prognosis in diffuse large B-cell lymphoma patients treated with R-CHOP chemotherapy. Blood, 2009, 114, 3533-3537. | 0.6 | 566 |
| 27 | Brentuximab Vedotin with Chemotherapy for Stage III or IV Hodgkin's Lymphoma. New England Journal of Medicine, 2018, 378, 331-344. | 13.9 | 564 |
| 28 | MHC class II transactivator CIITA is a recurrent gene fusion partner in lymphoid cancers. Nature, 2011, 471, 377-381. | 13.7 | 551 |
| 29 | Lymphomas with concurrent BCL2 and MYC translocations: the critical factors associated with survival. Blood, 2009, 114, 2273-2279. | 0.6 | 523 |
| 30 | Determining cell-of-origin subtypes of diffuse large B-cell lymphoma using gene expression in formalin-fixed paraffin-embedded tissue. Blood, 2014, 123, 1214-1217. | 0.6 | 518 |
| 31 | Integration of gene mutations in risk prognostication for patients receiving first-line immunochemotherapy for follicular lymphoma: a retrospective analysis of a prospective clinical trial and validation in a population-based registry. Lancet Oncology, The, 2015, 16, 1111-1122. | 5.1 | 483 |
| 32 | Mantle Cell Lymphoma: A Clinicopathologic Study of 80 Cases. Blood, 1997, 89, 2067-2078. | 0.6 | 448 |
| 33 | Randomized Comparison of ABVD and MOPP/ABV Hybrid for the Treatment of Advanced Hodgkin's Disease: Report of an Intergroup Trial. Journal of Clinical Oncology, 2003, 21, 607-614. | 0.8 | 438 |
| 34 | Gene expression signatures delineate biological and prognostic subgroups in peripheral T-cell lymphoma. Blood, 2014, 123, 2915-2923. | 0.6 | 435 |
| 35 | Analysis of multiple biomarkers shows that lymphoma-associated macrophage (LAM) content is an independent predictor of survival in follicular lymphoma (FL). Blood, 2005, 106, 2169-2174. | 0.6 | 427 |
| 36 | <i>De novo</i> transcriptome assembly with ABySS. Bioinformatics, 2009, 25, 2872-2877. | 1.8 | 371 |

| # | Article | IF | CITATIONS |
|----|--|------|-----------|
| 37 | ABVD Alone versus Radiation-Based Therapy in Limited-Stage Hodgkin's Lymphoma. New England Journal of Medicine, 2012, 366, 399-408. | 13.9 | 360 |
| 38 | Molecular Pathogenesis of Hodgkin's Lymphoma: Increasing Evidence of the Importance of the Microenvironment. Journal of Clinical Oncology, 2011, 29, 1812-1826. | 0.8 | 350 |
| 39 | Mutational and structural analysis of diffuse large B-cell lymphoma using whole-genome sequencing. Blood, 2013, 122, 1256-1265. | 0.6 | 349 |
| 40 | Diffuse large B-cell lymphoma subgroups have distinct genetic profiles that influence tumor biology and improve gene-expression-based survival prediction. Blood, 2005, 106, 3183-3190. | 0.6 | 348 |
| 41 | Survival of Patients With Peripheral T-Cell Lymphoma After First Relapse or Progression: Spectrum of Disease and Rare Long-Term Survivors. Journal of Clinical Oncology, 2013, 31, 1970-1976. | 0.8 | 335 |
| 42 | Prognostic Significance of Diffuse Large B-Cell Lymphoma Cell of Origin Determined by Digital Gene Expression in Formalin-Fixed Paraffin-Embedded Tissue Biopsies. Journal of Clinical Oncology, 2015, 33, 2848-2856. | 0.8 | 334 |
| 43 | Population-Based Analysis of Incidence and Outcome of Transformed Non-Hodgkin's Lymphoma. Journal of Clinical Oncology, 2008, 26, 5165-5169. | 0.8 | 333 |
| 44 | Five-year survival and durability results of brentuximab vedotin in patients with relapsed or refractory Hodgkin lymphoma. Blood, 2016, 128, 1562-1566. | 0.6 | 324 |
| 45 | Whole transcriptome sequencing reveals recurrent NOTCH1 mutations in mantle cell lymphoma. Blood, 2012, 119, 1963-1971. | 0.6 | 313 |
| 46 | CNS International Prognostic Index: A Risk Model for CNS Relapse in Patients With Diffuse Large B-Cell Lymphoma Treated With R-CHOP. Journal of Clinical Oncology, 2016, 34, 3150-3156. | 0.8 | 313 |
| 47 | Enteropathy-associated T-cell lymphoma: clinical and histological findings from the International Peripheral T-Cell Lymphoma Project. Blood, 2011, 118, 148-155. | 0.6 | 308 |
| 48 | Randomized Comparison of ABVD Chemotherapy With a Strategy That Includes Radiation Therapy in Patients With Limited-Stage Hodgkin's Lymphoma: National Cancer Institute of Canada Clinical Trials Group and the Eastern Cooperative Oncology Group. Journal of Clinical Oncology, 2005, 23, 4634-4642. | 0.8 | 305 |
| 49 | BCL2 Expression Is a Prognostic Marker for the Activated B-Cell–Like Type of Diffuse Large B-Cell Lymphoma. Journal of Clinical Oncology, 2006, 24, 961-968. | 0.8 | 277 |
| 50 | Cooperative Epigenetic Modulation by Cancer Amplicon Genes. Cancer Cell, 2010, 18, 590-605. | 7.7 | 263 |
| 51 | BCL2 Translocation Defines a Unique Tumor Subset within the Germinal Center B-Cell-Like Diffuse Large B-Cell Lymphoma. American Journal of Pathology, 2004, 165, 159-166. | 1.9 | 262 |
| 52 | Flavopiridol in Untreated or Relapsed Mantle-Cell Lymphoma: Results of a Phase II Study of the National Cancer Institute of Canada Clinical Trials Group. Journal of Clinical Oncology, 2003, 21, 1740-1745. | 0.8 | 261 |
| 53 | Double-Hit Gene Expression Signature Defines a Distinct Subgroup of Germinal Center B-Cell-Like Diffuse Large B-Cell Lymphoma. Journal of Clinical Oncology, 2019, 37, 190-201. | 0.8 | 257 |
| 54 | Randomized Phase III Trial of ABVD Versus Stanford V With or Without Radiation Therapy in Locally Extensive and Advanced-Stage Hodgkin Lymphoma: An Intergroup Study Coordinated by the Eastern Cooperative Oncology Group (E2496). Journal of Clinical Oncology, 2013, 31, 684-691. | 0.8 | 256 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Brentuximab vedotin combined with ABVD or AVD for patients with newly diagnosed Hodgkin's lymphoma: a phase 1, open-label, dose-escalation study. Lancet Oncology, The, 2013, 14, 1348-1356. | 5.1 | 251 |
| 56 | Point mutations and genomic deletions in CCND1 create stable truncated cyclin D1 mRNAs that are associated with increased proliferation rate and shorter survival. Blood, 2007, 109, 4599-4606. | 0.6 | 226 |
| 57 | State-of-the-Art Therapeutics: Hodgkin's Lymphoma. Journal of Clinical Oncology, 2005, 23, 6400-6408. | 0.8 | 221 |
| 58 | Treatment-Related Myelodysplasia and Acute Leukemia in Non-Hodgkin's Lymphoma Patients. Journal of Clinical Oncology, 2003, 21, 897-906. | 0.8 | 215 |
| 59 | Molecular and Genetic Characterization of MHC Deficiency Identifies EZH2 as Therapeutic Target for Enhancing Immune Recognition. Cancer Discovery, 2019, 9, 546-563. | 7.7 | 213 |
| 60 | Durable remissions in a pivotal phase 2 study of brentuximab vedotin in relapsed or refractory Hodgkin lymphoma. Blood, 2015, 125, 1236-1243. | 0.6 | 199 |
| 61 | Impact of Concordant and Discordant Bone Marrow Involvement on Outcome in Diffuse Large B-Cell Lymphoma Treated With R-CHOP. Journal of Clinical Oncology, 2011, 29, 1452-1457. | 0.8 | 197 |
| 62 | Primary CNS Lymphoma of T-Cell Origin: A Descriptive Analysis From the International Primary CNS Lymphoma Collaborative Group. Journal of Clinical Oncology, 2005, 23, 2233-2239. | 0.8 | 188 |
| 63 | Tumor-associated macrophages predict inferior outcomes in classic Hodgkin lymphoma: a correlative study from the E2496 Intergroup trial. Blood, 2012, 120, 3280-3287. | 0.6 | 188 |
| 64 | Histological Transformation and Progression in Follicular Lymphoma: A Clonal Evolution Study. PLoS Medicine, 2016, 13, e1002197. | 3.9 | 185 |
| 65 | Recurrent somatic mutations of PTPN1 in primary mediastinal B cell lymphoma and Hodgkin lymphoma. Nature Genetics, 2014, 46, 329-335. | 9.4 | 180 |
| 66 | Transformation to Aggressive Lymphoma in Nodular Lymphocyte-Predominant Hodgkin's Lymphoma. Journal of Clinical Oncology, 2010, 28, 793-799. | 0.8 | 178 |
| 67 | Clinicogenetic risk models predict early progression of follicular lymphoma after first-line immunochemotherapy. Blood, 2016, 128, 1112-1120. | 0.6 | 177 |
| 68 | Brief Chemotherapy and Involved Field Radiation Therapy for Limited-Stage, Histologically Aggressive Lymphoma. Annals of Internal Medicine, 1987, 107, 25. | 2.0 | 176 |
| 69 | Gene Expression–Based Model Using Formalin-Fixed Paraffin-Embedded Biopsies Predicts Overall Survival in Advanced-Stage Classical Hodgkin Lymphoma. Journal of Clinical Oncology, 2013, 31, 692-700. | 0.8 | 176 |
| 70 | Five-year results of brentuximab vedotin in patients with relapsed or refractory systemic anaplastic large cell lymphoma. Blood, 2017, 130, 2709-2717. | 0.6 | 176 |
| 71 | The architectural pattern of FOXP3-positive T cells in follicular lymphoma is an independent predictor of survival and histologic transformation. Blood, 2010, 115, 289-295. | 0.6 | 173 |
| 72 | International Prognostic Score in Advanced-Stage Hodgkin's Lymphoma: Altered Utility in the Modern Era. Journal of Clinical Oncology, 2012, 30, 3383-3388. | 0.8 | 171 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Small Noncleaved, Non-Burkitt's (Burkitt-Like) Lymphoma: Cytogenetics Predict Outcome and Reflect Clinical Presentation. Journal of Clinical Oncology, 1999, 17, 1558-1558. | 0.8 | 169 |
| 74 | High-grade B-cell lymphoma with MYC and BCL2 and/or BCL6 rearrangements with diffuse large B-cell lymphoma morphology. Blood, 2018, 131, 2060-2064. | 0.6 | 167 |
| 75 | Acquired <i>TNFRSF14</i> Mutations in Follicular Lymphoma Are Associated with Worse Prognosis. Cancer Research, 2010, 70, 9166-9174. | 0.4 | 160 |
| 76 | LMO2 Protein Expression Predicts Survival in Patients With Diffuse Large B-Cell Lymphoma Treated With Anthracycline-Based Chemotherapy With and Without Rituximab. Journal of Clinical Oncology, 2008, 26, 447-454. | 0.8 | 159 |
| 77 | Genome-wide copy number analysis of Hodgkin Reed-Sternberg cells identifies recurrent imbalances with correlations to treatment outcome. Blood, 2010, 116, 418-427. | 0.6 | 152 |
| 78 | BCL2 Predicts Survival in Germinal Center B-cell–like Diffuse Large B-cell Lymphoma Treated with CHOP-like Therapy and Rituximab. Clinical Cancer Research, 2011, 17, 7785-7795. | 3.2 | 152 |
| 79 | Analysis of Heritability and Shared Heritability Based on Genome-Wide Association Studies for Thirteen Cancer Types. Journal of the National Cancer Institute, 2015, 107, djv279. | 3.0 | 152 |
| 80 | Impact of dual expression of MYC and BCL2 by immunohistochemistry on the risk of CNS relapse in DLBCL. Blood, 2016, 127, 2182-2188. | 0.6 | 145 |
| 81 | Analysis of secondary chromosomal alterations in 165 cases of follicular lymphoma with t(14;18). Genes Chromosomes and Cancer, 2001, 30, 375-382. | 1.5 | 142 |
| 82 | Analysis of FOXO1 mutations in diffuse large B-cell lymphoma. Blood, 2013, 121, 3666-3674. | 0.6 | 139 |
| 83 | Prognostic Factors in Follicular Lymphoma. Journal of Clinical Oncology, 2010, 28, 2902-2913. | 0.8 | 136 |
| 84 | Diffuse large B-cell lymphoma: reduced CD20 expression is associated with an inferior survival. Blood, 2009, 113, 3773-3780. | 0.6 | 133 |
| 85 | Gene expression predicts overall survival in paraffin-embedded tissues of diffuse large B-cell lymphoma treated with R-CHOP. Blood, 2008, 112, 3425-3433. | 0.6 | 130 |
| 86 | Essential Role of the Linear Ubiquitin Chain Assembly Complex in Lymphoma Revealed by Rare Germline Polymorphisms. Cancer Discovery, 2014, 4, 480-493. | 7.7 | 130 |
| 87 | Prognostic Significance of Anaplastic Lymphoma Kinase (ALK) Protein Expression in Adults With Anaplastic Large Cell Lymphoma. Blood, 1999, 93, 3913-3921. | 0.6 | 130 |
| 88 | Expression of the FOXP1 transcription factor is strongly associated with inferior survival in patients with diffuse large B-cell lymphoma. Clinical Cancer Research, 2005, 11, 1065-72. | 3.2 | 130 |
| 89 | Recurrent targets of aberrant somatic hypermutation in lymphoma. Oncotarget, 2012, 3, 1308-1319. | 0.8 | 127 |
| 90 | Involved-Nodal Radiation Therapy As a Component of Combination Therapy for Limited-Stage Hodgkin's Lymphoma: A Question of Field Size. Journal of Clinical Oncology, 2008, 26, 5170-5174. | 0.8 | 126 |

| # | Article | IF | CITATIONS |
|-----|--|------|-----------|
| 91 | The E3 ubiquitin ligase UBR5 is recurrently mutated in mantle cell lymphoma. Blood, 2013, 121, 3161-3164. | 0.6 | 124 |
| 92 | Genome-wide profiling of follicular lymphoma by array comparative genomic hybridization reveals prognostically significant DNA copy number imbalances. Blood, 2009, 113, 137-148. | 0.6 | 122 |
| 93 | Gene expression profiling of microdissected Hodgkin Reed-Sternberg cells correlates with treatment outcome in classical Hodgkin lymphoma. Blood, 2012, 120, 3530-3540. | 0.6 | 122 |
| 94 | Organochlorines and risk of nonâ€Hodgkin lymphoma. International Journal of Cancer, 2007, 121, 2767-2775. | 2.3 | 121 |
| 95 | Follicular lymphoma lacking the t(14;18)(q32;q21): identification of two disease subtypes. British Journal of Haematology, 2003, 120, 424-433. | 1.2 | 118 |
| 96 | Identification of cytogenetic subgroups and karyotypic pathways of clonal evolution in follicular lymphomas. Genes Chromosomes and Cancer, 2004, 39, 195-204. | 1.5 | 114 |
| 97 | High-dose chemotherapy and autologous stem cell transplantation for primary refractory or relapsed Hodgkin lymphoma: long-term outcome in the first 100 patients treated in Vancouver. Blood, 2005, 106, 1473-1478. | 0.6 | 112 |
| 98 | Genetic profiling of MYC and BCL2 in diffuse large B-cell lymphoma determines cell-of-origin–specific clinical impact. Blood, 2017, 129, 2760-2770. | 0.6 | 112 |
| 99 | Treatment of post-transplant lymphoproliferative disease with rituximab monoclonal antibody after lung transplantation. Lancet, The, 1999, 354, 1698-1699. | 6.3 | 111 |
| 100 | The efficacy and tolerability of adriamycin, bleomycin, vinblastine, dacarbazine and <scp>S</scp> tanford <scp>V</scp> in older <scp>H</scp> odgkin lymphoma patients: a comprehensive analysis from the <scp>N</scp> orth <scp>A</scp> merican intergroup trial <scp>E</scp> 2496. British lournal of Haematology, 2013, 161, 76-86. | 1.2 | 111 |
| 101 | Genome-wide copy-number analyses reveal genomic abnormalities involved in transformation of follicular lymphoma. Blood, 2014, 123, 1681-1690. | 0.6 | 110 |
| 102 | Comprehensive miRNA sequence analysis reveals survival differences in diffuse large B-cell lymphoma patients. Genome Biology, 2015, 16, 18. | 3.8 | 107 |
| 103 | Hodgkin lymphoma. Nature Reviews Disease Primers, 2020, 6, 61. | 18.1 | 103 |
| 104 | New Molecular Assay for the Proliferation Signature in Mantle Cell Lymphoma Applicable to Formalin-Fixed Paraffin-Embedded Biopsies. Journal of Clinical Oncology, 2017, 35, 1668-1677. | 0.8 | 102 |
| 105 | Genome-wide discovery of somatic regulatory variants in diffuse large B-cell lymphoma. Nature Communications, 2018, 9, 4001. | 5.8 | 102 |
| 106 | The Prognostic Impact of CD163-Positive Macrophages in Follicular Lymphoma: A Study from the BC Cancer Agency and the Lymphoma Study Association. Clinical Cancer Research, 2015, 21, 3428-3435. | 3.2 | 101 |
| 107 | High microvessel density determines a poor outcome in patients with diffuse large B-cell lymphoma treated with rituximab plus chemotherapy. Haematologica, 2011, 96, 996-1001. | 1.7 | 100 |
| 108 | Rapid infusion rituximab in combination with corticosteroid-containing chemotherapy or as maintenance therapy is well tolerated and can safely be delivered in the community setting. Blood, 2007, 109, 4171-4173. | 0.6 | 98 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Integrative genomic analysis identifies key pathogenic mechanisms in primary mediastinal large B-cell lymphoma. Blood, 2019, 134, 802-813. | 0.6 | 96 |
| 110 | Longâ€ŧerm outcomes for patients with limited stage follicular lymphoma. Cancer, 2010, 116, 3797-3806. | 2.0 | 94 |
| 111 | Meta-analysis of genome-wide association studies discovers multiple loci for chronic lymphocytic leukemia. Nature Communications, 2016, 7, 10933. | 5.8 | 94 |
| 112 | Paraffin-based 6-gene model predicts outcome in diffuse large B-cell lymphoma patients treated with R-CHOP. Blood, 2008, 111, 5509-5514. | 0.6 | 93 |
| 113 | Autologous and Allogeneic Stem-Cell Transplantation for Transformed Follicular Lymphoma: A Report of the Canadian Blood and Marrow Transplant Group. Journal of Clinical Oncology, 2013, 31, 1164-1171. | 0.8 | 92 |
| 114 | Cell of origin of transformed follicular lymphoma. Blood, 2015, 126, 2118-2127. | 0.6 | 91 |
| 115 | Treatment strategies, outcomes and prognostic factors in 291 patients with secondary CNS involvement by diffuse large B-cell lymphoma. European Journal of Cancer, 2018, 93, 57-68. | 1.3 | 90 |
| 116 | Mantle cell lymphoma in the ocular adnexal region. Ophthalmology, 2005, 112, 114-119. | 2.5 | 89 |
| 117 | Brentuximab vedotin plus bendamustine in relapsed or refractory Hodgkin's lymphoma: an international, multicentre, single-arm, phase 1–2 trial. Lancet Oncology, The, 2018, 19, 257-266. | 5.1 | 89 |
| 118 | A gene signature that distinguishes conventional and leukemic nonnodal mantle cell lymphoma helps predict outcome. Blood, 2018, 132, 413-422. | 0.6 | 89 |
| 119 | Primary Adrenal Lymphoma. Clinical Lymphoma and Myeloma, 2003, 4, 154-160. | 2.1 | 87 |
| 120 | Brentuximab vedotin with chemotherapy for stage III/IV classical Hodgkin lymphoma: 3-year update of the ECHELON-1 study. Blood, 2020, 135, 735-742. | 0.6 | 86 |
| 121 | Treating limited-stage nodular lymphocyte predominant Hodgkin lymphoma similarly to classical Hodgkin lymphoma with ABVD may improve outcome. Blood, 2011, 118, 4585-4590. | 0.6 | 83 |
| 122 | Brentuximab vedotin with chemotherapy for stage III or IV classical Hodgkin lymphoma (ECHELON-1): 5-year update of an international, open-label, randomised, phase 3 trial. Lancet Haematology,the, 2021, 8, e410-e421. | 2.2 | 83 |
| 123 | Allogeneic haematopoietic stem-cell transplantation for relapsed and refractory aggressive histology non-Hodgkin lymphoma*. British Journal of Haematology, 2005, 131, 223-230. | 1.2 | 78 |
| 124 | Diffuse large Bâ€cell lymphoma with testicular involvement: outcome and risk of <scp>CNS</scp> relapse in the rituximab era. British Journal of Haematology, 2017, 176, 210-221. | 1.2 | 78 |
| 125 | Early progression after bendamustine-rituximab is associated with high risk of transformation in advanced stage follicular lymphoma. Blood, 2019, 134, 761-764. | 0.6 | 77 |
| 126 | Advanced-stage nodular lymphocyte predominant Hodgkin lymphoma compared with classical Hodgkin lymphoma: a matched pair outcome analysis. Blood, 2014, 123, 3567-3573. | 0.6 | 76 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 127 | Active Idiotypic Vaccination Versus Control Immunotherapy for Follicular Lymphoma. Journal of Clinical Oncology, 2014, 32, 1797-1803. | 0.8 | 75 |
| 128 | MACOP-B and VACOP-B in diffuse large cell lymphomas and MOPP/ABV in Hodgkin's disease. Annals of Oncology, 1991, 2, 17-23. | 0.6 | 74 |
| 129 | High-resolution architecture and partner genes of MYC rearrangements in lymphoma with DLBCL morphology. Blood Advances, 2018, 2, 2755-2765. | 2.5 | 74 |
| 130 | Outcome prediction by extranodal involvement, IPI, Râ€IPI, and NCCNâ€IPI in the PET/CT and rituximab era: A <scp>D</scp> anish– <scp>C</scp> anadian study of 443 patients with diffuseâ€large <scp>B</scp> â€cell lymphoma. American Journal of Hematology, 2015, 90, 1041-1046. | 2.0 | 71 |
| 131 | Diffuse large B-cell lymphoma with involvement of the kidney: outcome and risk of central nervous system relapse. Haematologica, 2011, 96, 1002-1007. | 1.7 | 69 |
| 132 | Prognostic Significance of Bax Protein Expression in Diffuse Aggressive Non-Hodgkin's Lymphoma. Blood, 1997, 90, 3173-3178. | 0.6 | 65 |
| 133 | The Spectrum of Lymphoma with 8q24 Aberrations: A Clinical, Pathological and Cytogenetic Study of 87 Consecutive Cases. Leukemia and Lymphoma, 2004, 45, 519-528. | 0.6 | 65 |
| 134 | Treatment and Outcomes in Patients With Primary Cutaneous B-Cell Lymphoma: The BC Cancer Agency Experience. International Journal of Radiation Oncology Biology Physics, 2013, 87, 719-725. | 0.4 | 65 |
| 135 | The number of extranodal sites assessed by PET/CT scan is a powerful predictor of CNS relapse for patients with diffuse large B-cell lymphoma: An international multicenter study of 1532 patients treated with chemoimmunotherapy. European Journal of Cancer, 2017, 75, 195-203. | 1.3 | 65 |
| 136 | Molecular classification of primary mediastinal large B-cell lymphoma using routinely available tissue specimens. Blood, 2018, 132, 2401-2405. | 0.6 | 64 |
| 137 | CHOP-R therapy overcomes the adverse prognostic influence of BCL-2 expression in diffuse large B-cell lymphoma. Leukemia and Lymphoma, 2007, 48, 1102-1109. | 0.6 | 63 |
| 138 | Correlations between BCL6 rearrangement and outcome in patients with diffuse large B-cell lymphoma treated with CHOP or R-CHOP. Haematologica, 2010, 95, 96-101. | 1.7 | 63 |
| 139 | Identification of Primary Mediastinal Large B-cell Lymphoma at Nonmediastinal Sites by Gene Expression Profiling. American Journal of Surgical Pathology, 2015, 39, 1322-1330. | 2.1 | 63 |
| 140 | High-Dose Therapy and Autologous Hematopoietic Stem-Cell Transplantation Does Not Increase the Risk of Second Neoplasms for Patients With Hodgkin's Lymphoma: A Comparison of Conventional Therapy Alone Versus Conventional Therapy Followed by Autologous Hematopoietic Stem-Cell Transplantation. Journal of Clinical Oncology, 2005, 23, 7994-8002. | 0.8 | 62 |
| 141 | TBL1XR1/TP63: a novel recurrent gene fusion in B-cell non-Hodgkin lymphoma. Blood, 2012, 119, 4949-4952. | 0.6 | 60 |
| 142 | Reduced telomere length variation in healthy oldest old. Mechanisms of Ageing and Development, 2008, 129, 638-641. | 2.2 | 59 |
| 143 | Outcomes in splenic marginal zone lymphoma: analysis of 107 patients treated in British Columbia. British Journal of Haematology, 2015, 169, 520-527. | 1.2 | 58 |
| 144 | Assessment of Capture and Amplicon-Based Approaches for the Development of a Targeted Next-Generation Sequencing Pipeline to Personalize Lymphoma Management. Journal of Molecular Diagnostics, 2018, 20, 203-214. | 1.2 | 58 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 145 | Long-term results of PET-guided radiation in patients with advanced-stage diffuse large B-cell lymphoma treated with R-CHOP. Blood, 2021, 137, 929-938. | 0.6 | 57 |
| 146 | Evaluation of the Risk of Relapse in Classical Hodgkin Lymphoma at Event-Free Survival Time Points and Survival Comparison With the General Population in British Columbia. Journal of Clinical Oncology, 2016, 34, 2493-2500. | 0.8 | 56 |
| 147 | Identification of highâ€risk <i><scp>DUSP</scp>22</i> â€rearranged <scp>ALK</scp> â€negative anaplastic large cell lymphoma. British Journal of Haematology, 2019, 186, e28-e31. | 1.2 | 56 |
| 148 | Primary paranasal sinus lymphoma: natural history and improved outcome with central nervous system chemoprophylaxis. Leukemia and Lymphoma, 2005, 46, 1721-1727. | 0.6 | 55 |
| 149 | Hodgkin's Lymphoma in Adolescents. Journal of Clinical Oncology, 2006, 24, 2520-2526. | 0.8 | 55 |
| 150 | Cytogenetic Analysis in Mantle Cell Lymphoma: A Review of 214 Cases. Leukemia and Lymphoma, 2002, 43, 783-791. | 0.6 | 54 |
| 151 | CD20 mutations involving the rituximab epitope are rare in diffuse large B-cell lymphomas and are not a significant cause of R-CHOP failure. Haematologica, 2009, 94, 423-427. | 1.7 | 53 |
| 152 | Prognostic implications of extranodal involvement in patients with diffuse large B-cell lymphoma treated with rituximab and cyclophosphamide, doxorubicin, vincristine, and prednisone. Leukemia and Lymphoma, 2010, 51, 1-10. | 0.6 | 53 |
| 153 | A new biologic prognostic model based on immunohistochemistry predicts survival in patients with diffuse large B-cell lymphoma. Blood, 2012, 120, 2290-2296. | 0.6 | 53 |
| 154 | High resolution analysis of follicular lymphoma genomes reveals somatic recurrent sites of copyâ€neutral loss of heterozygosity and copy number alterations that target single genes. Genes Chromosomes and Cancer, 2010, 49, 669-681. | 1.5 | 51 |
| 155 | GBV /hepatitis G virus infection and nonâ€Hodgkin lymphoma: a case control study. International Journal of Cancer, 2010, 126, 2885-2892. | 2.3 | 51 |
| 156 | Bone Marrow Involvement in T-Cell–Rich B-Cell Lymphoma. American Journal of Clinical Pathology, 1997, 108, 570-578. | 0.4 | 50 |
| 157 | Prognostic significance of secondary cytogenetic alterations in follicular lymphomas. Genes Chromosomes and Cancer, 2008, 47, 1038-1048. | 1.5 | 50 |
| 158 | International Assessment of Event-Free Survival at 24 Months and Subsequent Survival in Peripheral T-Cell Lymphoma. Journal of Clinical Oncology, 2017, 35, 4019-4026. | 0.8 | 50 |
| 159 | Limitedâ€stage diffuse large Bâ€cell lymphoma treated with abbreviated systemic therapy and consolidation radiotherapy. Cancer, 2012, 118, 4156-4165. | 2.0 | 49 |
| 160 | The treatment of primary central nervous system lymphoma in 122 immunocompetent patients. Cancer, 2005, 103, 1008-1017. | 2.0 | 48 |
| 161 | Prognostic Model to Predict Post-Autologous Stem-Cell Transplantation Outcomes in Classical Hodgkin Lymphoma. Journal of Clinical Oncology, 2017, 35, 3722-3733. | 0.8 | 48 |
| 162 | Outcomes in adolescents and young adults with Hodgkin lymphoma treated on US cooperative group protocols: An adult intergroup (E2496) and Children's Oncology Group (COG AHOD0031) comparative analysis. Cancer, 2018, 124, 136-144. | 2.0 | 47 |

| # | Article | IF | CITATIONS |
|-----|--|------|-----------|
| 163 | Long-term disease-free survival of patients with advanced follicular lymphoma after allogeneic bone marrow transplantation. British Journal of Haematology, 2004, 127, 311-321. | 1.2 | 46 |
| 164 | Hand-assisted laparoscopic splenectomy versus open splenectomy for massive splenomegaly: 20-year experience at a Canadian centre. Canadian Journal of Surgery, 2011, 54, 189-193. | 0.5 | 46 |
| 165 | Treatment and Outcomes of Primary Breast Lymphoma. Clinical Breast Cancer, 2012, 12, 412-419. | 1.1 | 46 |
| 166 | Comparison of American Joint Committee on Cancer TNM-based Staging System (7th edition) and Ann Arbor Classification for Predicting Outcome in Ocular Adnexal Lymphoma. Orbit, 2014, 33, 23-28. | 0.5 | 46 |
| 167 | TMEM30A loss-of-function mutations drive lymphomagenesis and confer therapeutically exploitable vulnerability in B-cell lymphoma. Nature Medicine, 2020, 26, 577-588. | 15.2 | 46 |
| 168 | Novel agents improve survival of transplant patients with multiple myeloma including those with high-risk disease defined by early relapse (<12 months). Leukemia and Lymphoma, 2011, 52, 34-41. | 0.6 | 45 |
| 169 | Vascularization predicts overall survival and risk of transformation in follicular lymphoma. Haematologica, 2010, 95, 2157-2160. | 1.7 | 44 |
| 170 | HLA-DR protein status predicts survival in patients with diffuse large B-cell lymphoma treated on the MACOP-B chemotherapy regimen. Leukemia and Lymphoma, 2007, 48, 542-546. | 0.6 | 43 |
| 171 | Hodgkin Lymphoma in Pregnancy. Current Hematologic Malignancy Reports, 2013, 8, 211-217. | 1.2 | 43 |
| 172 | Genetic Variation in Healthy Oldest-Old. PLoS ONE, 2009, 4, e6641. | 1.1 | 42 |
| 173 | Clinical Manifestations and Natural History of Hodgkin's Lymphoma. Cancer Journal (Sudbury, Mass), 2009, 15, 124-128. | 1.0 | 42 |
| 174 | Rituximab with highâ€dose methotrexate in primary central nervous system lymphoma. American Journal of Hematology, 2015, 90, 1149-1154. | 2.0 | 41 |
| 175 | COO and MYC/BCL2 status do not predict outcome among patients with stage I/II DLBCL: a retrospective multicenter study. Blood Advances, 2019, 3, 2013-2021. | 2.5 | 40 |
| 176 | Multicolour fluorescence in situ hybridization analysis of t(14;18)-positive follicular lymphoma and correlation with gene expression data and clinical outcome. British Journal of Haematology, 2003, 122, 745-759. | 1.2 | 39 |
| 177 | Outcome of Patients With Non-Hodgkin Lymphomas With Concurrent MYC and BCL2 Rearrangements Treated With CODOX-M/IVAC With Rituximab Followed by Hematopoietic Stem Cell Transplantation. Clinical Lymphoma, Myeloma and Leukemia, 2015, 15, 341-348. | 0.2 | 39 |
| 178 | Risk assessment in the management of newly diagnosed classical Hodgkin lymphoma. Blood, 2015, 125, 1693-1702. | 0.6 | 38 |
| 179 | Anaplastic large cell lymphoma: a clinicopathologic analysis. , 1999, 17, 137-148. | | 37 |
| 180 | Genetic Variation in H2AFX Contributes to Risk of Non–Hodgkin Lymphoma. Cancer Epidemiology Biomarkers and Prevention, 2007, 16, 1098-1106. | 1.1 | 37 |

| # | Article | IF | CITATIONS |
|-----|--|------|-----------|
| 181 | Five-year follow-up of brentuximab vedotin combined with ABVD or AVD for advanced-stage classical Hodgkin lymphoma. Blood, 2017, 130, 1375-1377. | 0.6 | 37 |
| 182 | Secondary Breast Cancer Risk by Radiation Volume in Women With Hodgkin Lymphoma. International Journal of Radiation Oncology Biology Physics, 2017, 97, 35-41. | 0.4 | 37 |
| 183 | Treatment of Elderly Hodgkin's Lymphoma Patients with a Novel 5-drug Regimen (ODBEP): A Phase II Study. Leukemia and Lymphoma, 2002, 43, 1395-1402. | 0.6 | 36 |
| 184 | Frequent occurrence of deletions in primary mediastinal Bâ \in cell lymphoma. Genes Chromosomes and Cancer, 2007, 46, 1090-1097. | 1.5 | 36 |
| 185 | Interaction between organochlorines and the AHR gene, and risk of non-Hodgkin lymphoma. Cancer Causes and Control, 2010, 21, 11-22. | 0.8 | 36 |
| 186 | Automated Analysis of Multidimensional Flow Cytometry Data Improves Diagnostic Accuracy Between Mantle Cell Lymphoma and Small Lymphocytic Lymphoma. American Journal of Clinical Pathology, 2012, 137, 75-85. | 0.4 | 36 |
| 187 | Primary, Unilateral Ocular Adnexal Lymphoma: Disease Progression and Long-Term Survival. Ophthalmic Plastic and Reconstructive Surgery, 2011, 27, 405-409. | 0.4 | 35 |
| 188 | Fludarabine and rituximab for relapsed or refractory hairy cell leukemia. Blood, 2012, 119, 1988-1991. | 0.6 | 35 |
| 189 | Hodgkin's Lymphoma — The Great Teacher. New England Journal of Medicine, 2011, 365, 264-265. | 13.9 | 34 |
| 190 | Prognostic factors for advancedâ€stage human immunodeficiency virusâ€associated classical Hodgkin lymphoma treated with doxorubicin, bleomycin, vinblastine, and dacarbazine plus combined antiretroviral therapy: A multiâ€institutional retrospective study. Cancer, 2015, 121, 423-431. | 2.0 | 34 |
| 191 | Uterine, but not ovarian, female reproductive organ involvement at presentation by diffuse large Bâ€cell lymphoma is associated with poor outcomes and a high frequency of secondary <scp>CNS</scp> involvement. British Journal of Haematology, 2016, 175, 876-883. | 1.2 | 34 |
| 192 | Splenectomy vs. alpha interferon: A randomized study in patients with previously untreated hairy cell leukemia. American Journal of Hematology, 1992, 41, 13-18. | 2.0 | 33 |
| 193 | Radioimmunotherapy — Hot New Treatment for Lymphoma. New England Journal of Medicine, 2005, 352, 496-498. | 13.9 | 33 |
| 194 | Gemcitabine, dexamethasone, and cisplatin (GDP) is an effective and well-tolerated salvage therapy for relapsed/refractory diffuse large B-cell lymphoma and Hodgkin lymphoma. Leukemia and Lymphoma, 2017, 58, 324-332. | 0.6 | 32 |
| 195 | Hodgkin Lymphoma: Current Status and Clinical Trial Recommendations. Journal of the National Cancer Institute, 2017, 109, djw249. | 3.0 | 31 |
| 196 | FOXP1 expression is a prognostic biomarker in follicular lymphoma treated with rituximab and chemotherapy. Blood, 2018, 131, 226-235. | 0.6 | 31 |
| 197 | Diffuse Malignant Lymphoma with Cerebriform Nuclei: A B-Cell Lymphoma Studied with Monoclonal Antibodies. American Journal of Clinical Pathology, 1985, 83, 753-759. | 0.4 | 30 |
| 198 | Characterization of the recurrent translocation t(1;1)(p36.3;q21.1-2) in non-Hodgkin lymphoma by multicolor banding and fluorescence in situ hybridization analysis. Genes Chromosomes and Cancer, 2003, 36, 375-381. | 1.5 | 30 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 199 | Salvage Therapy with Allogeneic Stem Cell Transplantation Results in Better Outcome for Patients with Relapsed/Refractory Follicular Lymphoma Compared to Those with Transformed Non-Hodgkin Lymphoma: A Population-Based Comparative Study Blood, 2008, 112, 975-975. | 0.6 | 29 |
| 200 | Results of a Pivotal Phase 2 Study of Brentuximab Vedotin (SGN-35) in Patients with Relapsed or Refractory Hodgkin Lymphoma. Blood, 2010, 116, 283-283. | 0.6 | 29 |
| 201 | The Outcome of Primary Mediastinal Large B-Cell Lymphoma (PMBCL) in the R-CHOP Treatment Era. Blood, 2012, 120, 303-303. | 0.6 | 29 |
| 202 | Genomic testing to determine drug response: measuring preferences of the public and patients using Discrete Choice Experiment (DCE). BMC Health Services Research, 2013, 13, 454. | 0.9 | 28 |
| 203 | R-CHOP with Etoposide Substituted for Doxorubicin (R-CEOP): Excellent Outcome in Diffuse Large B Cell Lymphoma for Patients with a Contraindication to Anthracyclines Blood, 2009, 114, 408-408. | 0.6 | 28 |
| 204 | Haematopoietic stem cell transplantation as primary therapy of sporadic adult Burkitt lymphoma*. British Journal of Haematology, 2006, 133, 634-637. | 1.2 | 27 |
| 205 | Hepatitis C virus and risk of nonâ€Hodgkin lymphoma in British Columbia, Canada. International Journal of Cancer, 2008, 122, 630-633. | 2.3 | 27 |
| 206 | Genetic variation in the NBS1, MRE11, RAD50 and BLM genes and susceptibility to non-Hodgkin lymphoma. BMC Medical Genetics, 2009, 10, 117. | 2.1 | 27 |
| 207 | Validation of a Prognostic Model to Assess the Risk of CNS Disease in Patients with Aggressive B-Cell Lymphoma. Blood, 2014, 124, 394-394. | 0.6 | 27 |
| 208 | Delineation of a minimal region of deletion at 6q16.3 in follicular lymphoma and construction of a bacterial artificial chromosome contig spanning a 6-megabase region of 6q16-q21. Genes Chromosomes and Cancer, 2004, 40, 60-65. | 1.5 | 26 |
| 209 | Effect of Place of Residence and Treatment on Survival Outcomes in Patients With Diffuse Large B-Cell Lymphoma in British Columbia. Oncologist, 2014, 19, 283-290. | 1.9 | 26 |
| 210 | Toward Personalized Lymphoma Immunotherapy: Identification of Common Driver Mutations Recognized by Patient CD8+ T Cells. Clinical Cancer Research, 2016, 22, 2226-2236. | 3.2 | 26 |
| 211 | Site of central nervous system (CNS) relapse in patients with diffuse large B ell lymphoma (DLBCL) by the CNSâ€IPI risk model. British Journal of Haematology, 2017, 179, 508-510. | 1.2 | 26 |
| 212 | Brentuximab Vedotin plus Chemotherapy in North American Subjects with Newly Diagnosed Stage III or IV Hodgkin Lymphoma. Clinical Cancer Research, 2019, 25, 1718-1726. | 3.2 | 26 |
| 213 | Cost-Effectiveness of the Addition of Rituximab to CHOP Chemotherapy in First-Line Treatment for Diffuse Large B-Cell Lymphoma in a Population-Based Observational Cohort in British Columbia, Canada. Value in Health, 2010, 13, 703-711. | 0.1 | 25 |
| 214 | Oral fludarabine and rituximab as initial therapy for chronic lymphocytic leukemia or small lymphocytic lymphoma: population-based experience matches clinical trials. Leukemia and Lymphoma, 2012, 53, 77-82. | 0.6 | 24 |
| 215 | An RCOR1 loss–associated gene expression signature identifies a prognostically significant DLBCL subgroup. Blood, 2015, 125, 959-966. | 0.6 | 24 |
| 216 | Three-Year Survival Results From An Ongoing Phase 2 Study Of Brentuximab Vedotin In Patients With Relapsed Or Refractory Systemic Anaplastic Large Cell Lymphoma. Blood, 2013, 122, 1809-1809. | 0.6 | 24 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 217 | The evolving role of etoposide in the management of lymphomas and hodgkin's disease. Cancer, 1991, 67, 271-280. | 2.0 | 23 |
| 218 | Clinical Significance of the t(14;18) AndBCL2Overexpression in Follicular Large Cell Lymphoma. Leukemia and Lymphoma, 2000, 36, 513-523. | 0.6 | 23 |
| 219 | Challenging Problems: Coincident Pregnancy, HIV Infection, and Older Age. Hematology American Society of Hematology Education Program, 2008, 2008, 334-339. | 0.9 | 23 |
| 220 | Allogeneic haematopoietic stem cell transplantation for chronic lymphocytic leukaemia: outcome in a 20â€year cohort. British Journal of Haematology, 2012, 158, 174-185. | 1.2 | 23 |
| 221 | Mapping the human T cell repertoire to recurrent driver mutations in MYD88 and EZH2 in lymphoma. Oncolmmunology, 2017, 6, e1321184. | 2.1 | 23 |
| 222 | Phase 2 Trial of Interim PET Scan-Tailored Therapy in Patients with Advanced Stage Diffuse Large B-Cell Lymphoma (DLBCL) in British Columbia (BC). Blood, 2014, 124, 392-392. | 0.6 | 23 |
| 223 | Mantle cell lymphoma: report of the 2010 Mantle Cell Lymphoma Consortium Workshop. Leukemia and Lymphoma, 2011, 52, 24-33. | 0.6 | 22 |
| 224 | Positron Emission Tomography in the Management of Hodgkin Lymphoma. Hematology American Society of Hematology Education Program, 2011, 2011, 317-322. | 0.9 | 22 |
| 225 | Molecular features of a large cohort of primary central nervous system lymphoma using tissue microarray. Blood Advances, 2019, 3, 3953-3961. | 2.5 | 22 |
| 226 | Uncovering novel inter- and intrachromosomal chromosome 1 aberrations in follicular lymphomas by using an innovative multicolor banding technique. Genes Chromosomes and Cancer, 2002, 34, 201-210. | 1.5 | 21 |
| 227 | Physical activity, obesity and survival in diffuse large Bâ€cell and follicular lymphoma cases. British Journal of Haematology, 2017, 178, 442-447. | 1.2 | 21 |
| 228 | Genetic Variation in Cell Death Genes and Risk of Non-Hodgkin Lymphoma. PLoS ONE, 2012, 7, e31560. | 1.1 | 21 |
| 229 | Outcome of primary cutaneous anaplastic large cell lymphoma: a 20â€year British Columbia Cancer Agency experience. British Journal of Haematology, 2017, 176, 234-240. | 1.2 | 20 |
| 230 | Advanced Stage Classical Hodgkin Lymphoma Patients with a Negative PET-Scan Following Treatment with ABVD Have Excellent Outcomes without the Need for Consolidative Radiotherapy Regardless of Disease Bulk at Presentation. Blood, 2015, 126, 579-579. | 0.6 | 20 |
| 231 | Three-Year Follow-Up Data and Characterization Of Long-Term Remissions From An Ongoing Phase 2 Study Of Brentuximab Vedotin In Patients With Relapsed Or Refractory Hodgkin Lymphoma. Blood, 2013, 122, 4382-4382. | 0.6 | 19 |
| 232 | Impact of MYC and BCL2 structural variants in tumors of DLBCL morphology and mechanisms of false-negative MYC IHC. Blood, 2021, 137, 2196-2208. | 0.6 | 18 |
| 233 | Development of adapted RECIST criteria to assess response in lymphoma and their comparison to the International Workshop Criteria. Leukemia and Lymphoma, 2007, 48, 513-520. | 0.6 | 17 |
| 234 | Elevated circulating t(14;18) translocation levels prior to diagnosis of follicular lymphoma. Blood, 2010, 116, 6146-6147. | 0.6 | 17 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 235 | A Canadian Evidence-Based Guideline for the First-Line Treatment of Follicular Lymphoma: Joint Consensus of the Lymphoma Canada Scientific Advisory Board. Clinical Lymphoma, Myeloma and Leukemia, 2015, 15, 59-74. | 0.2 | 17 |
| 236 | Long-term outcomes of R-CEOP show curative potential in patients with DLBCL and a contraindication to anthracyclines. Blood Advances, 2021, 5, 1483-1489. | 2.5 | 17 |
| 237 | Cure of ommaya reservoir associated staphylococcus epidermidis ventriculitis with a simple regimen of vancomycin and rifampin without reservoir removal. Medical and Pediatric Oncology, 1982, 10, 549-552. | 1.0 | 16 |
| 238 | TREATMENT CONSIDERATIONS IN THE ELDERLY PATIENT WITH LYMPHOMA. Hematology/Oncology Clinics of North America, 1997, 11, 949-961. | 0.9 | 16 |
| 239 | Favorable Outcomes from Allogeneic and Autologous Stem Cell Transplantation for Patients with Transformed Nonfollicular Indolent Lymphoma. Biology of Blood and Marrow Transplantation, 2014, 20, 1813-1818. | 2.0 | 16 |
| 240 | Interim PET-directed therapy in limited-stage Hodgkin lymphoma initially treated with ABVD. Haematologica, 2018, 103, e590-e593. | 1.7 | 16 |
| 241 | Brentuximab vedotin for the treatment of patients with relapsed or refractory Hodgkin lymphoma after autologous stem cell transplantation. British Journal of Haematology, 2020, 188, 540-549. | 1.2 | 16 |
| 242 | Limited-Stage Diffuse Large B-Cell Lymphoma (DLBCL) Patients with a Negative Pet Scan Following Three Cycles of R-CHOP Can Be Effectively Treated with Abbreviated Chemoimmunotherapy Alone Blood, 2007, 110, 787-787. | 0.6 | 16 |
| 243 | Gemcitabine, Dexamethasone, and Cisplatin (GDP) Is An Effective and Well-Tolerated out-Patient Salvage Therapy for Relapsed/Refractory Diffuse Large B-Cell Lymphoma (DLBCL) and Hodgkin Lymphoma (HL). Blood, 2010, 116, 113-113. | 0.6 | 16 |
| 244 | Hodgkin's Lymphoma: The Hazards of Success. Journal of Clinical Oncology, 2003, 21, 3388-3390. | 0.8 | 15 |
| 245 | Genomeâ€wide methylation analyses identify a subset of mantle cell lymphoma with a high number of methylated CpGs and aggressive clinicopathological features. International Journal of Cancer, 2013, 133, 2852-2863. | 2.3 | 15 |
| 246 | Clinical characteristics and outcomes of patients with Hodgkin lymphoma with central nervous system involvement: An international multicenter collaboration. American Journal of Hematology, 2016, 91, 894-899. | 2.0 | 15 |
| 247 | The Super-Seniors Study: Phenotypic characterization of a healthy 85+ population. PLoS ONE, 2018, 13, e0197578. | 1.1 | 15 |
| 248 | Excellent realâ€world outcomes of adults with Burkitt lymphoma treated with <scp>CODOX</scp> â€M/ <scp>IVAC</scp> plus or minus rituximab. British Journal of Haematology, 2018, 181, 782-790. | 1.2 | 15 |
| 249 | Comparison of the NCCNâ€IPI, the IPI and PIT scores as prognostic tools in peripheral Tâ€cell lymphomas. British Journal of Haematology, 2019, 186, e24-e27. | 1.2 | 15 |
| 250 | Long-term outcomes for patients with limited-stage follicular lymphoma: update of a population-based study. Blood, 2020, 136, 1006-1010. | 0.6 | 15 |
| 251 | Impact of time from diagnosis to initiation of curative chemotherapy on survival of patients with diffuse large B-cell lymphoma. Leukemia and Lymphoma, 2016, 57, 276-282. | 0.6 | 14 |
| 252 | Treatment patterns and costs of care for patients with relapsed and refractory Hodgkin lymphoma treated with brentuximab vedotin in the United States: A retrospective cohort study. PLoS ONE, 2017, 12, e0180261. | 1.1 | 14 |

| # | Article | IF | CITATIONS |
|-----|--|------|-----------|
| 253 | Addition of Rituximab to CHOP Chemotherapy Significantly Improves Survival of Patients with Transformed Lymphoma Blood, 2007, 110, 790-790. | 0.6 | 14 |
| 254 | Non-Hodgkin lymphoma: the clinician's perspective—a view from the receiving end. Modern Pathology, 2013, 26, S111-S118. | 2.9 | 13 |
| 255 | Diffuse large B-cell lymphoma cell-of-origin classification using the Lymph2Cx assay in the context of BCL2 and MYC expression status. Leukemia and Lymphoma, 2016, 57, 717-720. | 0.6 | 13 |
| 256 | Brentuximab Vedotin for Stage III or IV Hodgkin's Lymphoma. New England Journal of Medicine, 2018, 378, 1558-1561. | 13.9 | 13 |
| 257 | Brentuximab vedotin and bendamustine produce high complete response rates in patients with chemotherapy refractory Hodgkin lymphoma. British Journal of Haematology, 2018, 180, 757-760. | 1.2 | 13 |
| 258 | Genomic predictors of central nervous system relapse in primary testicular diffuse large B-cell lymphoma. Blood, 2021, 137, 1256-1259. | 0.6 | 13 |
| 259 | Gene Expression Differences between Low and High Stage Diffuse Large B Cell Lymphoma (DLBCL) Blood, 2006, 108, 809-809. | 0.6 | 13 |
| 260 | Complete Remissions with Brentuximab Vedotin (SGN-35) in Patients with Relapsed or Refractory Systemic Anaplastic Large Cell Lymphoma. Blood, 2010, 116, 961-961. | 0.6 | 13 |
| 261 | The Combination of Brentuximab Vedotin (Bv) and Bendamustine (B) Demonstrates Marked Activity in Heavily Treated Patients with Relapsed or Refractory Hodgkin Lymphoma (HL) and Anaplastic Large T-Cell Lymphoma (ALCL): Results of an International Multi Center Phase I/II Experience. Blood, 2015, 126, 586-586. | 0.6 | 13 |
| 262 | Alemtuzumab in clinical practice: A British Columbia experience. Leukemia and Lymphoma, 2008, 49, 218-226. | 0.6 | 12 |
| 263 | B Cells With High Side Scatter Parameter by Flow Cytometry Correlate With Inferior Survival in Diffuse Large B-Cell Lymphoma. American Journal of Clinical Pathology, 2012, 137, 805-814. | 0.4 | 12 |
| 264 | Single Cell Phenotypic Profiling of 27 DLBCL Cases Reveals Marked Intertumoral and Intratumoral Heterogeneity. Cytometry Part A: the Journal of the International Society for Analytical Cytology, 2020, 97, 620-629. | 1.1 | 12 |
| 265 | Brentuximab vedotin plus doxorubicin, vinblastine, and dacarbazine in patients with advancedâ€stage, classical Hodgkin lymphoma: A prespecified subgroup analysis of highâ€risk patients from the ECHELONâ€1 study. Hematological Oncology, 2021, 39, 185-195. | 0.8 | 12 |
| 266 | A Novel Prognostic Model Based on Tumor Microenvironment Biology in Relapse Biopsies Predicts Post-Autologous Stem Cell Transplantation Outcomes in Classical Hodgkin Lymphoma. Blood, 2016, 128, 1093-1093. | 0.6 | 12 |
| 267 | Gene Expression Profiling of Microdissected Hodgkin Reed Sternberg Cells: Molecular Subtypes and Treatment Outcome Correlations Blood, 2009, 114, 268-268. | 0.6 | 12 |
| 268 | Tumor-associated antigen PRAME exhibits dualistic functions that are targetable in diffuse large B cell lymphoma. Journal of Clinical Investigation, 2022, 132, . | 3.9 | 12 |
| 269 | A Literature Review of Single Agent Treatment of Multiply Relapsed Aggressive Non-Hodgkin's Lymphoma. Leukemia and Lymphoma, 2002, 43, 975-982. | 0.6 | 11 |
| 270 | Cytogenetic findings in reactive lymphoid hyperplasia: Significance of non-clonal t(3;14) and t(3;22). American Journal of Hematology, 2002, 70, 133-138. | 2.0 | 11 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 271 | Lifetime Physical Activity and the Risk of Non-Hodgkin Lymphoma. Cancer Epidemiology Biomarkers and Prevention, 2015, 24, 873-877. | 1.1 | 11 |
| 272 | Maintenance rituximab following induction R-CHOP chemotherapy in patients with composite or discordant, indolent and aggressive, B-cell non-Hodgkin lymphomas. Haematologica, 2016, 101, e411-e414. | 1.7 | 11 |
| 273 | Gene Expression Signatures Predict Overall Survial in Diffuse Large B Cell Lymphoma Treated with Rituximab and Chop-Like Chemotherapy Blood, 2007, 110, 348-348. | 0.6 | 11 |
| 274 | Brentuximab Vedotin (SGN-35) in Patients with Relapsed or Refractory Systemic Anaplastic Large Cell Lymphoma: A Phase 2 Study Update. Blood, 2011, 118, 443-443. | 0.6 | 11 |
| 275 | Brentuximab Vedotin Combined with ABVD or AVD for Patients with Newly Diagnosed Advanced Stage Hodgkin Lymphoma: Long Term Outcomes. Blood, 2014, 124, 292-292. | 0.6 | 11 |
| 276 | Cost–effectiveness of rituximab in follicular lymphoma. Expert Review of Pharmacoeconomics and Outcomes Research, 2012, 12, 569-577. | 0.7 | 10 |
| 277 | Impact of time from diagnosis to initiation of curative-intent chemotherapy on clinical outcomes in patients with classical Hodgkin lymphoma. Leukemia and Lymphoma, 2016, 57, 872-879. | 0.6 | 10 |
| 278 | High-dose Benda-EAM versus BEAM in patients with relapsed/refractory classical Hodgkin lymphoma undergoing autologous stem cell transplantation. Bone Marrow Transplantation, 2019, 54, 481-484. | 1.3 | 10 |
| 279 | Hodgkin lymphoma: outsmarting HRS cells. Blood, 2020, 136, 2362-2364. | 0.6 | 10 |
| 280 | Older Patients (pts) with Previously Untreated Classical Hodgkin Lymphoma (cHL): A Detailed Analysis from the Phase 3 ECHELON-1 Study. Blood, 2018, 132, 1618-1618. | 0.6 | 10 |
| 281 | Accurate Diagnosis of Aggressive B Cell Non-Hodgkin Lymphomas Using Gene Expression Profiling of Formalin-Fixed, Paraffin-Embedded Tissues. Blood, 2014, 124, 3016-3016. | 0.6 | 10 |
| 282 | Non-Hodgkin Lymphoma Risk and Variants in Genes Controlling Lymphocyte Development. PLoS ONE, 2013, 8, e75170. | 1.1 | 10 |
| 283 | Ketoconazole Versus Nystatin as Prophylaxis Against Fungal Infection for Lymphoma Patients Receiving Chemotherapy. American Journal of Clinical Oncology: Cancer Clinical Trials, 1987, 10, 355-359. | 0.6 | 9 |
| 284 | New Strategies for the Treatment of Early Stages of Hodgkin's Lymphoma. Hematology/Oncology Clinics of North America, 2007, 21, 871-880. | 0.9 | 9 |
| 285 | Improved survival outcomes with the addition of rituximab to initial therapy for chronic lymphocytic leukemia: a comparative effectiveness analysis in the province of British Columbia, Canada. Leukemia and Lymphoma, 2018, 59, 1356-1363. | 0.6 | 9 |
| 286 | Validation of a simplified international prognostic score (IPSâ€3) in patients with advancedâ€stage classic Hodgkin lymphoma. British Journal of Haematology, 2020, 189, 122-127. | 1.2 | 9 |
| 287 | Resources-Stratified Guidelines for Classical Hodgkin Lymphoma. International Journal of Environmental Research and Public Health, 2020, 17, 1783. | 1.2 | 9 |
| 288 | Outcome of relapsed and refractory nodular lymphocyteâ€predominant Hodgkin lymphoma: a North American analysis. British Journal of Haematology, 2021, 192, 560-567. | 1.2 | 9 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 289 | Rapid Infusion Rituximab in Combination with Steroid Containing Chemotherapy Can Be Given Safely and Substantially Reduces Resource Utilization Blood, 2004, 104, 1407-1407. | 0.6 | 9 |
| 290 | Mechanisms of Bcl-2 Protein Expression in Diffuse Large B-Cell Lymphoma (DLBCL) Blood, 2004, 104, 26-26. | 0.6 | 9 |
| 291 | FDG-PET Scan Guided Consolidative Radiation Therapy Optimizes Outcome In Patients with Advanced-Stage Diffuse Large B-Cell Lymphoma (DLBCL) with Residual Abnormalities on CT Scan Following R-CHOP. Blood, 2010, 116, 854-854. | 0.6 | 9 |
| 292 | Frontline Therapy with Brentuximab Vedotin Combined with ABVD or AVD in Patients with Newly Diagnosed Advanced Stage Hodgkin Lymphoma. Blood, 2012, 120, 798-798. | 0.6 | 9 |
| 293 | Five-Year Survival Data from a Pivotal Phase 2 Study of Brentuximab Vedotin in Patients with Relapsed or Refractory Systemic Anaplastic Large Cell Lymphoma. Blood, 2016, 128, 4144-4144. | 0.6 | 9 |
| 294 | Prognostic Significance of Bcl-2 Protein Expression and Bcl-2 Gene Rearrangement in Diffuse Aggressive Non-Hodgkin's Lymphoma. Blood, 1997, 90, 244-251. | 0.6 | 9 |
| 295 | Controversies and Future Directions in the Therapy of Lymphoma. Leukemia and Lymphoma, 1998, 30, 1-2. | 0.6 | 8 |
| 296 | A systematic evaluation of the <i>ataxia telangiectasia mutated</i> gene does not show an association with nonâ€Hodgkin lymphoma. International Journal of Cancer, 2007, 121, 1967-1975. | 2.3 | 8 |
| 297 | Autologous Stem Cell Transplantation Is Superior to Myeloablative Allogeneic SCT as a Salvage Therapy for Patients with Refractory/Relapsed Transformed Lymphoma. Blood, 2008, 112, 4459-4459. | 0.6 | 8 |
| 298 | Outcome in Unselected Patients with Relapsed/Refractory Diffuse Large B-Cell Lymphoma (DLBCL) Following R-CHOP When Stem Cell Transplantation Is Not Feasible. Blood, 2014, 124, 3069-3069. | 0.6 | 8 |
| 299 | Five-Year Survival Data Demonstrating Durable Responses from a Pivotal Phase 2 Study of Brentuximab Vedotin in Patients with Relapsed or Refractory Hodgkin Lymphoma. Blood, 2015, 126, 2736-2736. | 0.6 | 8 |
| 300 | Hodgkin lymphoma in the elderly, pregnant, and HIV-infected. Seminars in Hematology, 2016, 53, 203-208. | 1.8 | 7 |
| 301 | Nonrandom occurrence of lymphoid cancer types in 140 families. Leukemia and Lymphoma, 2017, 58, 2134-2143. | 0.6 | 7 |
| 302 | Leukocytoclastic vasculitis following lenalidomide during the treatment of follicular lymphoma. Leukemia and Lymphoma, 2017, 58, 711-714. | 0.6 | 7 |
| 303 | New Treatment Algorithms in Hodgkin Lymphoma: Too Much or Too Little?. American Society of Clinical Oncology Educational Book / ASCO American Society of Clinical Oncology Meeting, 2018, 38, 626-636. | 1.8 | 7 |
| 304 | Overall Survival Benefit for Patients with Relapsed Hodgkin Lymphoma Treated with Brentuximab Vedotin After Autologous Stem Cell Transplant. Blood, 2012, 120, 3701-3701. | 0.6 | 7 |
| 305 | A Clinicogenetic Risk Model (m7-FLIPI) Prospectively Identifies One-Half of Patients with Early Disease Progression of Follicular Lymphoma after First-Line Immunochemotherapy. Blood, 2015, 126, 333-333. | 0.6 | 7 |
| 306 | Prior Rituximab Reduces Relapse and Improves Survival Following High Dose Chemotherapy and Stem Cell Transplantation for Relapsed Composite Low and Intermediate Grade (Including Transformed) Lymphoma Blood, 2006, 108, 3662-3662. | 0.6 | 7 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 307 | The cured lymphoma patient. Postgraduate Medicine, 1982, 72, 53-59. | 0.9 | 6 |
| 308 | I. Hodgkin lymphoma: special challenges and solutions. Hematological Oncology, 2015, 33, 21-24. | 0.8 | 6 |
| 309 | Quantifying Benefit of Autologous Transplantation for Relapsed Follicular Lymphoma Patients via Instrumental Variable Analysis. Biology of Blood and Marrow Transplantation, 2016, 22, 941-948. | 2.0 | 6 |
| 310 | Outcome of patients with relapsed or refractory anaplastic large cell lymphoma who have failed brentuximab vedotin. Hematological Oncology, 2019, 37, 35-38. | 0.8 | 6 |
| 311 | Who Actually Detects Relapse in Hodgkin Lymphoma: Patient or Physician? Blood, 2004, 104, 3124-3124. | 0.6 | 6 |
| 312 | Mutations In MLL2 and MEF2B Genes In Follicular Lymphoma and Diffuse Large B-Cell Lymphoma. Blood, 2010, 116, 473-473. | 0.6 | 6 |
| 313 | The Outcome of Advanced Stage Nodular Lymphocyte Predominant Hodgkin's Lymphoma (NLPHL) Compared to Classical Hodgkin's Lymphoma (CHL): A Matched Pair Analysis. Blood, 2012, 120, 1531-1531. | 0.6 | 6 |
| 314 | Long-Term Remissions Observed in an Ongoing Phase 2 Study of Brentuximab Vedotin in Patients with Relapsed or Refractory Systemic Anaplastic Large Cell Lymphoma Blood, 2012, 120, 2745-2745. | 0.6 | 6 |
| 315 | Clinicopathological Analysis of Follicular Lymphoma with a Polyploid Karyotype. Leukemia and Lymphoma, 1998, 28, 383-389. | 0.6 | 5 |
| 316 | More Is Not Necessarily Better When Treating Hodgkin's Lymphoma. Journal of Clinical Oncology, 2011, 29, 4215-4216. | 0.8 | 5 |
| 317 | Chronic Lymphocytic Leukemia Patients With Deletion 11q Have a Short Time to Requirement of First-Line Therapy, But Long Overall Survival: Results of a Population-Based Cohort in British Columbia, Canada. Clinical Lymphoma, Myeloma and Leukemia, 2017, 17, 382-389. | 0.2 | 5 |
| 318 | Aberrant cytoplasmic expression of MHCII confers worse progression free survival in diffuse large B-cell lymphoma. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin, 2017, 470, 113-117. | 1.4 | 5 |
| 319 | Frontline Therapy with Bendamustine and Rituximab (BR) in Follicular Lymphoma: Prognosis Among Patients with Progression of Disease By 24 Months (POD24) Is Poor with Majority Having Transformed Lymphoma. Blood, 2018, 132, 2873-2873. | 0.6 | 5 |
| 320 | Autologous Stem-Cell Transplant with a Rituximab Purge and Maintenance vs. Standard Chemotherapy for Mantle Cell Lymphoma: Extended Follow-Up of a Matched Pair Analysis Blood, 2006, 108, 3051-3051. | 0.6 | 5 |
| 321 | Follicular Non-Hodgkin Lymphoma Grade 3a and 3b Subtypes Exhibit Similar Clinical Behaviour and Treatment Outcome. Blood, 2008, 112, 3777-3777. | 0.6 | 5 |
| 322 | Clinical Significance of Genetic Aberrations in Diffuse Large B Cell Lymphoma. Blood, 2014, 124, 703-703. | 0.6 | 5 |
| 323 | Brentuximab vedotin with chemotherapy for stage III or IV Hodgkin lymphoma (HL): Impact of cycle 2 PET result on modified progression-free survival (mPFS) Journal of Clinical Oncology, 2018, 36, 7539-7539. | 0.8 | 5 |
| 324 | Targeted Sequencing Reveals Novel Gene Mutations Associated with Transformation and Early Progression in Follicular Lymphoma. Blood, 2016, 128, 2919-2919. | 0.6 | 5 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 325 | Insulin Therapy in Diabetic Ketoacidosis. Annals of Internal Medicine, 1977, 86, 109. | 2.0 | 4 |
| 326 | Identification of Risk Factors in Patients Treated for First Relapse of Hodgkin's Disease. Leukemia and Lymphoma, 1994, 15, 189-200. | 0.6 | 4 |
| 327 | Treatment of Diffuse Large B-Cell Lymphoma: A Risk-Based Approach. Clinical Lymphoma and Myeloma, 2006, 7, S14-S19. | 1.4 | 4 |
| 328 | Genetic polymorphism at BCL2 as a predictor for rituximab, cyclophosphamide, doxorubicin, vincristine and prednisone efficacy in patients with diffuse large B-cell lymphoma. Haematologica, 2017, 102, e199-e202. | 1.7 | 4 |
| 329 | Outcome of limited-stage nodular lymphocyte-predominant Hodgkin lymphoma and the impact of a PET-adapted approach. Blood Advances, 2021, 5, 3647-3655. | 2.5 | 4 |
| 330 | CRIS: complete reconstruction of immunoglobulin <i>V-D-J</i> sequences from RNA-seq data. Bioinformatics Advances, 2021, 1, vbab021. | 0.9 | 4 |
| 331 | Outcome of Patients with Chemotherapy Refractory and Early Progressive Diffuse Large B Cell Lymphoma After R-CHOP Treatment. Blood, 2010, 116, 1751-1751. | 0.6 | 4 |
| 332 | Whole Transcriptome Sequencing Reveals Recurrent NOTCH1 Mutations in Mantle Cell Lymphoma. Blood, 2011, 118, 436-436. | 0.6 | 4 |
| 333 | Long-Term Survival Analyses of an Ongoing Phase 2 Study of Brentuximab Vedotin in Patients with Relapsed or Refractory Hodgkin Lymphoma. Blood, 2012, 120, 3689-3689. | 0.6 | 4 |
| 334 | ABVD Chemotherapy Results in Excellent Outcomes with Reduced Radiation Therapy Rates in Children, Adolescents and Young Adults with Hodgkin Lymphoma. Blood, 2015, 126, 2695-2695. | 0.6 | 4 |
| 335 | Phase 3 study of brentuximab vedotin plus doxorubicin, vinblastine, and dacarbazine (A+AVD) versus doxorubicin, bleomycin, vinblastine, and dacarbazine (ABVD) as front-line treatment for advanced classical Hodgkin lymphoma (HL): Echelon-1 study Journal of Clinical Oncology, 2014, 32, TPS8613-TPS8613. | 0.8 | 4 |
| 336 | Outcomes of Hodgkin variant Richter transformation in chronic lymphocytic leukaemia and small lymphocytic lymphoma in British Columbia. British Journal of Haematology, 2022, 198, 684-692. | 1.2 | 4 |
| 337 | Advanced-Stage Hodgkin Lymphoma. Cancer Journal (Sudbury, Mass), 2018, 24, 230-236. | 1.0 | 3 |
| 338 | Birth Order, Sibship Size, Childhood Environment and Immune-Related Disorders, and Risk of Lymphoma in Lymphoid Cancer Families. Cancer Epidemiology Biomarkers and Prevention, 2020, 29, 1168-1178. | 1.1 | 3 |
| 339 | Transformed Lymphoma: Incidence and Long-Term Outcome Blood, 2004, 104, 3253-3253. | 0.6 | 3 |
| 340 | Relapse Patterns and Subsequent Outcomes of Patients Treated on the NCIC CTG HD.6 (ECOG JHD06) Randomized Trial Evaluating ABVD Alone in Patients with Limited Stage Hodgkin Lymphoma (HL) Blood, 2005, 106, 817-817. | 0.6 | 3 |
| 341 | FDC-PET Guided Consolidative Radiotherapy in Patients with Advanced Stage Hodgkin Lymphoma with Residual Abnormalities on Post Chemotherapy CT Scan Blood, 2007, 110, 213-213. | 0.6 | 3 |
| 342 | The Architecural Pattern of FOXP3+ T Cells Predicts Risk of Transformation in Patients with Follicular Lymphoma (FL) Blood, 2007, 110, 358-358. | 0.6 | 3 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 343 | HLA-DR Protein Expression Correlates with Non-Neoplastic T-Cell Infiltration and Predicts Survival in Patients with Primary Mediastinal Large B Cell Lymphoma (PMBCL) Treated with CHOP Chemotherapy Blood, 2009, 114, 133-133. | 0.6 | 3 |
| 344 | Diffuse Large B-Cell Lymphoma with Testicular Involvement: Outcome and Risk of CNS Relapse in the Rituximab Era. Blood, 2011, 118, 780-780. | 0.6 | 3 |
| 345 | A Phase 1 -2 Study of Brentuximab Vedotin (Bv) and Bendamustine (B) in Patients with Relapsed or Refractory Hodgkin Lymphoma (HL) and Anaplastic Large T-Cell Lymphoma (ALCL). Blood, 2014, 124, 3084-3084. | 0.6 | 3 |
| 346 | Outcome of Observation As Initial Strategy in Patients with Mantle Cell Lymphoma. Blood, 2015, 126, 2699-2699. | 0.6 | 3 |
| 347 | Second Malignancies in Patients With Hairy Cell Leukemia in British Columbia: A 20-Year Experience. Blood, 1998, 92, 1160-1164. | 0.6 | 3 |
| 348 | Is There a Role for MACOP-B in the Treatment of Diffuse Large Cell Lymphoma?. Leukemia and Lymphoma, 1993, 10, 85-89. | 0.6 | 2 |
| 349 | The Case for Chemotherapy Alone for Limited-Stage Hodgkin's Lymphoma. Oncologist, 2012, 17, 1011-1013. | 1.9 | 2 |
| 350 | Identification of a novel non-immunoglobulin/MYC translocation t(8;12)(q24;p12) involving the LRMP gene in a primary B-cell lymphoma. A case report. Leukemia Research, 2012, 36, e22-e24. | 0.4 | 2 |
| 351 | Treatment of NLPHL. Blood, 2013, 122, 4288-4289. | 0.6 | 2 |
| 352 | Development and characterization of a Mantle Cell Lymphoma Cell Bank in the American Type Culture Collection. Leukemia and Lymphoma, 2015, 56, 2114-2122. | 0.6 | 2 |
| 353 | Response assessment in lymphoma: Concordance between independent central review and local evaluation in a clinical trial setting. Clinical Trials, 2016, 13, 545-554. | 0.7 | 2 |
| 354 | Tattoos and Hematologic Malignancies in British Columbia, Canada. Cancer Epidemiology Biomarkers and Prevention, 2020, 29, 2093-2095. | 1.1 | 2 |
| 355 | Anaplastic large cell lymphoma: a clinicopathologic analysis. Hematological Oncology, 1999, 17, 137-148. | 0.8 | 2 |
| 356 | Molecular and Genetic Characterization of MHC Deficiency Identifies EZH2 As a Therapeutic Target for Restoring MHC Expression in Diffuse Large B-Cell Lymphoma. Blood, 2018, 132, 1560-1560. | 0.6 | 2 |
| 357 | Hematopoietic Stem Cell Transplant (HSCT) as Primary Treatment for T-Cell Lymphobastic Lymphoma (T-LBL) : An Intention to Treat Analysis Blood, 2004, 104, 900-900. | 0.6 | 2 |
| 358 | Allogeneic Stem Cell Transplantation as Treatment for Relapsed and High-Risk Peripheral T-Cell Lymphoma Blood, 2007, 110, 3040-3040. | 0.6 | 2 |
| 359 | LMO2 Protein Expression Predicts Survival in Patients with Diffuse Large B-Cell Lymphoma in the Pre- and Post-Rituximab Treatment Eras Blood, 2007, 110, 52-52. | 0.6 | 2 |
| 360 | CARD11 as an Oncogene in Diffuse Large B Cell Lymphoma Blood, 2007, 110, 692-692. | 0.6 | 2 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 361 | Deletion in Chromosome 17p12 and Gains in Chromosome 9q33.3 by Array Comparative Hybridization Are Associated with R-CHOP Treatment Failure in Patients with Diffuse Large B Cell Lymphoma. Blood, 2008, 112, 477-477. | 0.6 | 2 |
| 362 | Number of Lymphoma-Associated-Macrophages (LAM) Is An Independent Predictor of Survival in Patients with Mantle Cell Lymphoma (MCL) Blood, 2009, 114, 3944-3944. | 0.6 | 2 |
| 363 | Incorporation of ABVD Increases Cure Rates of Patients with Limited Stage Nodular Lymphocyte Predominant Hodgkin Lymphoma (NLPHL). Blood, 2010, 116, 3887-3887. | 0.6 | 2 |
| 364 | Hodgkin Lymphoma Patients with Stage II B or Stage II Bulky Disease Have Advanced Disease and Should Not Be Included In Limited Stage Trials. Blood, 2010, 116, 417-417. | 0.6 | 2 |
| 365 | Final Analysis of a Randomized Comparison of ABVD Chemotherapy with a Strategy That Includes Radiation Therapy (RT) in Patients with Limited-Stage Hodgkin Lymphoma (HL): NCIC CTG/ECOG HD.6. Blood, 2011, 118, 590-590. | 0.6 | 2 |
| 366 | Survival of Peripheral T-Cell Lymphomas (PTCLs) Patients Following Relapse: Spectrum of Disease and Rare Long-Term Survivors. Blood, 2011, 118, 96-96. | 0.6 | 2 |
| 367 | An International Collaborative Study of Outcome and Prognostic Factors in Patients with Secondary CNS Involvement By Diffuse Large B-Cell Lymphoma. Blood, 2016, 128, 1874-1874. | 0.6 | 2 |
| 368 | Prognostic significance of the proliferation signature in mantle cell lymphoma measured using digital gene expression in formalin-fixed paraffin-embedded tissue biopsies Journal of Clinical Oncology, 2016, 34, 7510-7510. | 0.8 | 2 |
| 369 | EFS24 as a predictor of outcome in a population-based cohort of patients with DLBCL in British Columbia (BC) Journal of Clinical Oncology, 2016, 34, 7569-7569. | 0.8 | 2 |
| 370 | CD20 Mutations at the Rituximab Binding Site Are Rare and Are Not a Significant Cause of R-CHOP Resistance in Patients with De Novo Diffuse Large B-Cell Lymphoma Blood, 2007, 110, 686-686. | 0.6 | 2 |
| 371 | Concurrent BCL2 and MYC Protein Expression by Immunohistochemistry Determines Clinical Outcome In DLBCL Patients Treated with R-CHOP. Blood, 2010, 116, 2005-2005. | 0.6 | 2 |
| 372 | Hepatic Dysfunction During and After Lymphoma Chemotherapy in Patients With Hepatitis C. Journal of Clinical Gastroenterology, 2006, 40, 636-638. | 1.1 | 1 |
| 373 | Clinical Impact of Pathology Reviews of Outside Material: A Patient-Focused Approach. Journal of Clinical Oncology, 2011, 29, 4212-4213. | 0.8 | 1 |
| 374 | Sex- and Subtype-Specific Analysis of H2AFX Polymorphisms in Non-Hodgkin Lymphoma. PLoS ONE, 2013, 8, e74619. | 1.1 | 1 |
| 375 | Chemotherapy Only in Early Stage Hodgkin Lymphoma: More Relapses but the Same Survival – What to Do?. Current Hematologic Malignancy Reports, 2014, 9, 217-221. | 1.2 | 1 |
| 376 | Is the lymphoma better? Not easy to determine. Blood, 2016, 128, 2481-2482. | 0.6 | 1 |
| 377 | Immune Checkpoint Inhibition: A New Era in Lymphoma Treatment. Journal of Oncology Practice, 2016, 12, 109-110. | 2.5 | 1 |
| 378 | Hodgkin lymphoma, treatment and thrombosis: a dangerous mix. Annals of Oncology, 2019, 30, 1192-1193. | 0.6 | 1 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 379 | Molecular Features of Primary Central Nervous System Lymphoma in a Large Tissue Microarray. Blood, 2018, 132, 348-348. | 0.6 | 1 |
| 380 | The Double-Hit Gene Expression Signature Defines a Clinically and Biologically Distinct Subgroup within GCB-DLBCL. Blood, 2018, 132, 921-921. | 0.6 | 1 |
| 381 | Prolonged Overall Survival (OS) in a Subset of Responders to the Combination of Brentuximab Vedotin (Bv) and Bendamustine (B) in Heavily Treated Patients with Relapsed or Refractory Hodgkin Lymphoma (HL): Results of an International Multi- Center Phase I/II Experience. Blood, 2018, 132, 2907-2907. | 0.6 | 1 |
| 382 | Outcome of Limited Stage Nodular Lymphocyte Predominant Hodgkin Lymphoma (NLPHL) and Evaluation of a PET-Adapted Approach. Blood, 2019, 134, 2845-2845. | 0.6 | 1 |
| 383 | Lymphoma-Associated Macrophage (LAM) Content Is an Independent Predictor of Survival in Patients with Follicular Lymphoma (FL) Blood, 2004, 104, 3259-3259. | 0.6 | 1 |
| 384 | Survival of Limited Stage Peripheral T-Cell Lymphoma Is Similar To Diffuse Large B-Cell Lymphoma Blood, 2005, 106, 2817-2817. | 0.6 | 1 |
| 385 | Insights into Disease Evolution of Transformed Follicular Lymphoma Derived from Cytogenetics Blood, 2005, 106, 604-604. | 0.6 | 1 |
| 386 | Clinicopathologic Subtypes of Mantle Cell Lymphoma (MCL) Show Distinct Patterns of Genetic Copy Number Alteration Blood, 2006, 108, 2049-2049. | 0.6 | 1 |
| 387 | Comparison of Outcome Between Refractory/Relapsed De Novo Diffuse Large B-Cell and Transformed Lymphoma Using Related and Unrelated Allogeneic Hematopoietic SCT Blood, 2008, 112, 2173-2173. | 0.6 | 1 |
| 388 | The Percentage of Cytotoxic T-Cells in Mantle Cell Lymphoma (MCL) Biopsies Predicts Response to Rituximab Blood, 2009, 114, 2923-2923. | 0.6 | 1 |
| 389 | Identification of Genes Frequently Mutated In FL and DLBCL with Transcriptome, Genome and Exome Sequencing. Blood, 2010, 116, 804-804. | 0.6 | 1 |
| 390 | A Gene Expression Signature in Diagnostic Formalin Fixed Paraffin Embedded Tissue Predicts Overall Survival in Locally Advanced and Advanced Stage Classical Hodgkin Lymphoma – a Correlative Study From the E2496 Intergroup Trial. Blood, 2011, 118, 430-430. | 0.6 | 1 |
| 391 | FAS Mutations Accelerate Lymphoma Growth and Induce Therapeutic Resistance. Blood, 2014, 124, 3018-3018. | 0.6 | 1 |
| 392 | Rituximab with High Dose Methotrexate in the Management of Primary Central Nervous System Diffuse Large B-Cell Lymphoma. Blood, 2014, 124, 3090-3090. | 0.6 | 1 |
| 393 | The Absolute Number of Extranodal Sites Detected By PET-CT Is a Powerful Predictor of Secondary Central Nervous System Involvement in Patients with Diffuse Large B-Cell Lymphoma Treated with R-CHOP. Blood, 2015, 126, 3905-3905. | 0.6 | 1 |
| 394 | Both Discordant and Concordant Bone Marrow (BM) Involvement Predict for a Poorer Outcome Independent of the IPI in Patients with Diffuse Large B-Cell Lymphoma (DLBCL) Treated with R-CHOP Blood, 2007, 110, 1559-1559. | 0.6 | 1 |
| 395 | Molecular Signatures Implicate Innate Immune Cells, Fibrosis, and Angiogenesis in Survival Following R-CHOP Treatment of Diffuse Large B Cell Lymphoma. Blood, 2008, 112, 475-475. | 0.6 | 1 |
| 396 | Genetic Alterations Detected by High-Resolution Array Comparative Genomic Hybridization in Microdissected HRS Cells Correlate with Treatment Outcome in Classical Hodgkin Lymphoma. Blood, 2008, 112, 522-522. | 0.6 | 1 |

| # | Article | IF | CITATIONS |
|-----|---|--------------------|-------------------------------|
| 397 | TNFRSF14 Is Mutated in a Subset of Follicular Lymphoma and Correlated with Inferior Prognosis Blood, 2009, 114, 1919-1919. | 0.6 | 1 |
| 398 | Protein Tyrosine Phosphatase Type-1 (PTPN1) Is Frequently Mutated In Primary Mediastinal B Cell Lymphoma and Hodgkin Lymphoma. Blood, 2013, 122, 242-242. | 0.6 | 1 |
| 399 | Role of Bone Marrow Biopsy in the Staging of Diffuse Large B-Cell Lymphoma in the PET/CT Era. Blood, 2014, 124, 2960-2960. | 0.6 | 1 |
| 400 | Prognostic Impact of Extranodal Diffuse Large B-Cell Lymphoma in the Era of Immunochemotherapy and PET/CT Staging. Blood, 2014, 124, 1630-1630. | 0.6 | 1 |
| 401 | Outcomes in Adolescents and Young Adults (AYA) with Hodgkin Lymphoma (HL) Treated on US Cooperative Group Protocols: An Adult Intergroup (E2496) and Children's Oncology Group (COG) Tj ETQq1 | 10.78 ⊕3 ₫4 | rgBT1/Overlo <mark>c</mark> k |
| 402 | Frequent Genetic Alterations of PI3K-AKT Pathway and Their Clinical Significance in Germinal Center B-Cell-like Diffuse Large B-Cell Lymphoma. Blood, 2016, 128, 607-607. | 0.6 | 1 |
| 403 | Outcome of elderly patients with classical Hodgkin lymphoma (HL) in British Columbia Journal of Clinical Oncology, 2020, 38, 8031-8031. | 0.8 | 1 |
| 404 | Primary Testicular Lymphoma. Journal of Urology, 1987, 137, . | 0.2 | 0 |
| 405 | Reply to D.E. Roos. Journal of Clinical Oncology, 2012, 30, 1147-1147. | 0.8 | Ο |
| 406 | Interview: Learning from lymphocytes: a career studying lymphoid cancer. International Journal of Hematologic Oncology, 2014, 3, 15-17. | 0.7 | 0 |
| 407 | Introduction From the Guest Editor: Hodgkin Lymphoma. Cancer Journal (Sudbury, Mass), 2018, 24, 205-205. | 1.0 | Ο |
| 408 | Letter comments on a published article in the New England Journal of Medicine. European Journal of Cancer, 2018, 104, 250-251. | 1.3 | 0 |
| 409 | To treat or not to treat, that is the NLPHL question. Blood, 2019, 133, 2113-2114. | 0.6 | Ο |
| 410 | HLA-DR Protein Status Predicts Survival in Patients with Diffuse Large B Cell Lymphoma (DLBCL) Treated with the MACOP-B Chemotherapy Regimen Blood, 2004, 104, 3273-3273. | 0.6 | 0 |
| 411 | Acceptable Outcomes for Patients Receiving Allogeneic Hematopoietic Stem Cell Transplantation (AlloSCT) for Relapsed Aggressive Non-Hodgkin's Lymphoma (NHL) Blood, 2004, 104, 3325-3325. | 0.6 | Ο |
| 412 | High Cyclin D1 Expression Is Associated with Increased Proliferation Rate and Decreased Survival in Mantle Cell Lymphoma (MCL) and Is Caused by Genomic Deletions and Mutations that Enhance Stability of Cyclin D1 mRNA Blood, 2004, 104, 697-697. | 0.6 | 0 |
| 413 | Primary Central Nervous System Lymphoma of T Cell Origin: A Descriptive Analysis of 45 Cases from the International PCNSL Collaborative Group Blood, 2004, 104, 1372-1372. | 0.6 | 0 |
| 414 | Long Term Results of Myeloablative Allogeneic Stem Cell Transplantation Using Related and Unrelated Donors in Patients with Relapsed Composite Low and Intermediate Grade (Including Transformed) Lymphoma Blood, 2006, 108, 3139-3139. | 0.6 | 0 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 415 | Outcome in Patients with Diffuse Large B-Cell Lymphoma (DLBCL) Treated with CHOP-R Can Be Predicted by Stage and Serum Lactate Dehydrogenase (LDH) Level Blood, 2006, 108, 2739-2739. | 0.6 | 0 |
| 416 | Bone Marow (BM) Involvement in Diffuse Large B Cell Lymphoma (DLBCL): Clinical Impact of Discordant Disease Blood, 2006, 108, 2031-2031. | 0.6 | 0 |
| 417 | Myeloablative Allogeneic Stem Cell Transplantation for Relapsed Composite Low and Intermediate Grade Lymphoma Is Not Superior to Autologous Stem Cell Transplantation - Reduced Relapse Risk Is Offset by Higher Treatment Related Mortality Blood, 2006, 108, 3039-3039. | 0.6 | Ο |
| 418 | Long Term Follow-Up Results for Patients with Hodgkin Lymphoma after First Remission of Longer Than Ten Years: A Population Based Study of 954 Patients from the British Columbia Cancer Agency Blood, 2006, 108, 2269-2269. | 0.6 | 0 |
| 419 | Correlation of DNA Copy Number, Gene Expression, Protein Expression and Clinical Outcome in a Group of Patients with Diffuse Large B Cell Lymphoma (DLBCL) Treated with CHOP Plus Rituximab (CHOP-R) Blood, 2006, 108, 2027-2027. | 0.6 | Ο |
| 420 | Vascularization Predicts Overall Survival (OS) & Risk of Transformation (RT) in Uniformly Treated Patients with Follicular Lymphoma (FL) Blood, 2007, 110, 184-184. | 0.6 | 0 |
| 421 | Impact of Comorbidity Index on Outcome with Allogeneic Hematopoetic Stem Cell Transplantation for Chronic Lymphocytic Leukemia. Blood, 2008, 112, 3305-3305. | 0.6 | Ο |
| 422 | Genome-Wide Expression Profiling Predicts Treatment Outcome in Classical Hodgkin Lymphoma. Blood, 2008, 112, 520-520. | 0.6 | 0 |
| 423 | Base-Pair Resolution of Somatic and Germline-Derived Genome Rearrangement Breakpoints in Follicular Lymphoma Blood, 2009, 114, 439-439. | 0.6 | 0 |
| 424 | Exposure to Novel Agents Increases Post Relapse Survival in Patients with High Risk Myeloma Defined by Early Relapse (<12 months) Blood, 2009, 114, 2872-2872. | 0.6 | 0 |
| 425 | Fludarabine (F) and Rituximab (R) (FR) as Initial Therapy for Symptomatic Chronic Lymphocytic Leukemia (CLL) or Small Lymphocytic Lymphoma (SLL): Population-Based Experience Matches Clinical Trials Blood, 2009, 114, 2363-2363. | 0.6 | 0 |
| 426 | The Management of Hodgkin Lymphoma During Pregnancy. , 2011, , 241-248. | | 0 |
| 427 | The Prognosis of Limited Stage Peripheral T-Cell Lymphoma (PTCL): A Population-Based Analysis and Comparison to Diffuse Large B-Cell Lymphoma (DLBCL). Blood, 2010, 116, 4129-4129. | 0.6 | Ο |
| 428 | Fludarabine and Rituximab (FR) Is a Safe and Effective Treatment Alternative for Relapsed or Refractory Hairy Cell Leukemia (HCL). Blood, 2010, 116, 2454-2454. | 0.6 | 0 |
| 429 | Recurrent DNA Mutations In Non-Hodgkin Lymphomas Reveal Candidate Therapeutic Targets. Blood, 2010, 116, 632-632. | 0.6 | Ο |
| 430 | Front-Line Therapy with Rituximab, Cyclophosphamide, Vincristine, and Prednisone (R-CVP) Followed by 2 Years of Rituximab Maintenance for Follicular Lymphoma (FL) Is Associated with Excellent Outcomes and Improved Progression-Free Survival (PFS) In Comparison to No Maintenance. Blood, 2010, 116, 1803-1803. | 0.6 | 0 |
| 431 | BCL2 Is Highly Mutated In Diffuse Large B-Cell Lymphoma. Blood, 2010, 116, 4187-4187. | 0.6 | 0 |
| 432 | CSF1R Expression of Hodgkin Reed Sternberg Cells Is Associated with the Number of Macrophages in the Tumor Microenvironment and Is Correlated with Treatment Outcome. Blood, 2011, 118, 427-427. | 0.6 | 0 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 433 | Chemoresistance Can Be Reliably Overcome with High-Dose Therapy and Autologous Stem Cell Transplantation (HDT/ASCT) for Relapsed and Refractory Hodgkin's Lymphoma. Blood, 2011, 118, 2022-2022. | 0.6 | 0 |
| 434 | Population Survey of Mantle Cell Lymphoma (MCL) in British Columbia (BC) – A Heterogeneous Disorder with Improved Outcomes in the Modern Era. Blood, 2012, 120, 1602-1602. | 0.6 | 0 |
| 435 | FAS Mutations Are Associated with Therapeutic Resistance in Follicular Lymphoma. Blood, 2012, 120, 1549-1549. | 0.6 | Ο |
| 436 | Prognostic Factors in Patients with HIV-Associated Hodgkin Lymphoma: An Analysis of 199 Cases. Blood, 2012, 120, 1528-1528. | 0.6 | 0 |
| 437 | Large-Scale High Resolution Integration of Copy Number and Gene Expression in DLBCL Reveals Focal and Frequent Deletions in Chromatin Modifying Genes with Outcome Correlation. Blood, 2012, 120, 295-295. | 0.6 | 0 |
| 438 | Abstract A10: MLL2 interactions in follicular and diffuse large B-cell lymphoma. , 2013, , . | | 0 |
| 439 | Genetic Alterations In Immune Cell Crosstalk Genes In Diffuse Large B-Cell Lymphoma Predict Survival. Blood, 2013, 122, 500-500. | 0.6 | 0 |
| 440 | Determining Cell-Of-Origin Subtypes In Diffuse Large B-Cell Lymphoma Using Gene Expression Profiling On Formalin-Fixed Paraffin-Embedded Tissue – An L.L.M.P.P. Project. Blood, 2013, 122, 73-73. | 0.6 | 0 |
| 441 | The Use Of GDP (Gemcitabine, Dexamethasone and Cisplatin) in The Primary Therapy Of Peripheral T-Cell Lymphomas. Blood, 2013, 122, 1804-1804. | 0.6 | 0 |
| 442 | The outcome of patients with "nodal" peripheral T-cell lymphomas in a complete response following standard chemotherapy Journal of Clinical Oncology, 2014, 32, 8555-8555. | 0.8 | 0 |
| 443 | Abstract 959: FAS mutations induce therapeutic resistance in non-Hodgkin lymphomas. , 2014, , . | | 0 |
| 444 | Analysis of Relapse Biopsies in Classical Hodgkin Lymphoma Reveals Correlations with Outcome after Autologous Stem Cell Transplantation. Blood, 2014, 124, 136-136. | 0.6 | 0 |
| 445 | Cell-of-Origin Subtype Classification of Diffuse Large B-Cell Lymphoma Using the Lymph2Cx Assay Retains Relevance in the Context of BCL2 and MYC Expression Status. Blood, 2014, 124, 1667-1667. | 0.6 | Ο |
| 446 | Primary Cutaneous Anaplastic Large Cell Lymphoma: The British Columbia Cancer Agency Experience. Blood, 2014, 124, 3076-3076. | 0.6 | 0 |
| 447 | Maintenance Rituximab Following R-CHOP Chemotherapy in Patients with Composite or Discordant, Indolent and Aggressive, B-Cell Non-Hodgkin Lymphomas. Blood, 2015, 126, 3950-3950. | 0.6 | Ο |
| 448 | An International Assessment of Event-Free Survival at 24 Months (EFS24) and Subsequent Survival in Peripheral T-Cell Lymphoma. Blood, 2016, 128, 920-920. | 0.6 | 0 |
| 449 | Outcome of Hodgkin Lymphoma after Progression Following Autologous Stem Cell Transplantation. Blood, 2016, 128, 2994-2994. | 0.6 | 0 |
| 450 | Divergent Modes of Tumor Evolution Underlie Histological Transformation and Early Progression of Follicular Lymphoma. Blood, 2016, 128, 1091-1091. | 0.6 | 0 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 451 | Outcome of Patients with Peripheral T-Cell Lymphoma Undergoing Allogeneic Stem Cell in British Columbia. Blood, 2016, 128, 5852-5852. | 0.6 | Ο |
| 452 | Population-Based Survival Outcomes in Adult Patients with Burkitt Lymphoma (BL) Treated with Cyclophosphamide, Vincristine, Doxorubicin, High-Dose Methotrexate (CODOX-M)/Ifosfamide, Etoposide and High-Dose Cytarabine (IVAC) Plus or Minus Rituximab (R) in British Columbia (BC), Canada. Blood, 2016, 128, 1113-1113. | 0.6 | 0 |
| 453 | Abstract 2445: Integrative genetic analysis identifies therapeutic relevance of cell of origin-specific genetic alterations in diffuse large B-cell lymphoma. , 2017, , . | | 0 |
| 454 | Validation of a Simplified International Prognostic Score (IPS-3) in Patients with Advanced Stage Hodgkin Lymphoma. Blood, 2018, 132, 2916-2916. | 0.6 | 0 |
| 455 | Hodgkin Variant of Richter's Transformation (HvRT) Among Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL) Patients in British Columbia (BC), Canada. Blood, 2020, 136, 13-15. | 0.6 | 0 |
| 456 | Cardiac Morbidity in Adolescents and Young Adult Survivors of Hodgkin Lymphoma. Blood, 2020, 136, 18-19. | 0.6 | 0 |
| 457 | Real-World Characterization of Ibrutinib Therapy for Chronic Lymphocytic Leukemia (CLL) and Small Lymphocytic Lymphoma (SLL) Patients in British Columbia (BC). Blood, 2020, 136, 33-34. | 0.6 | 0 |
| 458 | Cost-Effectiveness of Molecularly Guided Treatment in Diffuse Large B-Cell Lymphoma (DLBCL) in Patients under 60. Cancers, 2022, 14, 908. | 1.7 | 0 |
| 459 | Abstract 3480: <i>TMEM30A</i> loss-of-function mutations drive lymphomagenesis and confer therapeutically exploitable vulnerability in B-cell lymphoma. , 2019, , . | | 0 |
| 460 | Radioimmunotherapy for orbital marginal zone lymphoma: a retrospective review. Leukemia and Lymphoma, 2022, , 1-4. | 0.6 | 0 |
| 461 | Outcomes after initial refusal of curative treatment in patients with classic Hodgkin lymphoma. Leukemia and Lymphoma, 2022, 63, 2739-2742. | 0.6 | 0 |