

Peter E Wright

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3050365/publications.pdf>

Version: 2024-02-01

341
papers

47,694
citations

1457

107
h-index

1895

208
g-index

364
all docs

364
docs citations

364
times ranked

31397
citing authors

#	ARTICLE	IF	CITATIONS
1	Multivalency enables unidirectional switch-like competition between intrinsically disordered proteins. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2022, 119, .	3.3	22
2	Interactions of a Long Noncoding RNA with Domains of NF- κ B and I κ B β : Implications for the Inhibition of Non-Signal-Related Phosphorylation. <i>Biochemistry</i> , 2022, 61, 367-376.	1.2	4
3	Structural Biology â€“ Painting the Mechanistic Landscape of Biomolecules. <i>Journal of Molecular Biology</i> , 2022, 434, 167566.	2.0	1
4	Client Specificity of an ATPâ€independent Chaperone is Regulated by a Temperature Sensitive Switch. <i>FASEB Journal</i> , 2022, 36, .	0.2	0
5	A transthyretin monomer intermediate undergoes local unfolding and transient interaction with oligomers in a kinetically concerted aggregation pathway. <i>Journal of Biological Chemistry</i> , 2022, 298, 102162.	1.6	5
6	Modeling of Hidden Structures Using Sparse Chemical Shift Data from NMR Relaxation Dispersion. <i>Biophysical Journal</i> , 2021, 120, 296-305.	0.2	4
7	Thermodynamic Stability and Aggregation Kinetics of EF Helix and EF Loop Variants of Transthyretin. <i>Biochemistry</i> , 2021, 60, 756-764.	1.2	14
8	Role of Active Site Loop Dynamics in Mediating Ligand Release from <i>E. coli</i> Dihydrofolate Reductase. <i>Biochemistry</i> , 2021, 60, 2663-2671.	1.2	4
9	The molecular basis of allostery in a facilitated dissociation process. <i>Structure</i> , 2021, 29, 1327-1338.e5.	1.6	6
10	NMR illuminates intrinsic disorder. <i>Current Opinion in Structural Biology</i> , 2021, 70, 44-52.	2.6	60
11	A phosphorylation-dependent switch in the disordered p53 transactivation domain regulates DNA binding. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	3.3	33
12	Characterization of the High-Affinity Fuzzy Complex between the Disordered Domain of the E7 Oncoprotein from High-Risk HPV and the TAZ2 Domain of CBP. <i>Biochemistry</i> , 2021, 60, 3887-3898.	1.2	9
13	A Disorder-to-Order Transition Activates an ATP-Independent Membrane Protein Chaperone. <i>Journal of Molecular Biology</i> , 2020, 432, 166708.	2.0	8
14	An allosteric peptide inhibitor of HIF-1 α regulates hypoxia-induced retinal neovascularization. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 28297-28306.	3.3	31
15	RNA Binding by the KTS Splice Variants of Wilmsâ€™ Tumor Suppressor Protein WT1. <i>Biochemistry</i> , 2020, 59, 3889-3901.	1.2	4
16	A Conformational Switch in the Zinc Finger Protein Kaiso Mediates Differential Readout of Specific and Methylated DNA Sequences. <i>Biochemistry</i> , 2020, 59, 1909-1926.	1.2	7
17	Determining Binding Kinetics of Intrinsically Disordered Proteins by NMR Spectroscopy. <i>Methods in Molecular Biology</i> , 2020, 2141, 663-681.	0.4	3
18	Comparison of backbone dynamics of the p50 dimerization domain of NF κ B in the homodimeric transcription factor NF κ B1 and in its heterodimeric complex with RelA (p65). <i>Protein Science</i> , 2019, 28, 2064-2072.	3.1	7

#	ARTICLE	IF	CITATIONS
19	Perspective: the essential role of NMR in the discovery and characterization of intrinsically disordered proteins. <i>Journal of Biomolecular NMR</i> , 2019, 73, 651-659.	1.6	48
20	Role of Backbone Dynamics in Modulating the Interactions of Disordered Ligands with the TAZ1 Domain of the CREB-Binding Protein. <i>Biochemistry</i> , 2019, 58, 1354-1362.	1.2	33
21	A Dynamic Switch in Inactive p38 ^β Leads to an Excited State on the Pathway to an Active Kinase. <i>Biochemistry</i> , 2019, 58, 5160-5172.	1.2	7
22	Expanding the Paradigm: Intrinsically Disordered Proteins and Allosteric Regulation. <i>Journal of Molecular Biology</i> , 2018, 430, 2309-2320.	2.0	105
23	Slow Dynamics of Tryptophan-Water Networks in Proteins. <i>Journal of the American Chemical Society</i> , 2018, 140, 675-682.	6.6	26
24	CH ₂ -O Hydrogen Bonds Mediate Highly Specific Recognition of Methylated CpG Sites by the Zinc Finger Protein Kaiso. <i>Biochemistry</i> , 2018, 57, 2109-2120.	1.2	19
25	Long-range regulation of p53 DNA binding by its intrinsically disordered N-terminal transactivation domain. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018, 115, E11302-E11310.	3.3	93
26	Mispacking of the Phe87 Side Chain Reduces the Kinetic Stability of Human Transthyretin. <i>Biochemistry</i> , 2018, 57, 6919-6922.	1.2	8
27	Structural Basis for Graded Inhibition of CREB:DNA Interactions by Multisite Phosphorylation. <i>Biochemistry</i> , 2018, 57, 6964-6972.	1.2	7
28	Structural basis for cooperative regulation of KIX-mediated transcription pathways by the HTLV-1 HBZ activation domain. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018, 115, 10040-10045.	3.3	18
29	How Do Intrinsically Disordered Viral Proteins Hijack the Cell?. <i>Biochemistry</i> , 2018, 57, 4045-4046.	1.2	22
30	Tight complexes from disordered proteins. <i>Nature</i> , 2018, 555, 37-38.	13.7	3
31	NMR Measurements Reveal the Structural Basis of Transthyretin Destabilization by Pathogenic Mutations. <i>Biochemistry</i> , 2018, 57, 4421-4430.	1.2	30
32	Kinetic analysis of the multistep aggregation pathway of human transthyretin. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018, 115, E6201-E6208.	3.3	29
33	Hypersensitive termination of the hypoxic response by a disordered protein switch. <i>Nature</i> , 2017, 543, 447-451.	13.7	140
34	Role of the CBP catalytic core in intramolecular SUMOylation and control of histone H3 acetylation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, E5335-E5342.	3.3	56
35	How Does Your Protein Fold? Elucidating the Apomyoglobin Folding Pathway. <i>Accounts of Chemical Research</i> , 2017, 50, 105-111.	7.6	44
36	Fluorotryptophan Incorporation Modulates the Structure and Stability of Transthyretin in a Site-Specific Manner. <i>Biochemistry</i> , 2017, 56, 5570-5581.	1.2	20

#	ARTICLE	IF	CITATIONS
37	Defining the Structural Basis for Allosteric Product Release from <i>E. coli</i> Dihydrofolate Reductase Using NMR Relaxation Dispersion. <i>Journal of the American Chemical Society</i> , 2017, 139, 11233-11240.	6.6	27
38	Structural Basis for Interaction of the Tandem Zinc Finger Domains of Human Muscleblind with Cognate RNA from Human Cardiac Troponin T. <i>Biochemistry</i> , 2017, 56, 4154-4168.	1.2	27
39	Finding Our Way in the Dark Proteome. <i>Journal of the American Chemical Society</i> , 2016, 138, 9730-9742.	6.6	111
40	NMR Characterization of Information Flow and Allosteric Communities in the MAP Kinase p38 β . <i>Scientific Reports</i> , 2016, 6, 28655.	1.6	19
41	Solid-State NMR Studies Reveal Native-like β -Sheet Structures in Transthyretin Amyloid. <i>Biochemistry</i> , 2016, 55, 5272-5278.	1.2	25
42	Mapping the interactions of adenoviral E1A proteins with the p160 nuclear receptor coactivator binding domain of CBP. <i>Protein Science</i> , 2016, 25, 2256-2267.	3.1	18
43	Recognition of the disordered p53 transactivation domain by the transcriptional adapter zinc finger domains of CREB-binding protein. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, E1853-62.	3.3	94
44	Role of Intrinsic Protein Disorder in the Function and Interactions of the Transcriptional Coactivators CREB-binding Protein (CBP) and p300. <i>Journal of Biological Chemistry</i> , 2016, 291, 6714-6722.	1.6	251
45	Conformational dynamics of a membrane protein chaperone enables spatially regulated substrate capture and release. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, E1615-24.	3.3	33
46	Multi-probe relaxation dispersion measurements increase sensitivity to protein dynamics. <i>Physical Chemistry Chemical Physics</i> , 2016, 18, 5789-5798.	1.3	9
47	Functional advantages of dynamic protein disorder. <i>FEBS Letters</i> , 2015, 589, 2433-2440.	1.3	162
48	Cofactor-Mediated Conformational Dynamics Promote Product Release From <i>Escherichia coli</i> Dihydrofolate Reductase via an Allosteric Pathway. <i>Journal of the American Chemical Society</i> , 2015, 137, 9459-9468.	6.6	45
49	Conformational propensities of intrinsically disordered proteins influence the mechanism of binding and folding. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, 9614-9619.	3.3	222
50	Intrinsically disordered proteins in cellular signalling and regulation. <i>Nature Reviews Molecular Cell Biology</i> , 2015, 16, 18-29.	16.1	1,849
51	pE-DB: a database of structural ensembles of intrinsically disordered and of unfolded proteins. <i>Nucleic Acids Research</i> , 2014, 42, D326-D335.	6.5	195
52	Combinatorial regulation of a signal-dependent activator by phosphorylation and acetylation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 17116-17121.	3.3	20
53	Assemblages: Functional units formed by cellular phase separation. <i>Journal of Cell Biology</i> , 2014, 206, 579-588.	2.3	227
54	Integrated description of protein dynamics from room-temperature X-ray crystallography and NMR. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, E445-54.	3.3	142

#	ARTICLE	IF	CITATIONS
55	The High-Risk HPV16 E7 Oncoprotein Mediates Interaction between the Transcriptional Coactivator CBP and the Retinoblastoma Protein pRb. <i>Journal of Molecular Biology</i> , 2014, 426, 4030-4048.	2.0	61
56	Side Chain Conformational Averaging in Human Dihydrofolate Reductase. <i>Biochemistry</i> , 2014, 53, 1134-1145.	1.2	8
57	Probing the Non-Native H Helix Translocation in Apomyoglobin Folding Intermediates. <i>Biochemistry</i> , 2014, 53, 3767-3780.	1.2	16
58	Accurate scoring of non-uniform sampling schemes for quantitative NMR. <i>Journal of Magnetic Resonance</i> , 2014, 246, 31-35.	1.2	57
59	Classification of Intrinsically Disordered Regions and Proteins. <i>Chemical Reviews</i> , 2014, 114, 6589-6631.	23.0	1,618
60	Structural Characterization of Interactions between the Double-Stranded RNA-Binding Zinc Finger Protein JAZ and Nucleic Acids. <i>Biochemistry</i> , 2014, 53, 1495-1510.	1.2	20
61	Automated identification of functional dynamic contact networks from X-ray crystallography. <i>Nature Methods</i> , 2013, 10, 896-902.	9.0	130
62	Fast and accurate fitting of relaxation dispersion data using the flexible software package GLOVE. <i>Journal of Biomolecular NMR</i> , 2013, 56, 275-283.	1.6	71
63	The CH2 domain of CBP/p300 is a novel zinc finger. <i>FEBS Letters</i> , 2013, 587, 2506-2511.	1.3	12
64	Divergent evolution of protein conformational dynamics in dihydrofolate reductase. <i>Nature Structural and Molecular Biology</i> , 2013, 20, 1243-1249.	3.6	153
65	Localized Structural Fluctuations Promote Amyloidogenic Conformations in Transthyretin. <i>Journal of Molecular Biology</i> , 2013, 425, 977-988.	2.0	65
66	A Distal Mutation Perturbs Dynamic Amino Acid Networks in Dihydrofolate Reductase. <i>Biochemistry</i> , 2013, 52, 4605-4619.	1.2	77
67	Modulation of allostery by protein intrinsic disorder. <i>Nature</i> , 2013, 498, 390-394.	13.7	295
68	Side-Chain Conformational Heterogeneity of Intermediates in the <i>Escherichia coli</i> Dihydrofolate Reductase Catalytic Cycle. <i>Biochemistry</i> , 2013, 52, 3464-3477.	1.2	16
69	Analysis of the RelA:CBP/p300 Interaction Reveals Its Involvement in NF- κ B-Driven Transcription. <i>PLoS Biology</i> , 2013, 11, e1001647.	2.6	118
70	Mechanisms of Transthyretin Inhibition of β -Amyloid Aggregation <i>In Vitro</i> . <i>Journal of Neuroscience</i> , 2013, 33, 19423-19433.	1.7	118
71	Functional Interactions of Intrinsically Disordered Proteins in Signaling Networks. <i>FASEB Journal</i> , 2013, 27, 459.3.	0.2	0
72	Molecular basis for recognition of methylated and specific DNA sequences by the zinc finger protein Kaiso. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, 15229-15234.	3.3	101

#	ARTICLE	IF	CITATIONS
73	Quantitative Analysis of Multisite Protein-Ligand Interactions by NMR: Binding of Intrinsically Disordered p53 Transactivation Subdomains with the TAZ2 Domain of CBP. <i>Journal of the American Chemical Society</i> , 2012, 134, 3792-3803.	6.6	123
74	Kaiso uses all three zinc fingers and adjacent sequence motifs for high affinity binding to sequence-specific and methyl-CpG DNA targets. <i>FEBS Letters</i> , 2012, 586, 734-739.	1.3	17
75	Structural Basis for Cooperative Transcription Factor Binding to the CBP Coactivator. <i>FASEB Journal</i> , 2012, 26, lb266.	0.2	0
76	A Dynamic Knockout Reveals That Conformational Fluctuations Influence the Chemical Step of Enzyme Catalysis. <i>Science</i> , 2011, 332, 234-238.	6.0	414
77	Consequences of Stabilizing the Natively Disordered F Helix for the Folding Pathway of Apomyoglobin. <i>Journal of Molecular Biology</i> , 2011, 411, 248-263.	2.0	16
78	Identification of endogenous ligands bound to bacterially expressed human and <i>E. coli</i> dihydrofolate reductase by 2D NMR. <i>FEBS Letters</i> , 2011, 585, 3528-3532.	1.3	14
79	Measurement of protein unfolding/refolding kinetics and structural characterization of hidden intermediates by NMR relaxation dispersion. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 9078-9083.	3.3	70
80	Leu628 of the KIX domain of CBP is a key residue for the interaction with the MLL transactivation domain. <i>FEBS Letters</i> , 2010, 584, 4500-4504.	1.3	32
81	3P040 Mapping the Interactions of the Intrinsically Disordered p53 Transactivation Subdomains with the TAZ2 Domain of CBP by NMR (Protein: Structure & Function, The 48th Annual Meeting of the IUPAC) 10.784614 rgBT (Overl	1.0	0
82	Graded enhancement of p53 binding to CREB-binding protein (CBP) by multisite phosphorylation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 19290-19295.	3.3	188
83	Millisecond timescale fluctuations in dihydrofolate reductase are exquisitely sensitive to the bound ligands. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 1373-1378.	3.3	133
84	Structure of the p53 Transactivation Domain in Complex with the Nuclear Receptor Coactivator Binding Domain of CREB Binding Protein. <i>Biochemistry</i> , 2010, 49, 9964-9971.	1.2	162
85	Energetic Frustration of Apomyoglobin Folding: Role of the B Helix. <i>Journal of Molecular Biology</i> , 2010, 396, 1319-1328.	2.0	17
86	Cooperative regulation of p53 by modulation of ternary complex formation with CBP/p300 and HDM2. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 6591-6596.	3.3	197
87	Structural basis for subversion of cellular control mechanisms by the adenoviral E1A oncoprotein. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 13260-13265.	3.3	119
88	Linking folding and binding. <i>Current Opinion in Structural Biology</i> , 2009, 19, 31-38.	2.6	932
89	The role of dynamic conformational ensembles in biomolecular recognition. <i>Nature Chemical Biology</i> , 2009, 5, 789-796.	3.9	1,649
90	Diagnostic chemical shift markers for loop conformation and substrate and cofactor binding in dihydrofolate reductase complexes. <i>Protein Science</i> , 2009, 12, 2230-2238.	3.1	38

#	ARTICLE	IF	CITATIONS
91	Mapping the Interactions of the p53 Transactivation Domain with the KIX Domain of CBP. <i>Biochemistry</i> , 2009, 48, 2115-2124.	1.2	109
92	Prion Proteins with Pathogenic and Protective Mutations Show Similar Structure and Dynamics. <i>Biochemistry</i> , 2009, 48, 8120-8128.	1.2	53
93	Editorial from the Editor-in-Chief. <i>Journal of Molecular Biology</i> , 2009, 385, 1-2.	2.0	1
94	Structural basis for recruitment of CBP/p300 coactivators by STAT1 and STAT2 transactivation domains. <i>EMBO Journal</i> , 2009, 28, 948-958.	3.5	147
95	Prediction of the Rotational Tumbling Time for Proteins with Disordered Segments. <i>Journal of the American Chemical Society</i> , 2009, 131, 6814-6821.	6.6	48
96	Structural characterization of partially folded intermediates of apomyoglobin H64F. <i>Protein Science</i> , 2008, 17, 313-321.	3.1	16
97	How Do Proteins Interact?. <i>Science</i> , 2008, 320, 1429-1430.	6.0	174
98	The Kinetic and Equilibrium Molten Globule Intermediates of Apoleghemoglobin Differ in Structure. <i>Journal of Molecular Biology</i> , 2008, 378, 715-725.	2.0	26
99	Overexpression of post-translationally modified peptides in <i>Escherichia coli</i> by co-expression with modifying enzymes. <i>Protein Expression and Purification</i> , 2008, 57, 108-115.	0.6	30
100	Hierarchical folding mechanism of apomyoglobin revealed by ultra-fast H/D exchange coupled with 2D NMR. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 13859-13864.	3.3	89
101	NMR Relaxation Study of the Complex Formed Between CBP and the Activation Domain of the Nuclear Hormone Receptor Coactivator ACTR. <i>Biochemistry</i> , 2008, 47, 1299-1308.	1.2	86
102	The Intrinsically Disordered RNR Inhibitor Sml1 Is a Dynamic Dimer. <i>Biochemistry</i> , 2008, 47, 13428-13437.	1.2	53
103	Conformational Relaxation following Hydride Transfer Plays a Limiting Role in Dihydrofolate Reductase Catalysis. <i>Biochemistry</i> , 2008, 47, 9227-9233.	1.2	53
104	Modeling transient collapsed states of an unfolded protein to provide insights into early folding events. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 6278-6283.	3.3	105
105	S03A3 Metastable structure detected by relaxation dispersion NMR spectroscopy(Visualising) Tj ETQq1 1 0.784314 rgBT /Overlock 10 T	0.9	0
106	PRAK Is Essential for ras-Induced Senescence and Tumor Suppression. <i>Cell</i> , 2007, 128, 295-308.	13.5	286
107	Embryonic Neural Inducing Factor Churchill Is not a DNA-binding Zinc Finger Protein: Solution Structure Reveals a Solvent-exposed Î²-Sheet and Zinc Binuclear Cluster. <i>Journal of Molecular Biology</i> , 2007, 371, 1274-1289.	2.0	21
108	Structure of the Wilms Tumor Suppressor Protein Zinc Finger Domain Bound to DNA. <i>Journal of Molecular Biology</i> , 2007, 372, 1227-1245.	2.0	91

#	ARTICLE	IF	CITATIONS
109	Tailoring Relaxation Dispersion Experiments for Fast-Associating Protein Complexes. <i>Journal of the American Chemical Society</i> , 2007, 129, 13406-13407.	6.6	52
110	Mechanism of coupled folding and binding of an intrinsically disordered protein. <i>Nature</i> , 2007, 447, 1021-1025.	13.7	984
111	The Dynamic Energy Landscape of Dihydrofolate Reductase Catalysis. <i>Science</i> , 2006, 313, 1638-1642.	6.0	877
112	An NMR Perspective on Enzyme Dynamics. <i>Chemical Reviews</i> , 2006, 106, 3055-3079.	23.0	424
113	NMR Solution Structure of the Peptide Fragment 1 [~] 30, Derived from Unprocessed Mouse Doppel Protein, in DHPC Micelles. <i>Biochemistry</i> , 2006, 45, 159-166.	1.2	19
114	Structural Basis for Cooperative Transcription Factor Binding to the CBP Coactivator. <i>Journal of Molecular Biology</i> , 2006, 355, 1005-1013.	2.0	166
115	Structure of the <i>Escherichia coli</i> Quorum Sensing Protein SdiA: Activation of the Folding Switch by Acyl Homoserine Lactones. <i>Journal of Molecular Biology</i> , 2006, 355, 262-273.	2.0	162
116	Identification of Native and Non-native Structure in Kinetic Folding Intermediates of Apomyoglobin. <i>Journal of Molecular Biology</i> , 2006, 355, 139-156.	2.0	112
117	Induced Fit and "Lock and Key" Recognition of 5S RNA by Zinc Fingers of Transcription Factor IIIA. <i>Journal of Molecular Biology</i> , 2006, 357, 275-291.	2.0	72
118	Solution Structure of the Hdm2 C2H2C4 RING, a Domain Critical for Ubiquitination of p53. <i>Journal of Molecular Biology</i> , 2006, 363, 433-450.	2.0	120
119	According to current textbooks, a well-defined three-dimensional structure is a prerequisite for the function of a protein. Is this correct?. <i>IUBMB Life</i> , 2006, 58, 107-109.	1.5	20
120	The role of hydrophobic interactions in initiation and propagation of protein folding. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 13057-13061.	3.3	266
121	Localization of Sites of Interaction between p23 and Hsp90 in Solution. <i>Journal of Biological Chemistry</i> , 2006, 281, 14457-14464.	1.6	58
122	Structure and Function of the CBP/p300 TAZ Domains. , 2005, , 114-120.		4
123	Intrinsically unstructured proteins and their functions. <i>Nature Reviews Molecular Cell Biology</i> , 2005, 6, 197-208.	16.1	3,403
124	Defining the role of active-site loop fluctuations in dihydrofolate reductase catalysis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 5032-5037.	3.3	152
125	Enhanced picture of protein-folding intermediates using organic solvents in H/D exchange and quench-flow experiments. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 4765-4770.	3.3	62
126	Inhibition of DNA Binding by Human Estrogen-Related Receptor 2 and Estrogen Receptor β with Minor Groove Binding Polyamides. <i>Biochemistry</i> , 2005, 44, 4196-4203.	1.2	42

#	ARTICLE	IF	CITATIONS
127	CBP/p300 TAZ1 Domain Forms a Structured Scaffold for Ligand Binding. <i>Biochemistry</i> , 2005, 44, 490-497.	1.2	76
128	Solution Structure of the N-terminal Zinc Fingers of the <i>Xenopus laevis</i> double-stranded RNA-binding Protein ZFa. <i>Journal of Molecular Biology</i> , 2005, 351, 718-730.	2.0	18
129	Sequence Determinants of a Protein Folding Pathway. <i>Journal of Molecular Biology</i> , 2005, 351, 383-392.	2.0	54
130	Elucidation of the Protein Folding Landscape by NMR. <i>Methods in Enzymology</i> , 2005, 394, 299-321.	0.4	90
131	Interaction of the TAZ1 Domain of the CREB-Binding Protein with the Activation Domain of CITED2. <i>Journal of Biological Chemistry</i> , 2004, 279, 3042-3049.	1.6	97
132	Recognition of the mRNA AU-rich element by the zinc finger domain of TIS11d. <i>Nature Structural and Molecular Biology</i> , 2004, 11, 257-264.	3.6	320
133	Model-free Analysis of Protein Dynamics: Assessment of Accuracy and Model Selection Protocols Based on Molecular Dynamics Simulation. <i>Journal of Biomolecular NMR</i> , 2004, 29, 243-257.	1.6	60
134	Unfolded Proteins and Protein Folding Studied by NMR. <i>ChemInform</i> , 2004, 35, no.	0.1	1
135	The LEF-1 High-Mobility Group Domain Undergoes a Disorder-to-Order Transition upon Formation of a Complex with Cognate DNA. <i>Biochemistry</i> , 2004, 43, 8725-8734.	1.2	62
136	Effect of Cofactor Binding and Loop Conformation on Side Chain Methyl Dynamics in Dihydrofolate Reductase. <i>Biochemistry</i> , 2004, 43, 374-383.	1.2	73
137	Conformational Changes in the Active Site Loops of Dihydrofolate Reductase during the Catalytic Cycle. <i>Biochemistry</i> , 2004, 43, 16046-16055.	1.2	119
138	Unfolded Proteins and Protein Folding Studied by NMR. <i>Chemical Reviews</i> , 2004, 104, 3607-3622.	23.0	596
139	Packing, specificity, and mutability at the binding interface between the p160 coactivator and CREB-binding protein. <i>Protein Science</i> , 2004, 13, 203-210.	3.1	66
140	Introduction: Biological Nuclear Magnetic Resonance. <i>Chemical Reviews</i> , 2004, 104, 3517-3518.	23.0	2
141	Solution Structure of the KIX Domain of CBP Bound to the Transactivation Domain of c-Myb. <i>Journal of Molecular Biology</i> , 2004, 337, 521-534.	2.0	181
142	Structural Characterization of Unfolded States of Apomyoglobin using Residual Dipolar Couplings. <i>Journal of Molecular Biology</i> , 2004, 340, 1131-1142.	2.0	165
143	ZZ Domain of CBP: an Unusual Zinc Finger Fold in a Protein Interaction Module. <i>Journal of Molecular Biology</i> , 2004, 343, 1081-1093.	2.0	81
144	Structure, Dynamics, and Catalytic Function of Dihydrofolate Reductase. <i>Annual Review of Biophysics and Biomolecular Structure</i> , 2004, 33, 119-140.	18.3	444

#	ARTICLE	IF	CITATIONS
145	The CBP/p300 TAZ1 domain in its native state is not a binding partner of MDM2. <i>Biochemical Journal</i> , 2004, 381, 685-691.	1.7	41
146	Changes in structure and dynamics of the Fv fragment of a catalytic antibody upon binding of inhibitor. <i>Protein Science</i> , 2003, 12, 1386-1394.	3.1	14
147	Role of a solvent-exposed tryptophan in the recognition and binding of antibiotic substrates for a metallo- β -lactamase. <i>Protein Science</i> , 2003, 12, 1368-1375.	3.1	56
148	Structure of the Nuclear Factor κ B: Insights into Post-Transcriptional Regulatory and mRNA Nuclear Export Processes. <i>Biochemistry</i> , 2003, 42, 7348-7357.	1.2	20
149	Role of the B Helix in Early Folding Events in Apomyoglobin: Evidence from Site-directed Mutagenesis for Native-like Long Range Interactions. <i>Journal of Molecular Biology</i> , 2003, 334, 293-307.	2.0	51
150	Monomeric Complex of Human Orphan Estrogen Related Receptor-2 with DNA: A Pseudo-dimer Interface Mediates Extended Half-site Recognition. <i>Journal of Molecular Biology</i> , 2003, 327, 819-832.	2.0	97
151	Folding of a β -sheet Protein Monitored by Real-time NMR Spectroscopy. <i>Journal of Molecular Biology</i> , 2003, 328, 1161-1171.	2.0	29
152	Roles of Phosphorylation and Helix Propensity in the Binding of the KIX Domain of CREB-binding Protein by Constitutive (c-Myb) and Inducible (CREB) Activators. <i>Journal of Biological Chemistry</i> , 2002, 277, 42241-42248.	1.6	134
153	Structural basis for Hif-1 α /CBP recognition in the cellular hypoxic response. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 5271-5276.	3.3	376
154	Molecular Hinges in Protein Folding: The Urea-Denatured State of Apomyoglobin. <i>Biochemistry</i> , 2002, 41, 12681-12686.	1.2	130
155	Insights into the structure and dynamics of unfolded proteins from nuclear magnetic resonance. <i>Advances in Protein Chemistry</i> , 2002, 62, 311-340.	4.4	208
156	The Apomyoglobin Folding Pathway Revisited: Structural Heterogeneity in the Kinetic Burst Phase Intermediate. <i>Journal of Molecular Biology</i> , 2002, 322, 483-489.	2.0	89
157	Cooperativity in Transcription Factor Binding to the Coactivator CREB-binding Protein (CBP). <i>Journal of Biological Chemistry</i> , 2002, 277, 43168-43174.	1.6	166
158	High Pressure NMR Reveals that Apomyoglobin is an Equilibrium Mixture from the Native to the Unfolded. <i>Journal of Molecular Biology</i> , 2002, 320, 311-319.	2.0	81
159	Mapping Long-range Contacts in a Highly Unfolded Protein. <i>Journal of Molecular Biology</i> , 2002, 322, 655-662.	2.0	140
160	Coupling of folding and binding for unstructured proteins. <i>Current Opinion in Structural Biology</i> , 2002, 12, 54-60.	2.6	1,223
161	Mutual synergistic folding in recruitment of CBP/p300 by p160 nuclear receptor coactivators. <i>Nature</i> , 2002, 415, 549-553.	13.7	423
162	Comparison of protein solution structures refined by molecular dynamics simulation in vacuum, with a generalized Born model, and with explicit water. <i>Journal of Biomolecular NMR</i> , 2002, 22, 317-331.	1.6	112

#	ARTICLE	IF	CITATIONS
163	Assignment of a 15 kDa protein complex formed between the p160 coactivator ACTR and CREB binding protein. <i>Journal of Biomolecular NMR</i> , 2002, 22, 377-378.	1.6	6
164	Structural and dynamic characterization of an unfolded state of poplar apo-plastocyanin formed under non-denaturing conditions. <i>Protein Science</i> , 2001, 10, 1056-1066.	3.1	79
165	NMR Structural and Dynamic Characterization of the Acid-Unfolded State of Apomyoglobin Provides Insights into the Early Events in Protein Folding. <i>Biochemistry</i> , 2001, 40, 3561-3571.	1.2	212
166	Backbone Dynamics in Dihydrofolate Reductase Complexes: A Role of Loop Flexibility in the Catalytic Mechanism. <i>Biochemistry</i> , 2001, 40, 9846-9859.	1.2	246
167	Nuclear Magnetic Resonance Methods for Elucidation of Structure and Dynamics in Disordered States. <i>Methods in Enzymology</i> , 2001, 339, 258-270.	0.4	148
168	Sequence-Dependent Correction of Random Coil NMR Chemical Shifts. <i>Journal of the American Chemical Society</i> , 2001, 123, 2970-2978.	6.6	562
169	Solution structure of <i>Escherichia coli</i> glutaredoxin-2 shows similarity to mammalian glutathione-S-transferases. <i>Journal of Molecular Biology</i> , 2001, 310, 907-918.	2.0	71
170	Solution structure of the third immunoglobulin domain of the neural cell adhesion molecule N-CAM: can solution studies define the mechanism of homophilic binding?. <i>Journal of Molecular Biology</i> , 2001, 311, 161-172.	2.0	25
171	Conformational and Dynamic Characterization of the Molten Globule State of an Apomyoglobin Mutant with an Altered Folding Pathway. <i>Biochemistry</i> , 2001, 40, 14459-14467.	1.2	45
172	Local Structural Plasticity of the Prion Protein. Analysis of NMR Relaxation Dynamics. <i>Biochemistry</i> , 2001, 40, 2743-2753.	1.2	171
173	SANE (Structure Assisted NOE Evaluation): an automated model-based approach for NOE assignment. <i>Journal of Biomolecular NMR</i> , 2001, 19, 321-329.	1.6	113
174	Anisotropic rotational diffusion in model-free analysis for a ternary DHFR complex. , 2001, 19, 209-230.		43
175	Potential bias in NMR relaxation data introduced by peak intensity analysis and curve fitting methods. <i>Journal of Biomolecular NMR</i> , 2001, 21, 1-9.	1.6	44
176	Zinc finger proteins: new insights into structural and functional diversity. <i>Current Opinion in Structural Biology</i> , 2001, 11, 39-46.	2.6	1,240
177	Conservation of folding pathways in evolutionarily distant globin sequences. <i>Nature Structural Biology</i> , 2000, 7, 679-686.	9.7	95
178	Assessment of zinc finger orientations by residual dipolar coupling constants. <i>Journal of Biomolecular NMR</i> , 2000, 16, 9-21.	1.6	43
179	Backbone H(N), N, C α , C' and C β assignments of the 19 kDa DHFR/NADPH complex at 9 degrees C and pH 7.6. <i>Journal of Biomolecular NMR</i> , 2000, 16, 349-350.	1.6	3
180	Assignment of ^1H , ^{13}C and ^{15}N resonances of the I-domain of human leukocyte function associated antigen-1. <i>Journal of Biomolecular NMR</i> , 2000, 16, 271-272.	1.6	7

#	ARTICLE	IF	CITATIONS
181	Random coil chemical shifts in acidic 8 M urea: implementation of random coil shift data in NMRView. <i>Journal of Biomolecular NMR</i> , 2000, 18, 43-48.	1.6	272
182	DNA-induced α -helix capping in conserved linker sequences is a determinant of binding affinity in Cys2-His2 zinc fingers. <i>Journal of Molecular Biology</i> , 2000, 295, 719-727.	2.0	137
183	NMR solution structure of the inserted domain of human leukocyte function associated antigen-1. <i>Journal of Molecular Biology</i> , 2000, 295, 1251-1264.	2.0	74
184	Solution Structure of the Cysteine-rich Domain of the Escherichia coli Chaperone Protein DnaJ. <i>Journal of Molecular Biology</i> , 2000, 300, 805-818.	2.0	121
185	NMR and molecular dynamics studies of the hydration of a zinc finger-DNA complex 1 Edited by M. F. Summers. <i>Journal of Molecular Biology</i> , 2000, 302, 1101-1117.	2.0	57
186	Solution structure of the TAZ2 (CH3) domain of the transcriptional adaptor protein CBP. <i>Journal of Molecular Biology</i> , 2000, 303, 243-253.	2.0	121
187	Solution structure and acetyl-lysine binding activity of the GCN5 bromodomain. <i>Journal of Molecular Biology</i> , 2000, 304, 355-370.	2.0	141
188	Structure of the PHD Zinc Finger from Human Williams-Beuren Syndrome Transcription Factor. <i>Journal of Molecular Biology</i> , 2000, 304, 723-729.	2.0	142
189	Identification of the regions involved in DNA binding by the mouse PEBP2 α protein. <i>FEBS Letters</i> , 2000, 470, 125-130.	1.3	15
190	Dynamics of the Metallo- β -Lactamase from <i>Bacteroides fragilis</i> in the Presence and Absence of a Tight-Binding Inhibitor. <i>Biochemistry</i> , 2000, 39, 13356-13364.	1.2	59
191	Alternative Splicing of Wilms' Tumor Suppressor Protein Modulates DNA Binding Activity through Isoform-Specific DNA-Induced Conformational Changes. <i>Biochemistry</i> , 2000, 39, 5341-5348.	1.2	58
192	Changes in the Apomyoglobin Folding Pathway Caused by Mutation of the Distal Histidine Residue. <i>Biochemistry</i> , 2000, 39, 11227-11237.	1.2	68
193	Native and Non-native Secondary Structure and Dynamics in the pH 4 Intermediate of Apomyoglobin. <i>Biochemistry</i> , 2000, 39, 2894-2901.	1.2	121
194	Amide proton hydrogen exchange rates for sperm whale myoglobin obtained from ^{15}N - ^1H NMR spectra. <i>Protein Science</i> , 2000, 9, 186-193.	3.1	23
195	Characterization of monomeric and dimeric B domain of Staphylococcal protein A. <i>Chemical Biology and Drug Design</i> , 1999, 54, 344-352.	1.2	5
196	Inherent flexibility in a potent inhibitor of blood coagulation, recombinant nematode anticoagulant protein c2. <i>FEBS Journal</i> , 1999, 265, 539-548.	0.2	42
197	Backbone resonance assignments for the Fv fragment of the catalytic antibody NPN43C9 with bound p-nitrophenol. <i>Journal of Biomolecular NMR</i> , 1999, 15, 83-84.	1.6	4
198	Improved low pH bicelle system for orienting macromolecules over a wide temperature range. <i>Journal of Biomolecular NMR</i> , 1999, 13, 387-391.	1.6	84

#	ARTICLE	IF	CITATIONS
199	Assignment of ¹ H, ¹³ C, and ¹⁵ N resonances of reduced Escherichia coli glutaredoxin 2. Journal of Biomolecular NMR, 1999, 14, 197-198.	1.6	3
200	Association between the first two immunoglobulin-like domains of the neural cell adhesion molecule N-CAM. FEBS Letters, 1999, 451, 162-168.	1.3	30
201	Determination of Fe ²⁺ Ligand Bond Lengths and the Fe ²⁺ N ³⁺ O Bond Angles in Soybean Ferrous and Ferric NitrosylhemoglobinaUsing Multiple-Scattering XAFS Analyses. Biochemistry, 1999, 38, 16491-16499.	1.2	20
202	NMR Characterization of the Metallo-β-lactamase from Bacteroides fragilis and Its Interaction with a Tight-Binding Inhibitor: Role of an Active-Site Loop. Biochemistry, 1999, 38, 14507-14514.	1.2	104
203	Effect of H helix destabilizing mutations on the kinetic and equilibrium folding of apomyoglobin 1 Edited by F. Cohen. Journal of Molecular Biology, 1999, 285, 269-282.	2.0	79
204	Structural analyses of CREB-CBP transcriptional activator-coactivator complexes by NMR spectroscopy: implications for mapping the boundaries of structural domains. Journal of Molecular Biology, 1999, 287, 859-865.	2.0	68
205	Identification of a minimal domain of 5 S ribosomal RNA sufficient for high affinity interactions with the RNA-specific zinc fingers of transcription factor IIIA 1 Edited by D. E. Draper. Journal of Molecular Biology, 1999, 291, 549-560.	2.0	19
206	Intrinsically unstructured proteins: re-assessing the protein structure-function paradigm. Journal of Molecular Biology, 1999, 293, 321-331.	2.0	2,668
207	Role of Secondary Structure in Discrimination between Constitutive and Inducible Activators. Molecular and Cellular Biology, 1999, 19, 5601-5607.	1.1	127
208	Quench-flow experiments combined with mass spectrometry show apomyoglobin folds through an obligatory intermediate. Protein Science, 1999, 8, 45-49.	3.1	93
209	Distributed torsion angle grid search in high dimensions: a systematic approach to NMR structure determination. Journal of Biomolecular NMR, 1998, 11, 241-263.	1.6	20
210	A NOESY-HSQC simulation program, SPIRIT. Journal of Biomolecular NMR, 1998, 11, 17-29.	1.6	17
211	¹ H, ¹³ C and ¹⁵ N NMR backbone assignments of 25.5 kDa metallo-beta-lactamase from Bacteroides fragilis. Journal of Biomolecular NMR, 1998, 12, 201-202.	1.6	5
212	Recommendations for the presentation of NMR structures of proteins and nucleic acids. IUPAC-IUBMB-IUPAB Inter-Union Task Group on the Standardization of Data Bases of Protein and Nucleic Acid Structures Determined by NMR Spectroscopy. Journal of Biomolecular NMR, 1998, 12, 1-23.	1.6	347
213	Chemical shift as a probe of molecular interfaces: NMR studies of DNA binding by the three amino-terminal zinc finger domains from transcription factor IIIA. Journal of Biomolecular NMR, 1998, 12, 51-71.	1.6	110
214	Structural and dynamic characterization of partially folded states of apomyoglobin and implications for protein folding. Nature Structural Biology, 1998, 18, 148-155.	9.7	344
215	Equilibrium NMR studies of unfolded and partially folded proteins. Nature Structural Biology, 1998, 5, 499-503.	9.7	187
216	Sequence requirements for stabilization of a peptide reverse turn in water solution. Proline is not essential for stability. FEBS Journal, 1998, 255, 462-471.	0.2	25

#	ARTICLE	IF	CITATIONS
217	Recommendations for the presentation of NMR structures of proteins and nucleic acids. IUPAC-IUBMB-IUPAB inter-union task group on the standardization of data bases of protein and nucleic acid structures determined by NMR spectroscopy. FEBS Journal, 1998, 256, 1-15.	0.2	137
218	High-resolution solution structure of Bacillus subtilis IIAGlc. , 1998, 31, 258-270.		4
219	The identification of metal-binding ligand residues in metalloproteins using nuclear magnetic resonance spectroscopy. Protein Science, 1998, 7, 2476-2479.	3.1	6
220	Analysis of an Activator:Coactivator Complex Reveals an Essential Role for Secondary Structure in Transcriptional Activation. Molecular Cell, 1998, 2, 353-359.	4.5	155
221	Conformational preferences in the Ser133 -phosphorylated and non-phosphorylated forms of the kinase inducible transactivation domain of CREB. FEBS Letters, 1998, 430, 317-322.	1.3	137
222	Recommendations for the presentation of NMR structures of proteins and nucleic acids. Journal of Molecular Biology, 1998, 280, 933-952.	2.0	217
223	High-resolution solution structure of the retinoid X receptor DNA-binding domain. Journal of Molecular Biology, 1998, 281, 271-284.	2.0	58
224	DNA-induced conformational changes are the basis for cooperative dimerization by the DNA binding domain of the retinoid X receptor. Journal of Molecular Biology, 1998, 284, 533-539.	2.0	66
225	NMR Spectroscopic Studies of the DNA-binding Domain of the Monomer-binding Nuclear Orphan Receptor, Human Estrogen Related Receptor-2. Journal of Biological Chemistry, 1997, 272, 18038-18043.	1.6	31
226	Contribution of Increased Length and Intact Capping Sequences to the Conformational Preference for Helix in a 31-Residue Peptide from the C Terminus of Myohemerythrin. Biochemistry, 1997, 36, 5234-5244.	1.2	44
227	Solution structure of the first three zinc fingers of TFIIIA bound to the cognate DNA sequence: determinants of affinity and sequence specificity. Journal of Molecular Biology, 1997, 273, 183-206.	2.0	182
228	Solution Structure of the KIX Domain of CBP Bound to the Transactivation Domain of CREB: A Model for Activator:Coactivator Interactions. Cell, 1997, 91, 741-752.	13.5	705
229	Populating the equilibrium molten globule state of apomyoglobin under conditions suitable for structural characterization by NMR. FEBS Letters, 1997, 417, 92-96.	1.3	53
230	Chemical shift dispersion and secondary structure prediction in unfolded and partly folded proteins. FEBS Letters, 1997, 419, 285-289.	1.3	135
231	Domain packing and dynamics in the DNA complex of the N-terminal zinc fingers of TFIIIA. Nature Structural Biology, 1997, 4, 605-608.	9.7	116
232	PCR-based gene synthesis and protein NMR spectroscopy. Structure, 1997, 5, 1407-1412.	1.6	22
233	Folding propensities of peptide fragments of myoglobin. Protein Science, 1997, 6, 706-716.	3.1	82
234	Absence of a stable intermediate on the folding pathway of protein A. Protein Science, 1997, 6, 1449-1457.	3.1	117

#	ARTICLE	IF	CITATIONS
235	Probing protein structure using biochemical and biophysical methods. <i>Journal of Chromatography A</i> , 1997, 777, 23-30.	1.8	58
236	INSIGHTS INTO PROTEIN FOLDING FROM NMR. <i>Annual Review of Physical Chemistry</i> , 1996, 47, 369-395.	4.8	144
237	Probing Protein/Protein Interactions with Mass Spectrometry and Isotopic Labeling:Â Analysis of the p21/Cdk2 Complex. <i>Journal of the American Chemical Society</i> , 1996, 118, 5320-5321.	6.6	62
238	Is Apomyoglobin a Molten Globule? Structural Characterization by NMR. <i>Journal of Molecular Biology</i> , 1996, 263, 531-538.	2.0	242
239	Gene synthesis, high-level expression and assignment of backbone ¹⁵ N and ¹³ C resonances of soybean leghemoglobin. <i>FEBS Letters</i> , 1996, 399, 283-289.	1.3	12
240	Biophysical methods Faster and bigger. <i>Current Opinion in Structural Biology</i> , 1996, 6, 583-584.	2.6	1
241	Hydrogen Exchange in the Carbon Monoxide Complex of Soybean Leghemoglobin. <i>FEBS Journal</i> , 1996, 237, 212-220.	0.2	17
242	Measurement of intrinsic exchange rates of amide protons in a ¹⁵ N-labeled peptide. <i>Journal of Biomolecular NMR</i> , 1996, 7, 341.	1.6	0
243	Assessment of major and minor groove DNA interactions by the zinc fingers of <i>Xenopus</i> transcription factor IIIA. <i>Nucleic Acids Research</i> , 1996, 24, 2567-2574.	6.5	12
244	NMR Structural Studies of Flexible Molecules. , 1996, , 245-249.		1
245	â€ Random coilâ€™ ¹ H chemical shifts obtained as a function of temperature and trifluoroethanol concentration for the peptide series GCXGC. <i>Journal of Biomolecular NMR</i> , 1995, 5, 14-24.	1.6	476
246	Structural basis for DNA bending by the architectural transcription factor LEF-1. <i>Nature</i> , 1995, 376, 791-795.	13.7	582
247	Antigenic peptides. <i>FASEB Journal</i> , 1995, 9, 37-42.	0.2	96
248	Interaction of the RNA binding Fingers of <i>Xenopus</i> Transcription Factor IIIA with Specific Regions of 5 S Ribosomal RNA. <i>Journal of Molecular Biology</i> , 1995, 248, 44-57.	2.0	47
249	Dynamics of the Dihydrofolate Reductase-Folate Complex: Catalytic Sites and Regions Known To Undergo Conformational Change Exhibit Diverse Dynamical Features. <i>Biochemistry</i> , 1995, 34, 11037-11048.	1.2	162
250	¹ H and ¹⁵ N resonance assignments and secondary structure of the carbon monoxide complex of sperm whale myoglobin. <i>Journal of Biomolecular NMR</i> , 1994, 4, 491-504.	1.6	49
251	¹ H, ¹⁵ N and ¹³ C resonance assignments, secondary structure, and the conformation of substrate in the binary folate complex of <i>Escherichia coli</i> dihydrofolate reductase. <i>Journal of Biomolecular NMR</i> , 1994, 4, 349-366.	1.6	28
252	High-resolution solution structures of oxidized and reduced <i>Escherichia coli</i> thioredoxin. <i>Structure</i> , 1994, 2, 853-868.	1.6	281

#	ARTICLE	IF	CITATIONS
253	NMR Assignments and Secondary Structure of the Retinoid X Receptor alpha DNA-binding Domain. Evidence for the Novel C-terminal Helix. FEBS Journal, 1994, 224, 639-650.	0.2	15
254	¹ H resonance assignments and secondary structure of the carbon monoxide complex of soybean leghemoglobin determined by homonuclear two-dimensional and three-dimensional NMR spectroscopy. FEBS Journal, 1994, 219, 611-626.	0.2	10
255	Dynamics of a flexible loop in dihydrofolate reductase from Escherichia coli and its implication for catalysis. Biochemistry, 1994, 33, 439-442.	1.2	150
256	Relative Contributions of the Zinc Fingers of Transcription Factor IIIA to the Energetics of DNA Binding. Journal of Molecular Biology, 1994, 244, 23-25.	2.0	66
257	Solution Structure of Carbonmonoxy Myoglobin Determined from Nuclear Magnetic Resonance Distance and Chemical Shift Constraints. Journal of Molecular Biology, 1994, 244, 183-197.	2.0	103
258	Stabilization of a type VI turn in a family of linear peptides in water solution. Journal of Molecular Biology, 1994, 243, 736-753.	2.0	152
259	Three-dimensional structure of a type VI turn in a linear peptide in water solution Evidence for stacking of aromatic rings as a major stabilizing factor. Journal of Molecular Biology, 1994, 243, 754-766.	2.0	118
260	[13] Use of chemical shifts and coupling constants in nuclear magnetic resonance structural studies on peptides and proteins. Methods in Enzymology, 1994, 239, 392-416.	0.4	111
261	NMR Order Parameters of Biomolecules: A New Analytical Representation and Application to the Gaussian Axial Fluctuation Model. Journal of the American Chemical Society, 1994, 116, 8426-8427.	6.6	131
262	Differential Side Chain Hydration in a Linear Peptide Containing a Type VI Turn. Journal of the American Chemical Society, 1994, 116, 12051-12052.	6.6	35
263	Peptide models of protein folding initiation sites. 3. The G-H helical hairpin of myoglobin. Biochemistry, 1993, 32, 6356-6364.	1.2	109
264	Peptide models of protein folding initiation sites. 1. Secondary structure formation by peptides corresponding to the G- and H-helices of myoglobin. Biochemistry, 1993, 32, 6337-6347.	1.2	209
265	Peptide models of protein folding initiation sites. 2. The G-H turn region of myoglobin acts as a helix stop signal. Biochemistry, 1993, 32, 6348-6355.	1.2	92
266	Peptide conformation and protein folding. Current Opinion in Structural Biology, 1993, 3, 60-65.	2.6	232
267	Biophysical methods Editorial overview. Current Opinion in Structural Biology, 1993, 3, 723-724.	2.6	1
268	Characterization of a folding intermediate of apoplastocyanin trapped by proline isomerization. Biochemistry, 1993, 32, 12299-12310.	1.2	103
269	Assignment of proton, nitrogen-15, and carbon-13 resonances, identification of elements of secondary structure and determination of the global fold of the DNA-binding domain of GAL4. Biochemistry, 1993, 32, 2144-2153.	1.2	30
270	Mapping of the binding interfaces of the proteins of the bacterial phosphotransferase system, HPr and IIAGlc. Biochemistry, 1993, 32, 32-37.	1.2	141

#	ARTICLE	IF	CITATIONS
271	Comparison of backbone and tryptophan side-chain dynamics of reduced and oxidized Escherichia coli thioredoxin using nitrogen-15 NMR relaxation measurements. <i>Biochemistry</i> , 1993, 32, 426-435.	1.2	148
272	NMR evidence for multiple conformations in a highly helical model peptide. <i>Biochemistry</i> , 1993, 32, 13089-13097.	1.2	61
273	Electrostatic calculations of side-chain pKa values in myoglobin and comparison with NMR data for histidines. <i>Biochemistry</i> , 1993, 32, 8045-8056.	1.2	237
274	Backbone dynamics of the Bacillus subtilis glucose permease IIA domain determined from nitrogen-15 NMR relaxation measurements. <i>Biochemistry</i> , 1992, 31, 4394-4406.	1.2	233
275	Suppression of the effects of cross-correlation between dipolar and anisotropic chemical shift relaxation mechanisms in the measurement of spin-spin relaxation rates. <i>Molecular Physics</i> , 1992, 75, 699-711.	0.8	287
276	Assignment of the aliphatic proton and carbon-13 resonances of the Bacillus subtilis glucose permease IIA domain using double- and triple-resonance heteronuclear three-dimensional NMR spectroscopy. <i>Biochemistry</i> , 1992, 31, 4413-4425.	1.2	55
277	Immunogenic peptides corresponding to the dominant antigenic region alanine-597 to cysteine-619 in the transmembrane protein of simian immunodeficiency virus have a propensity to fold in aqueous solution. <i>Biochemistry</i> , 1992, 31, 1458-1463.	1.2	15
278	Functional role of a mobile loop of Escherichia coli dihydrofolate reductase in transition-state stabilization. <i>Biochemistry</i> , 1992, 31, 7826-7833.	1.2	100
279	Low resolution solution structure of the Bacillus subtilis glucose permease IIA domain derived from heteronuclear three-dimensional NMR spectroscopy. <i>FEBS Letters</i> , 1992, 296, 148-152.	1.3	47
280	Structural determinants of Cys2 His2 zinc fingers. <i>FEBS Letters</i> , 1992, 296, 11-15.	1.3	20
281	The zinc finger motif Conservation of chemical shifts and correlation with structure. <i>FEBS Letters</i> , 1992, 309, 29-32.	1.3	23
282	Specific interaction of the first three zinc fingers of TFIIIA with the internal control region of the Xenopus 5 S RNA gene. <i>Journal of Molecular Biology</i> , 1992, 223, 857-871.	2.0	106
283	Folding of peptide fragments comprising the complete sequence of proteins. <i>Journal of Molecular Biology</i> , 1992, 226, 795-817.	2.0	385
284	Folding of peptide fragments comprising the complete sequence of proteins. <i>Journal of Molecular Biology</i> , 1992, 226, 819-835.	2.0	226
285	Improved resolution in three-dimensional constant-time triple resonance NMR spectroscopy of proteins. <i>Journal of Biomolecular NMR</i> , 1992, 2, 103-108.	1.6	52
286	Relationship between ¹ H and ¹³ C NMR chemical shifts and the secondary and tertiary structure of a zinc finger peptide. <i>Journal of Biomolecular NMR</i> , 1992, 2, 307-322.	1.6	39
287	Evidence for two interconverting protein isomers in the methotrexate complex of dihydrofolate reductase from Escherichia coli. <i>Biochemistry</i> , 1991, 30, 2184-2191.	1.2	50
288	Intramolecular motions of a zinc finger DNA-binding domain from Xfin characterized by proton-detected natural abundance carbon-13 heteronuclear NMR spectroscopy. <i>Journal of the American Chemical Society</i> , 1991, 113, 4371-4380.	6.6	616

#	ARTICLE	IF	CITATIONS
289	Polypeptide backbone resonance assignments and secondary structure of Bacillus subtilis enzyme IIIgIc determined by two-dimensional and three-dimensional heteronuclear NMR spectroscopy. <i>Biochemistry</i> , 1991, 30, 6896-6907.	1.2	56
290	Zinc is required for folding and binding of a single zinc finger to DNA. <i>FEBS Letters</i> , 1991, 279, 289-294.	1.3	28
291	High-resolution solution structure of reduced French bean plastocyanin and comparison with the crystal structure of poplar plastocyanin. <i>Journal of Molecular Biology</i> , 1991, 221, 533-555.	2.0	114
292	Kinetics and mechanisms of reduction of Cu(II) and Fe(III) complexes by soybean leghemoglobin. <i>BBA - Proteins and Proteomics</i> , 1991, 1079, 182-196.	2.1	18
293	Measurement of relaxation time constants for methyl groups by proton-detected heteronuclear NMR spectroscopy. <i>Chemical Physics Letters</i> , 1991, 185, 41-46.	1.2	61
294	Three-dimensional solution structure of the reduced form of Escherichia coli thioredoxin determined by nuclear magnetic resonance spectroscopy. <i>Biochemistry</i> , 1990, 29, 4129-4136.	1.2	177
295	Partial proton NMR assignments of the Escherichia coli dihydrofolate reductase complex with folate: evidence for a unique conformation of bound folate. <i>Biochemistry</i> , 1990, 29, 9667-9677.	1.2	35
296	Computational methods for determining protein structures from NMR data. <i>Biochemical Pharmacology</i> , 1990, 40, 15-22.	2.0	72
297	Antigen-antibody interactions: An NMR approach. <i>Biochemical Pharmacology</i> , 1990, 40, 83-88.	2.0	21
298	THE H-HELIX OF MYOGLOBIN AS A POTENTIAL INDEPENDENT PROTEIN FOLDING DOMAIN. , 1990, , 283-293.		7
299	Proton NMR studies of the solution conformations of an analog of the C-peptide of ribonuclease A. <i>Biochemistry</i> , 1989, 28, 7059-7064.	1.2	162
300	Assignment of the proton NMR spectrum of reduced and oxidized thioredoxin: sequence-specific assignments, secondary structure, and global fold. <i>Biochemistry</i> , 1989, 28, 7074-7087.	1.2	102
301	Conformation of a T cell stimulating peptide in aqueous solution. <i>FEBS Letters</i> , 1989, 250, 400-404.	1.3	49
302	Complete assignment of the ¹ H NMR spectrum of a synthetic zinc finger from Xfin Sequential resonance assignments and secondary structure. <i>FEBS Letters</i> , 1989, 254, 159-164.	1.3	24
303	What can two-dimensional NMR tell us about proteins?. <i>Trends in Biochemical Sciences</i> , 1989, 14, 255-260.	3.7	19
304	[6] Multiple-quantum nuclear magnetic resonance. <i>Methods in Enzymology</i> , 1989, 176, 114-134.	0.4	28
305	NMR studies of the conformations of leghemoglobins from soybean and lupin. <i>FEBS Journal</i> , 1988, 178, 419-435.	0.2	15
306	Proton NMR studies of human C3a anaphylatoxin in solution: sequential resonance assignments, secondary structure, and global fold. <i>Biochemistry</i> , 1988, 27, 9139-9148.	1.2	39

#	ARTICLE	IF	CITATIONS
307	Folding of immunogenic peptide fragments of proteins in water solution. <i>Journal of Molecular Biology</i> , 1988, 201, 161-200.	2.0	685
308	Folding of immunogenic peptide fragments of proteins in water solution. <i>Journal of Molecular Biology</i> , 1988, 201, 201-217.	2.0	477
309	Conformation of peptide fragments of proteins in aqueous solution: implications for initiation of protein folding. <i>Biochemistry</i> , 1988, 27, 7167-7175.	1.2	505
310	Complete assignment of the ¹ H nuclear magnetic resonance spectrum of French bean plastocyanin. <i>Journal of Molecular Biology</i> , 1988, 202, 603-622.	2.0	83
311	Complete assignment of the ¹ H nuclear magnetic resonance spectrum of French bean plastocyanin. <i>Journal of Molecular Biology</i> , 1988, 202, 623-636.	2.0	67
312	Complete assignment of lysine resonances in ¹ H NMR spectra of proteins as probes of surface structure and dynamics. <i>FEBS Letters</i> , 1987, 222, 109-114.	1.3	20
313	¹ H resonances of proximal histidine in CO complexes of hemoglobins provide a sensitive probe of coordination geometry. <i>FEBS Letters</i> , 1987, 213, 289-292.	1.3	8
314	Assignment of resonances in the ¹ H nuclear magnetic resonance spectrum of the carbon monoxide complex of sperm whale myoglobin by phase-sensitive two-dimensional techniques. <i>Journal of Molecular Biology</i> , 1987, 194, 313-327.	2.0	90
315	Electron transfer from cytochrome b5 to iron and copper complexes. <i>Biochemistry</i> , 1987, 26, 7102-7107.	1.2	34
316	A modified strategy for identification of ¹ H spin systems in proteins. <i>Biopolymers</i> , 1987, 26, 973-977.	1.2	52
317	Kinetics and mechanisms of the oxidation of myoglobin by Fe(III) and Cu(II) complexes. <i>BBA - Proteins and Proteomics</i> , 1987, 912, 384-397.	2.1	41
318	NMR studies of the heme pocket conformations of monomeric hemoglobins from <i>Glycera dibranchiata</i> . Implications for ligand binding. <i>FEBS Journal</i> , 1987, 166, 399-408.	0.2	17
319	Structural consequences of heme isomerism in monomeric hemoglobins from <i>Glycera dibranchiata</i> . <i>FEBS Journal</i> , 1987, 166, 409-414.	0.2	8
320	Proton NMR studies of plastocyanin: assignment of aromatic and methyl group resonances from two-dimensional spectra. <i>Biochemistry</i> , 1986, 25, 2364-2374.	1.2	17
321	¹ H NMR studies of heme pocket conformation in zinc-substituted leghemoglobin, a diamagnetic analog of deoxyleghemoglobin. <i>Journal of Inorganic Biochemistry</i> , 1986, 28, 303-309.	1.5	9
322	The immunodominant site of a synthetic immunogen has a conformational preference in water for a type-II reverse turn. <i>Nature</i> , 1985, 318, 480-483.	13.7	246
323	Differences in amino acid composition and heme electronic structure of the multiple monomeric hemoglobin components of <i>Glycera dibranchiata</i> . <i>BBA - Proteins and Proteomics</i> , 1985, 832, 357-364.	2.1	25
324	Heme orientation in the major monomeric hemoglobins of <i>Glycera dibranchiata</i> . <i>BBA - Proteins and Proteomics</i> , 1985, 832, 365-372.	2.1	21

#	ARTICLE	IF	CITATIONS
325	Assignment of heme and distal amino acid resonances in the ¹ H-NMR spectra of the carbon monoxide and oxygen complexes of sperm whale myoglobin. <i>BBA - Proteins and Proteomics</i> , 1985, 832, 175-185.	2.1	62
326	Simplification of ¹ H NMR spectra of proteins by one-dimensional multiple quantum filtration. <i>Biochemical and Biophysical Research Communications</i> , 1985, 131, 1094-1102.	1.0	17
327	Conformational disorder of the distal leucine in monomeric Glycerahemoglobins and implications for oxygen binding. <i>FEBS Letters</i> , 1985, 187, 219-223.	1.3	15
328	The Proof by ¹³ C-NMR Spectroscopy of the Predominance of the C5 Pathway over the Shemin Pathway in Chlorophyll Biosynthesis in Higher Plants and of the Formation of the Methyl Ester Group of Chlorophyll from Glycine. <i>FEBS Journal</i> , 1983, 130, 509-516.	0.2	58
329	Assignment of heme and distal amino acid resonances in the proton NMR spectra of the oxygen and carbon monoxide complexes of soybean leghemoglobin. <i>BBA - Proteins and Proteomics</i> , 1983, 744, 281-290.	2.1	14
330	NMR studies of oxyleghemoglobin. <i>BBA - Proteins and Proteomics</i> , 1983, 749, 281-288.	2.1	12
331	A two-dimensional NMR method for assignment of imidazole ring proton resonances of histidine residues in proteins. <i>Biochemical and Biophysical Research Communications</i> , 1982, 106, 559-565.	1.0	20
332	Differences in the heme environment of soybean leghemoglobin components shown by ¹ H-NMR spectroscopy. <i>BBA - Proteins and Proteomics</i> , 1982, 700, 171-177.	2.1	7
333	Measurement of heme accessibility in soybean ferric leghemoglobin a and its complexes by proton magnetic relaxation. <i>Biochemistry</i> , 1981, 20, 587-594.	1.2	26
334	¹ H-NMR studies of ferric soybean leghemoglobin Assignment of hyperfine shifted resonances of complexes with cyanide, nicotinate, pyridine and azide. <i>Biochimica Et Biophysica Acta (BBA) - Protein Structure</i> , 1980, 625, 202-220.	1.7	27
335	Electron transfer reagent binding sites on plastocyanin. <i>Nature</i> , 1980, 283, 682-683.	13.7	48
336	NMR study of the interaction of plastocyanin with chromium(III) analogues of inorganic electron transfer reagents. <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 1980, 591, 162-176.	0.5	83
337	Molecular basis for proton-dependent anion binding by soybean leghaemoglobin a. <i>Nature</i> , 1979, 280, 87-88.	13.7	23
338	Conformational studies of peroxidase-substrate complexes. Structure of the indolepropionic acid-horseradish peroxidase complex. <i>Journal of the Chemical Society Chemical Communications</i> , 1975, ,	2.0	38
339	E.s.r. and optical spectral properties of copper(II) complexes with Schiff-base ligands derived from o-aminobenzaldehyde. <i>Journal of the Chemical Society Dalton Transactions</i> , 1975, , 867.	1.1	13
340	Conformational influences in copper co-ordination compounds. Part V. Crystal and molecular structure of {1,2-bis-[(2-aminobenzylidene)-amino]propanato(2-)}copper(II). <i>Journal of the Chemical Society Dalton Transactions</i> , 1973, , 1508-1512.	1.1	13
341	Structural and dynamic studies of DNA recognition by NF- κ B p50 RHR homodimer using methyl NMR spectroscopy. <i>Nucleic Acids Research</i> , 0, ,	6.5	1