

Marko Lucijanic

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3052380/marko-lucijanic-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

110
papers

368
citations

9
h-index

13
g-index

118
ext. papers

559
ext. citations

5.2
avg, IF

4.43
L-index

#	Paper	IF	Citations
110	Clinical and prognostic significance of C-reactive protein to albumin ratio in hospitalized coronavirus disease 2019 (COVID-19) patients : Data on 2309 patients from a tertiary center and validation in an independent cohort.. <i>Wiener Klinische Wochenschrift</i> , 2022 , 1	2.3	4
109	Remdesivir associated bradycardia might be a sign of good prognosis in COVID-19 patients.. <i>Clinical Microbiology and Infection</i> , 2022 ,	9.5	0
108	Baseline and progressive adipopenia in newly diagnosed patients with diffuse large B-cell lymphoma with unfavorable features are associated with worse clinical outcomes.. <i>Leukemia and Lymphoma</i> , 2022 , 1-10	1.9	0
107	Vaccination provides protection from respiratory deterioration and death among hospitalized COVID-19 patients: differences between vector and mRNA vaccines.. <i>Journal of Medical Virology</i> , 2022 ,	19.7	2
106	Higher ferritin levels in COVID-19 patients are associated with hyperinflammation, worse prognosis, and more bacterial infections without pronounced features of hemophagocytosis.. <i>Annals of Hematology</i> , 2022 , 101, 1119	3	0
105	High platelet-to-lymphocyte ratio may differentiate polycythemia vera from secondary polycythemia.. <i>Wiener Klinische Wochenschrift</i> , 2022 , 1	2.3	0
104	Serum osmolarity in polycythemia vera and its clinical associations.. <i>International Journal of Laboratory Hematology</i> , 2022 ,	2.5	
103	Red cell distribution width is a potent prognostic parameter for in-hospital and post-discharge mortality in hospitalized coronavirus disease 2019 patients: a registry-based cohort study on 3941 patients. <i>Croatian Medical Journal</i> , 2022 , 63, 44-52	1.6	3
102	Gradual increase in RDW is similarly prognostic for in-hospital mortality in both anemic and non-anemic COVID-19 patients.. <i>Journal of Medical Virology</i> , 2022 ,	19.7	
101	Comparison of IMPROVE, modified IMPROVE, IMPROVEDD, Padua and CHA2DS2-VASC risk scores for venous and arterial thrombotic events prediction in hospitalized COVID-19 patients.. <i>Thrombosis Research</i> , 2022 , 214, 37-39	8.2	0
100	Incidence and risk factors for venous and arterial thromboses in hospitalized patients with coronavirus disease 2019: data on 4014 patients from a tertiary center registry.. <i>Croatian Medical Journal</i> , 2022 , 63, 16-26	1.6	1
99	The associations of age, sex, and comorbidities with survival of hospitalized patients with coronavirus disease 2019: data from 4014 patients from a tertiary-center registry.. <i>Croatian Medical Journal</i> , 2022 , 63, 36-43	1.6	1
98	Red cell distribution width is a potent prognostic parameter for in-hospital and post-discharge mortality in hospitalized coronavirus disease 2019 patients: a registry-based cohort study on 3941 patients.. <i>Croatian Medical Journal</i> , 2022 , 63, 44-52	1.6	2
97	Atrial fibrillation in COVID-19 patients receiving remdesivir, matched case-control analysis.. <i>American Journal of Emergency Medicine</i> , 2022 ,	2.9	0
96	Remdesivir use in COVID-19 patients might predispose bacteremia, matched case-control analysis.. <i>Journal of Infection</i> , 2022 ,	18.9	2
95	Neutrophil-to-lymphocyte and platelet-to-lymphocyte ratios as prognostic biomarkers in polycythemia vera.. <i>International Journal of Laboratory Hematology</i> , 2021 ,	2.5	2
94	Bradycardia during remdesivir treatment might be associated with improved survival in patients with COVID-19: a retrospective cohort study on 473 patients from a tertiary centre. <i>Postgraduate Medical Journal</i> , 2021 ,	2	3

93	Pharmacists' interventions improve health-related quality of life of rural older person on warfarin: a randomized controlled trial. <i>Scientific Reports</i> , 2021 , 11, 21897	4.9	3
92	Hematocrit to hemoglobin ratio as a prognostic marker in polycythemia vera. <i>Wiener Klinische Wochenschrift</i> , 2021 , 1	2.3	0
91	Thrombotic risk in secondary polycythemia resembles low-risk polycythemia vera and increases in specific subsets of patients. <i>Thrombosis Research</i> , 2021 , 209, 47-50	8.2	1
90	An update to the custom-made MS Excel workbook performing the log-rank test with extended functionality and a new original COVID-19 training data set. <i>Croatian Medical Journal</i> , 2021 , 62, 531-533	1.6	2
89	Higher red blood cell distribution width predicts thrombosis risk in primary and secondary myelofibrosis. <i>Annals of Hematology</i> , 2021 , 1	3	3
88	Serologic Responses Following Sars-Cov-2 Vaccination and Factors Influencing It in Patients with Hematological Malignant and Non-Malignant Diseases - a Prospective Multicentric Study. <i>Blood</i> , 2021 , 138, 47-47	2.2	0
87	Asymptomatic deep vein thromboses in prolonged hospitalized COVID-19 patients. <i>Wiener Klinische Wochenschrift</i> , 2021 , 133, 1281	2.3	2
86	Thrombocytosis in COVID-19 patients without myeloproliferative neoplasms is associated with better prognosis but higher rate of venous thromboembolism. <i>Blood Cancer Journal</i> , 2021 , 11, 189	7	7
85	Ruxolitinib withdrawal due to the COVID-19. <i>Leukemia</i> , 2021 , 35, 1218	10.7	1
84	The Risk of Falling and Consequences of Falling in Patients with Atrial Fibrillation Receiving Different Types of Anticoagulant. <i>Drugs and Aging</i> , 2021 , 38, 417-425	4.7	2
83	Prognostic impact of psoas muscle index in patients with diffuse large B-cell lymphoma might be dependent on the immunochemotherapy type. <i>Leukemia and Lymphoma</i> , 2021 , 62, 2535-2538	1.9	2
82	Patients with dementia and atrial fibrillation less frequently receive direct oral anticoagulants (DOACs) and experience higher thrombotic and mortality risk. <i>Acta Clinica Belgica</i> , 2021 , 1-6	1.8	1
81	Psoas muscle index at the time of diagnosis might reflect the prognosis of classical Hodgkin's lymphoma patients. <i>Wiener Klinische Wochenschrift</i> , 2021 , 1	2.3	4
80	Perirenal and subcutaneous fat differently affect outcomes in newly diagnosed classical Hodgkin lymphoma patients. <i>Hematological Oncology</i> , 2021 , 39, 575-579	1.3	1
79	Remdesivir might induce changes in electrocardiogram beyond bradycardia in patients with coronavirus disease 2019-The pilot study. <i>Journal of Medical Virology</i> , 2021 , 93, 5724-5725	19.7	7
78	Once-Weekly Semaglutide in Adults with Overweight or Obesity. <i>New England Journal of Medicine</i> , 2021 , 385, e4	59.2	4
77	Ruxolitinib treatment improves muscle mass in patients with myelofibrosis. <i>Annals of Hematology</i> , 2021 , 100, 1105-1106	3	4
76	Differences in anticoagulation pattern, thrombotic and mortality risk in patients with atrial fibrillation and mid-range ejection fraction. <i>Cardiologia Croatica</i> , 2021 , 16, 59-59	0	

75	Low-density lipoprotein (LDL) and the risk of thrombotic events in essential thrombocythemia and polycythemia vera. <i>Annals of Hematology</i> , 2021 , 100, 1335-1336	3	1
74	Does body mass index influence clinical outcomes in patients with atrial fibrillation who receive direct oral anticoagulant therapy. <i>Cardiologia Croatica</i> , 2021 , 16, 67-67	0	
73	Large single center registry of anticoagulation therapy in atrial fibrillation: who is protected among a thousand patients?. <i>Cardiologia Croatica</i> , 2021 , 16, 58-58	0	
72	Modifying the Effects of Cytoreductive Drugs on Kidney Function in Myeloproliferative Neoplasms. <i>Pharmacology</i> , 2021 , 106, 451-452	2.3	0
71	Patients with dementia and atrial fibrillation are less likely to receive direct oral anticoagulants. <i>Cardiologia Croatica</i> , 2021 , 16, 60-60	0	
70	No Impact of the Increased Number of Cardiovascular Risk Factors on Thrombosis and Survival in Polycythemia Vera. <i>Oncology Research and Treatment</i> , 2021 , 44, 201-203	2.8	2
69	Higher serum uric acid is associated with higher risks of thrombosis and death in patients with primary myelofibrosis. <i>Wiener Klinische Wochenschrift</i> , 2021 , 1	2.3	3
68	Beneficial effect of ACE inhibitors on kidney function in polycythemia vera. <i>Wiener Klinische Wochenschrift</i> , 2021 , 133, 808-815	2.3	1
67	Changes of recommended anticoagulation therapy in patients with atrial fibrillation and high thrombotic risk: long-term follow-up data from two hospital centers. <i>Expert Review of Cardiovascular Therapy</i> , 2021 , 19, 857-863	2.5	0
66	Prevalence and Prognostic Impact of Deranged Liver Blood Tests in COVID-19: Experience from the Regional COVID-19 Center over the Cohort of 3812 Hospitalized Patients. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	2
65	FibroScan-AST Score Predicts 30-Day Mortality or Need for Mechanical Ventilation among Patients Hospitalized with COVID-19. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	2
64	More Pronounced Muscle Loss During Immunochemotherapy is Associated with Worse Clinical Outcomes in Newly Diagnosed Patients with Diffuse Large B-Cell Lymphoma with Unfavorable Features. <i>Therapeutics and Clinical Risk Management</i> , 2021 , 17, 1037-1044	2.9	1
63	The risk of falling and falling consequences in patients with atrial fibrillation receiving different types of anticoagulant drugs. <i>Cardiologia Croatica</i> , 2021 , 16, 61-61	0	
62	Needlestick and sharp injuries among healthcare workers prior to and during the coronavirus disease 2019 (COVID-19) pandemic.. <i>Infection Control and Hospital Epidemiology</i> , 2021 , 1-2	2	1
61	On interactions between body mass index, type of anticoagulation and outcomes. <i>International Journal of Cardiology</i> , 2020 , 312, 107	3.2	
60	Progressive multifocal leukoencephalopathy during rituximab maintenance after rituximab and bendamustine treatment for relapsed follicular lymphoma. <i>Annals of Hematology</i> , 2020 , 99, 2191-2192	3	0
59	Effect of age and comorbidities on the decline of glomerular filtration rate in chronic myeloid leukemia patients discontinuing tyrosine kinase inhibitors. <i>International Journal of Hematology</i> , 2020 , 112, 264	2.3	
58	Patients with atrial fibrillation and mid-range ejection fraction differ in anticoagulation pattern, thrombotic and mortality risk independently of CHADS-VAS score. <i>Heart and Vessels</i> , 2020 , 35, 1243-1249 ¹	3.1	3

57	Hyperuricemia might promote thrombosis in essential thrombocythemia and polycythemia vera. <i>Leukemia and Lymphoma</i> , 2020 , 61, 1744-1747	1.9	3
56	Association of Early Palliative Care With Survival in Patients With Advanced Lung Cancer. <i>JAMA Oncology</i> , 2020 , 6, 442-443	13.4	1
55	Comment on Nauck et al. Effects of Liraglutide Compared With Placebo on Events of Acute Gallbladder or Biliary Disease in Patients With Type 2 Diabetes at High Risk for Cardiovascular Events in the LEADER Randomized Trial. <i>Diabetes Care</i> 2019;42:1912-1920. <i>Diabetes Care</i> , 2020 , 43, e28-e29	14.6	
54	The effect of smoking on disease activity in rheumatoid arthritis - our experience. <i>Acta Clinica Croatica</i> , 2020 , 59, 312-317	0.8	
53	Patterns of anticoagulation therapy in atrial fibrillation: results from a large real-life single-center registry. <i>Croatian Medical Journal</i> , 2020 , 61, 440-449	1.6	4
52	A validation of the Croatian version of Zarit Burden Interview and clinical predictors of caregiver burden in informal caregivers of patients with dementia: a cross-sectional study. <i>Croatian Medical Journal</i> , 2020 , 61, 527-537	1.6	4
51	CORRELATION BETWEEN THE LEVEL OF COLORECTAL ANASTOMOSIS AND ANORECTAL FUNCTION. <i>Acta Clinica Croatica</i> , 2020 , 59, 703-711	0.8	
50	Patients with higher body mass index treated with direct / novel oral anticoagulants (DOAC / NOAC) for atrial fibrillation experience worse clinical outcomes. <i>International Journal of Cardiology</i> , 2020 , 301, 90-95	3.2	16
49	Neutral effect of Glioma-associated oncogene-1 expression on survival in myelofibrosis. <i>Wiener Klinische Wochenschrift</i> , 2020 , 132, 464-466	2.3	1
48	Prolonged methylprednisolone premedication prior to obinutuzumab in patients with chronic lymphocytic leukemia. <i>Leukemia and Lymphoma</i> , 2020 , 61, 934-939	1.9	3
47	Higher AURKA and PLK1 expression are associated with inferior overall survival in patients with myelofibrosis. <i>Blood Cells, Molecules, and Diseases</i> , 2020 , 81, 102396	2.1	3
46	Dyslipidaemia, a factor worthy of adjustment. <i>European Journal of Heart Failure</i> , 2020 , 22, 564	12.3	
45	C reactive protein to albumin ratio as prognostic marker in primary and secondary myelofibrosis. <i>Leukemia and Lymphoma</i> , 2020 , 61, 2969-2974	1.9	3
44	Iron Chelation in Transfusion-Dependent Patients With Low- to Intermediate-1-Risk Myelodysplastic Syndromes. <i>Annals of Internal Medicine</i> , 2020 , 173, 595	8	1
43	CDC25c expression in patients with myelofibrosis is associated with stronger myeloproliferation and shorter overall survival. <i>Wiener Klinische Wochenschrift</i> , 2020 , 1	2.3	
42	Reduced renal function strongly affects survival and thrombosis in patients with myelofibrosis. <i>Annals of Hematology</i> , 2020 , 99, 2779-2785	3	9
41	The possible role of insulin and glucagon in patients with heart failure and Type 2 diabetes. <i>European Heart Journal</i> , 2020 , 41, 325	9.5	4
40	Dietary Supplement Use and Patient Outcomes in High-Risk Early-Stage Breast Cancer. <i>Journal of Clinical Oncology</i> , 2020 , 38, 2110	2.2	3

39	Pharmacists' Influence on adverse reactions to warfarin: a randomised controlled trial in elderly rural patients. <i>International Journal of Clinical Pharmacy</i> , 2019 , 41, 1166-1173	2.3	6
38	DIFFERENTIATED THYROID CANCER IN PEDIATRIC POPULATION (18 YEARS): POSTOPERATIVE TREATMENT WITH RADIOACTIVE IODINE (I-131). <i>Acta Clinica Croatica</i> , 2019 , 58, 119-127	0.8	1
37	Catheter Ablation Compared With Drug Therapy for Atrial Fibrillation. <i>JAMA - Journal of the American Medical Association</i> , 2019 , 322, 1105	27.4	
36	Longer Survival With Anti-Programmed Cell Death 1 and Development of Cutaneous Toxic Effects, an Expected Association. <i>JAMA Oncology</i> , 2019 , 5, 1642	13.4	
35	Aspirin Use and the Risk of Cancer. <i>JAMA Oncology</i> , 2019 , 5, 912-913	13.4	5
34	Could haemochromatosis (HFE) gene mutations affect response to iron chelation in myelodysplastic syndrome?. <i>British Journal of Haematology</i> , 2019 , 186, 640-641	4.5	
33	Platelet Counts and Risk of Major Bleeding With Ibrutinib. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2019 , 19, e399	2	3
32	Above What Age Should Ibrutinib Not Be Given With R-CHOP to Patients With Non-GBC DLBCL?. <i>Journal of Clinical Oncology</i> , 2019 , 37, 2583-2584	2.2	2
31	Second primary malignancies in myeloproliferative neoplasms and the role of aspirin. <i>Leukemia</i> , 2019 , 33, 2554	10.7	1
30	Liraglutide in Children and Teens with Type 2 Diabetes. <i>New England Journal of Medicine</i> , 2019 , 381, 1786-1787	17.4	
29	Epidemiological and clinical features of primary biliary cholangitis in two Croatian regions: a retrospective study. <i>Croatian Medical Journal</i> , 2019 , 60, 494-502	1.6	6
28	Patterns of anticoagulation and clinical outcomes in atrial fibrillation: insights from a single institution registry. <i>Cardiologia Croatica</i> , 2019 , 14, 239-239	0	
27	Bendamustine-based conditioning prior to autologous stem cell transplantation is associated with high rate of febrile neutropenia and higher mortality. <i>American Journal of Hematology</i> , 2019 , 94, E42-E43	7.1	2
26	Progressive multifocal leukoencephalopathy developing after obinutuzumab treatment for chronic lymphocytic leukemia. <i>Annals of Hematology</i> , 2019 , 98, 1509-1510	3	8
25	Prognostic implications of low transferrin saturation in patients with primary myelofibrosis. <i>Leukemia Research</i> , 2018 , 66, 89-95	2.7	4
24	Assessing serum albumin concentration, lymphocyte count and prognostic nutritional index might improve prognostication in patients with myelofibrosis. <i>Wiener Klinische Wochenschrift</i> , 2018 , 130, 126-133	2.3	13
23	High red cell distribution width at the time of ST segment elevation myocardial infarction is better at predicting diastolic than systolic left ventricular dysfunction: A single-center prospective cohort study. <i>Medicine (United States)</i> , 2018 , 97, e0601	1.8	4
22	Higher Sclerostin/SOST expression is associated with lower percentage of circulatory blasts and better prognosis in patients with myelofibrosis. <i>Annals of Hematology</i> , 2018 , 97, 1293-1294	3	

21	Increased mean platelet volume (MPV) is an independent predictor of inferior survival in patients with primary and secondary myelofibrosis. <i>International Journal of Hematology</i> , 2018 , 107, 166-172	2.3	6
20	High absolute basophil count is a powerful independent predictor of inferior overall survival in patients with primary myelofibrosis. <i>Hematology</i> , 2018 , 23, 201-207	2.2	4
19	Risk stratification according to insulin type. <i>European Journal of Heart Failure</i> , 2018 , 20, 1497	12.3	2
18	Pharmacist ^B interventions improve time in therapeutic range of elderly rural patients on warfarin therapy: a randomized trial. <i>International Journal of Clinical Pharmacy</i> , 2018 , 40, 1078-1085	2.3	12
17	Ultrasound Grade of Liver Steatosis Is Independently Associated with the Risk of Metabolic Syndrome. <i>Canadian Journal of Gastroenterology and Hepatology</i> , 2018 , 2018, 8490242	2.8	13
16	Combining information on C reactive protein and serum albumin into the Glasgow Prognostic Score strongly discriminates survival of myelofibrosis patients. <i>Blood Cells, Molecules, and Diseases</i> , 2018 , 72, 14-16	2.1	3
15	Elevated Neutrophil-to-Lymphocyte-ratio and Platelet-to-Lymphocyte Ratio in Myelofibrosis: Inflammatory Biomarkers or Representatives of Myeloproliferation Itself?. <i>Anticancer Research</i> , 2018 , 38, 3157-3163	2.3	6
14	THE ASSOCIATION OF LOW TRIIODOTHYRONINE SYNDROME, PROTEIN-ENERGY WASTING AND CHRONIC INFLAMMATION WITH MORTALITY IN PREVALENT DIALYSIS PATIENTS. <i>Acta Clinica Croatica</i> , 2018 , 57, 449-457	0.8	
13	Impact of pharmacotherapeutic education on medication adherence and adverse outcomes in patients with type 2 diabetes mellitus: a prospective, randomized study. <i>Croatian Medical Journal</i> , 2018 , 59, 290-297	1.6	6
12	The role of glucagon in the possible mechanism of cardiovascular mortality reduction in type 2 diabetes patients. <i>International Journal of Clinical Practice</i> , 2018 , 72, e13274	2.9	5
11	Heat shock protein 27 (HSP27/HSPB1) expression is increased in patients with primary and secondary myelofibrosis and may be affecting their survival. <i>Leukemia and Lymphoma</i> , 2017 , 58, 2497-2508	1.8	6
10	Clinical and prognostic significance of anisocytosis measured as a red cell distribution width in patients with colorectal cancer. <i>QJM - Monthly Journal of the Association of Physicians</i> , 2017 , 110, 361-367	2.7	17
9	Comment to: "Management and outcome of gastrointestinal bleeding in patients taking oral anticoagulants or antiplatelet drugs". <i>Journal of Gastroenterology</i> , 2017 , 52, 1075-1076	6.9	1
8	Rituximab with dose-adjusted EPOCH as first-line treatment in patients with highly aggressive diffuse large B-cell lymphoma and autologous stem cell transplantation in selected patients. <i>Croatian Medical Journal</i> , 2017 , 58, 40-48	1.6	10
7	Survival analysis, more than meets the eye. <i>Biochemia Medica</i> , 2017 , 27, 14-18	2.5	14
6	Correlation between red cell distribution width, systolic function and atrial fibrillation occurrence in patients with acute ST segment elevation myocardial infarction. <i>Cardiologia Croatica</i> , 2017 , 12, 346-348	0	
5	Survival analysis in clinical practice: analyze your own data using an Excel workbook. <i>Croatian Medical Journal</i> , 2016 , 57, 77-9	1.6	32
4	Canonical Wnt/ECatenin Signaling Pathway Is Dysregulated in Patients With Primary and Secondary Myelofibrosis. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2016 , 16, 523-526	2	6

3	Hemochromatosis gene mutations may affect the survival of patients with myelodysplastic syndrome. <i>Hematology</i> , 2016 , 21, 170-4	2.2	5
2	The Degree of Anisocytosis Predicts Survival in Patients with Primary Myelofibrosis. <i>Acta Haematologica</i> , 2016 , 136, 98-100	2.7	12
1	Analysis of censored data. <i>Biochemia Medica</i> , 2012 , 22, 151-5	2.5	8