

Gerald S Treiman

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3054339/publications.pdf>

Version: 2024-02-01

13
papers

310
citations

933447

10
h-index

1125743

13
g-index

13
all docs

13
docs citations

13
times ranked

519
citing authors

#	ARTICLE	IF	CITATIONS
1	Intraluminal Thrombus, Intraplaque Hemorrhage, Plaque Thickness, and Current Smoking Optimally Predict Carotid Stroke. <i>Stroke</i> , 2015, 46, 84-90.	2.0	69
2	Prediction of Carotid Intraplaque Hemorrhage Using Adventitial Calcification and Plaque Thickness on CTA. <i>American Journal of Neuroradiology</i> , 2016, 37, 1496-1503.	2.4	46
3	Carotid MRI Detection of Intraplaque Hemorrhage at 3T and 1.5T. <i>Journal of Neuroimaging</i> , 2015, 25, 390-396.	2.0	26
4	Carotid Magnetization-Prepared Rapid Acquisition With Gradient-Echo Signal Is Associated With Acute Territorial Cerebral Ischemic Events Detected by Diffusion-Weighted MRI. <i>Circulation: Cardiovascular Imaging</i> , 2012, 5, 376-382.	2.6	25
5	In vivo and ex vivo measurements of the mean ADC values of lipid necrotic core and hemorrhage obtained from diffusion weighted imaging in human atherosclerotic plaques. <i>Journal of Magnetic Resonance Imaging</i> , 2011, 34, 1167-1175.	3.4	24
6	Magnetic Resonance Imaging Detection of Intraplaque Hemorrhage. <i>Magnetic Resonance Insights</i> , 2017, 10, 1178623X1769415.	2.5	23
7	Magnetic Resonance Imaging and Clinical Factors Associated With Ischemic Stroke in Patients Suspected of Cervical Artery Dissection. <i>Stroke</i> , 2018, 49, 2337-2344.	2.0	22
8	Diffusion-weighted imaging of human carotid artery using 2D single-shot interleaved multislice inner volume diffusion-weighted echo planar imaging (2D ss-EPI) at 3T: Diffusion measurement in atherosclerotic plaque. <i>Journal of Magnetic Resonance Imaging</i> , 2009, 30, 1068-1077.	3.4	19
9	Correlation of Carotid Intraplaque Hemorrhage and Stroke Using 1.5 T and 3 T Mri. <i>Magnetic Resonance Insights</i> , 2015, 8s1, MRI.S23560.	2.5	18
10	Interchangeable neck shape-specific coils for a clinically realizable anterior neck phased array system. <i>Magnetic Resonance in Medicine</i> , 2017, 78, 2460-2468.	3.0	14
11	Motion-insensitive carotid intraplaque hemorrhage imaging using 3D inversion recovery preparation stack of stars (IR-prep SOS) technique. <i>Journal of Magnetic Resonance Imaging</i> , 2017, 45, 410-417.	3.4	12
12	Vitamin D and Vulnerable Carotid Plaque. <i>American Journal of Neuroradiology</i> , 2016, 37, 2092-2099.	2.4	9
13	Differentiation of symptomatic and asymptomatic carotid intraplaque hemorrhage using 3D high-resolution diffusion-weighted stack of stars imaging. <i>NMR in Biomedicine</i> , 2021, 34, e4582.	2.8	3