

Fabiano Elias Xavier

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3059866/fabiano-elias-xavier-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

9

papers

99

citations

5

h-index

9

g-index

9

ext. papers

112

ext. citations

3.7

avg, IF

1.6

L-index

#	Paper	IF	Citations
9	Chronic cyclooxygenase-2 inhibition prevents the worsening of hypertension and endothelial dysfunction induced by ouabain in resistance arteries of spontaneously hypertensive rats. <i>Vascular Pharmacology</i> , 2021 , 139, 106880	5.9	0
8	Mechanisms underlying the vasorelaxant effect of trans-4-methoxy- β -nitrostyrene in the rat mesenteric resistance arteries. <i>European Journal of Pharmacology</i> , 2019 , 853, 201-209	5.3	5
7	Hyperglycaemia in pregnant rats causes sex-related vascular dysfunction in adult offspring: role of cyclooxygenase-2. <i>Experimental Physiology</i> , 2017 , 102, 1019-1036	2.4	9
6	Losartan reverses COX-2-dependent vascular dysfunction in offspring of hyperglycaemic rats. <i>Life Sciences</i> , 2017 , 184, 71-80	6.8	5
5	Enhanced Na ⁺ , K ⁺ -ATPase activity and endothelial modulation decrease phenylephrine-induced contraction in aorta from ouabain-treated normotensive and hypertensive rats. <i>Hormone Molecular Biology and Clinical Investigation</i> , 2014 , 18, 113-22	1.3	2
4	Vasorelaxant effects of 1-nitro-2-phenylethane, the main constituent of the essential oil of Aniba canelilla, in superior mesenteric arteries from spontaneously hypertensive rats. <i>European Journal of Pharmaceutical Sciences</i> , 2013 , 48, 709-16	5.1	21
3	Cardiovascular effects of 1-nitro-2-phenylethane, the main constituent of the essential oil of Aniba canelilla, in spontaneously hypertensive rats. <i>Fundamental and Clinical Pharmacology</i> , 2011 , 25, 661-9	3.1	21
2	Long-term ouabain treatment impairs vascular function in resistance arteries. <i>Journal of Vascular Research</i> , 2011 , 48, 316-26	1.9	19
1	Ouabain at nanomolar concentration promotes synthesis and release of angiotensin II from the endothelium of the tail vascular bed of spontaneously hypertensive rats. <i>Journal of Cardiovascular Pharmacology</i> , 2004 , 44, 372-80	3.1	17