

# Carolina Bravo

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/306275/publications.pdf>

Version: 2024-02-01

20  
papers

385  
citations

840776

11  
h-index

794594

19  
g-index

21  
all docs

21  
docs citations

21  
times ranked

505  
citing authors

#	ARTICLE	IF	CITATIONS
1	Detectability and predator strategy affect egg depredation rates: Implications for mitigating nest depredation in farmlands. <i>Science of the Total Environment</i> , 2022, 829, 154558.	8.0	8
2	Burrowing Parrots <i>Cyanoliseus patagonus</i> as Long-Distance Seed Dispersers of Keystone Algarrobos, Genus <i>Prosopis</i> , in the Monte Desert. <i>Diversity</i> , 2021, 13, 204.	1.7	7
3	Herb endozoochory by cockatoos: Is "foliage the fruit"? <i>Austral Ecology</i> , 2020, 45, 122-126.	1.5	11
4	Physiological dormancy broken by endozoochory: Austral parakeets ( <i>Enicognathus ferrugineus</i> ) as legitimate dispersers of calafate ( <i>Berberis microphylla</i> ) in the Patagonian Andes. <i>Journal of Plant Ecology</i> , 2020, 13, 538-544.	2.3	13
5	Revisiting an old question: Which predators eat eggs of ground-nesting birds in farmland landscapes?. <i>Science of the Total Environment</i> , 2020, 744, 140895.	8.0	16
6	Feeding functional responses in a sexually size-dimorphic bird. <i>Acta Oecologica</i> , 2019, 101, 103487.	1.1	5
7	Effects of farming practices on nesting success of steppe birds in dry cereal farmland. <i>European Journal of Wildlife Research</i> , 2018, 64, 1.	1.4	18
8	Food Availability But Not Sex Determines Morning Foraging Area Size in the Great Bustard <i>Otis tarda</i> , the Most Sexually Size-Dimorphic Bird Species. <i>Ardeola</i> , 2017, 64, 289.	0.7	6
9	Cantharidin is conserved across phylogeographic lineages and present in both morphs of Iberian <i>Berberomeloe blister</i> beetles (Coleoptera, Meloidae). <i>Zoological Journal of the Linnean Society</i> , 2017, 180, 790-804.	2.3	15
10	Diet composition of a declining steppe bird the Little Bustard ( <i>Tetrax tetrax</i> ) in relation to farming practices. <i>Avian Conservation and Ecology</i> , 2017, 12, .	0.8	14
11	Dietary divergence in the most sexually size-dimorphic bird. <i>Auk</i> , 2016, 133, 178-197.	1.4	19
12	Internal seed dispersal by parrots: an overview of a neglected mutualism. <i>PeerJ</i> , 2016, 4, e1688.	2.0	51
13	Males of a Strongly Polygynous Species Consume More Poisonous Food than Females. <i>PLoS ONE</i> , 2014, 9, e111057.	2.5	27
14	Effects of agricultural environmental schemes on farmland birds: do food availability measurements improve patterns obtained from simple habitat models?. <i>Ecology and Evolution</i> , 2014, 4, 2834-2847.	1.9	25
15	Effects of great bustard ( <i>Otis tarda</i> ) gut passage on black nightshade ( <i>Solanum nigrum</i> ) seed germination. <i>Seed Science Research</i> , 2014, 24, 265-271.	1.7	12
16	Habitat preferences of sympatric sandgrouse during the breeding season in Spain: a multi-scale approach. <i>European Journal of Wildlife Research</i> , 2014, 60, 625-636.	1.4	4
17	Faecal sexual steroids in sex typing and endocrine status of great bustards. <i>European Journal of Wildlife Research</i> , 2013, 59, 815-822.	1.4	16
18	Diet of young Great Bustards <i>Otis tarda</i> in Spain: sexual and seasonal differences. <i>Bird Study</i> , 2012, 59, 243-251.	1.0	24

#	ARTICLE	IF	CITATIONS
19	Side effects of rodent control on non-target species: Rodenticides increase parasite and pathogen burden in great bustards. <i>Science of the Total Environment</i> , 2011, 409, 4729-4734.	8.0	34
20	Effects of organic farming on plant and arthropod communities: A case study in Mediterranean dryland cereal. <i>Agriculture, Ecosystems and Environment</i> , 2011, 141, 193-201.	5.3	58