

Henis Yi

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3069712/henis-yi-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

49
papers

2,772
citations

25
h-index

50
g-index

50
ext. papers

3,011
ext. citations

5
avg, IF

4.65
L-index

#	Paper	IF	Citations
49	Competition between type I activin and BMP receptors for binding to ACVR2A regulates signaling to distinct Smad pathways.. <i>BMC Biology</i> , 2022 , 20, 50	7.3	0
48	Complex Formation Among TGF- β Receptors in Live Cell Membranes Measured by Patch-FRAP.. <i>Methods in Molecular Biology</i> , 2022 , 2488, 23-34	1.4	
47	Autophagy is induced and modulated by cholesterol depletion through transcription of autophagy-related genes and attenuation of flux. <i>Cell Death Discovery</i> , 2021 , 7, 320	6.9	1
46	ALK1 regulates the internalization of endoglin and the type III TGF- β receptor. <i>Molecular Biology of the Cell</i> , 2021 , 32, 605-621	3.5	1
45	Formation of self-organizing functionally distinct Rho of plants domains involves a reduced mobile population. <i>Plant Physiology</i> , 2021 , 187, 2485-2508	6.6	1
44	Ras Diffusion and Interactions with the Plasma Membrane Measured by FRAP Variations. <i>Methods in Molecular Biology</i> , 2021 , 2262, 185-197	1.4	
43	Noncatalytic Bruton's tyrosine kinase activates PLC γ variants mediating ibrutinib resistance in human chronic lymphocytic leukemia cells. <i>Journal of Biological Chemistry</i> , 2020 , 295, 5717-5736	5.4	16
42	Phase separation of TAZ compartmentalizes the transcription machinery to promote gene expression. <i>Nature Cell Biology</i> , 2020 , 22, 453-464	23.4	95
41	Zeb2 regulates the balance between retinal interneurons and Müller glia by inhibition of BMP-Smad signaling. <i>Developmental Biology</i> , 2020 , 468, 80-92	3.1	3
40	Cholesterol depletion enhances TGF- β Smad signaling by increasing c-Jun expression through a PKR-dependent mechanism. <i>Molecular Biology of the Cell</i> , 2018 , 29, 2494-2507	3.5	7
39	The residue at position 5 of the N-terminal region of Src and Fyn modulates their myristoylation, palmitoylation, and membrane interactions. <i>Molecular Biology of the Cell</i> , 2016 , 27, 3926-3936	3.5	9
38	Dynamin-dependent endocytosis of Bone Morphogenetic Protein2 (BMP2) and its receptors is dispensable for the initiation of Smad signaling. <i>International Journal of Biochemistry and Cell Biology</i> , 2016 , 76, 51-63	5.6	10
37	Rac-mediated Stimulation of Phospholipase C β Amplifies B Cell Receptor-induced Calcium Signaling. <i>Journal of Biological Chemistry</i> , 2015 , 290, 17056-72	5.4	14
36	Regulation of TGF- β receptor hetero-oligomerization and signaling by endoglin. <i>Molecular Biology of the Cell</i> , 2015 , 26, 3117-27	3.5	28
35	T β RIII independently binds type I and type II TGF- β receptors to inhibit TGF- β signaling. <i>Molecular Biology of the Cell</i> , 2015 , 26, 3535-45	3.5	20
34	Oligomeric interactions of TGF- β and BMP receptors. <i>FEBS Letters</i> , 2012 , 586, 1885-96	3.8	55
33	Coated pit-mediated endocytosis of the type I transforming growth factor- β (TGF- β) receptor depends on a di-leucine family signal and is not required for signaling. <i>Journal of Biological Chemistry</i> , 2012 , 287, 26876-89	5.4	21

32	Accurate quantification of diffusion and binding kinetics of non-integral membrane proteins by FRAP. <i>Traffic</i> , 2011 , 12, 1648-57	5.7	22
31	Formation of stable homomeric and transient heteromeric bone morphogenetic protein (BMP) receptor complexes regulates Smad protein signaling. <i>Journal of Biological Chemistry</i> , 2011 , 286, 19287-96	5.4	25
30	Raft protein clustering alters N-Ras membrane interactions and activation pattern. <i>Molecular and Cellular Biology</i> , 2011 , 31, 3938-52	4.8	39
29	Differential effects of prenylation and s-acylation on type I and II ROPS membrane interaction and function. <i>Plant Physiology</i> , 2011 , 155, 706-20	6.6	53
28	Differential regulation of phospholipase C-beta2 activity and membrane interaction by Galphaq, Gbeta1gamma2, and Rac2. <i>Journal of Biological Chemistry</i> , 2010 , 285, 3905-3915	5.4	35
27	An S-acylation switch of conserved G domain cysteines is required for polarity signaling by ROP GTPases. <i>Current Biology</i> , 2010 , 20, 914-20	6.3	62
26	Different domains regulate homomeric and heteromeric complex formation among type I and type II transforming growth factor-beta receptors. <i>Journal of Biological Chemistry</i> , 2009 , 284, 7843-52	5.4	22
25	Ras acylation, compartmentalization and signaling nanoclusters (Review). <i>Molecular Membrane Biology</i> , 2009 , 26, 80-92	3.4	94
24	A role for the juxtamembrane cytoplasm in the molecular dynamics of focal adhesions. <i>PLoS ONE</i> , 2009 , 4, e4304	3.7	62
23	Differential interference of chlorpromazine with the membrane interactions of oncogenic K-Ras and its effects on cell growth. <i>Journal of Biological Chemistry</i> , 2008 , 283, 27279-88	5.4	24
22	Src kinase activity and SH2 domain regulate the dynamics of Src association with lipid and protein targets. <i>Journal of Cell Biology</i> , 2007 , 178, 675-86	7.3	48
21	Clustering of raft-associated proteins in the external membrane leaflet modulates internal leaflet H-ras diffusion and signaling. <i>Molecular and Cellular Biology</i> , 2006 , 26, 7190-200	4.8	62
20	Different routes of bone morphogenetic protein (BMP) receptor endocytosis influence BMP signaling. <i>Molecular and Cellular Biology</i> , 2006 , 26, 7791-805	4.8	194
19	FRAP beam-size analysis to measure palmitoylation-dependent membrane association dynamics and microdomain partitioning of Ras proteins. <i>Methods</i> , 2006 , 40, 183-90	4.6	46
18	Cyclodextrins but not compactin inhibit the lateral diffusion of membrane proteins independent of cholesterol. <i>Traffic</i> , 2006 , 7, 917-26	5.7	58
17	Individual palmitoyl residues serve distinct roles in H-ras trafficking, microlocalization, and signaling. <i>Molecular and Cellular Biology</i> , 2005 , 25, 6722-33	4.8	177
16	Pathway- and expression level-dependent effects of oncogenic N-Ras: p27(Kip1) mislocalization by the Ral-GEF pathway and Erk-mediated interference with Smad signaling. <i>Molecular and Cellular Biology</i> , 2005 , 25, 8239-50	4.8	50
15	Differently anchored influenza hemagglutinin mutants display distinct interaction dynamics with mutual rafts. <i>Journal of Cell Biology</i> , 2003 , 163, 879-88	7.3	98

14	Transforming growth factor-beta receptors interact with AP2 by direct binding to beta2 subunit. <i>Molecular Biology of the Cell</i> , 2002 , 13, 4001-12	3.5	106
13	Activated K-Ras and H-Ras display different interactions with saturable nonraft sites at the surface of live cells. <i>Journal of Cell Biology</i> , 2002 , 157, 865-72	7.3	197
12	The mode of bone morphogenetic protein (BMP) receptor oligomerization determines different BMP-2 signaling pathways. <i>Journal of Biological Chemistry</i> , 2002 , 277, 5330-8	5.4	415
11	Bone morphogenetic protein receptor complexes on the surface of live cells: a new oligomerization mode for serine/threonine kinase receptors. <i>Molecular Biology of the Cell</i> , 2000 , 11, 1023-35	3.5	245
10	Membrane interactions of a constitutively active GFP-Ki-Ras 4B and their role in signaling. Evidence from lateral mobility studies. <i>Journal of Biological Chemistry</i> , 1999 , 274, 1606-13	5.4	102
9	Transforming growth factor-beta induces formation of a dithiothreitol-resistant type I/Type II receptor complex in live cells. <i>Journal of Biological Chemistry</i> , 1999 , 274, 5716-22	5.4	51
8	Oligomeric structure of type I and type II transforming growth factor beta receptors: homodimers form in the ER and persist at the plasma membrane. <i>Journal of Cell Biology</i> , 1998 , 140, 767-77	7.3	128
7	Dynamic or stable interactions of influenza hemagglutinin mutants with coated pits. Dependence on the internalization signal but not on aggregation. <i>Journal of Biological Chemistry</i> , 1995 , 270, 21075-81	5.4	26
6	Mobility of enzymes on insoluble substrates: the beta-amylase-starch gel system. <i>Biopolymers</i> , 1988 , 27, 123-38	2.2	20
5	Interaction of Sendai virions with resealed human erythrocyte ghosts. Lateral mobility of the viral glycoproteins in the cell membrane following fusion. <i>FEBS Letters</i> , 1988 , 228, 281-4	3.8	1
4	Enzyme diffusion and action on soluble and insoluble substrate biopolymers. <i>Biopolymers</i> , 1985 , 24, 257-77	7.7	12
3	Detection of Sendai virus fusion with human erythrocytes by fluorescence photobleaching recovery. <i>FEBS Letters</i> , 1983 , 151, 134-8	3.8	8
2	Interactions between antagonist-occupied muscarinic binding sites in rat adenohypophysis. <i>FEBS Letters</i> , 1982 , 140, 173-6	3.8	7
1	The mechanism of negative cooperativity in rabbit muscle glyceraldehyde-3-phosphate dehydrogenase. <i>Annals of the New York Academy of Sciences</i> , 1981 , 366, 217-36	6.5	2