Naik Vietti Violi

List of Publications by Citations

Source: https://exaly.com/author-pdf/3071793/naik-vietti-violi-publications-by-citations.pdf

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

17
papers285
citations9
h-index16
g-index21
ext. papers405
ext. citations5.5
avg, IF3.08
L-index

#	Paper	IF	Citations
17	Efficacy of microwave ablation versus radiofrequency ablation for the treatment of hepatocellular carcinoma in patients with chronic liver disease: a randomised controlled phase 2 trial. <i>The Lancet Gastroenterology and Hepatology</i> , 2018 , 3, 317-325	18.8	116
16	Prevalence and clinical importance of mesenteric venous thrombosis in the Swiss Inflammatory Bowel Disease Cohort. <i>American Journal of Roentgenology</i> , 2014 , 203, 62-9	5.4	28
15	Hepatocellular carcinoma detection in liver cirrhosis: diagnostic performance of contrast-enhanced CT vs. MRI with extracellular contrast vs. gadoxetic acid. <i>European Radiology</i> , 2020 , 30, 1020-1030	8	23
14	Hepatocellular carcinoma abutting large vessels: comparison of four percutaneous ablation systems. <i>International Journal of Hyperthermia</i> , 2018 , 34, 1171-1178	3.7	19
13	Gadoxetate-enhanced abbreviated MRI is highly accurate for hepatocellular carcinoma screening. <i>European Radiology</i> , 2020 , 30, 6003-6013	8	17
12	Local recurrence rate in patients with colorectal cancer liver metastasis after wedge resection or percutaneous radiofrequency ablation. <i>International Journal of Hyperthermia</i> , 2018 , 34, 1020-1028	3.7	17
11	Gadoxetate-enhanced Abbreviated MRI for Hepatocellular Carcinoma Surveillance: Preliminary Experience. <i>Radiology Imaging Cancer</i> , 2019 , 1, e190010	1.4	15
10	Acute mesenteric vein thrombosis: factors associated with evolution to chronic mesenteric vein thrombosis. <i>American Journal of Roentgenology</i> , 2014 , 203, 54-61	5.4	14
9	Value of Tumor Growth Rate (TGR) as an Early Biomarker Predictor of PatientsdOutcome in Neuroendocrine Tumors (NET)-The GREPONET Study. <i>Oncologist</i> , 2019 , 24, e1082-e1090	5.7	13
8	Tumor Growth Rate as a Validated Early Radiological Biomarker Able to Reflect Treatment-Induced Changes in Neuroendocrine Tumors: The GREPONET-2 Study. <i>Clinical Cancer Research</i> , 2019 , 25, 6692-6	6 99 9	7
7	Tumor Growth Rate to Predict the Outcome of Patients with Neuroendocrine Tumors: Performance and Sources of Variability. <i>Neuroendocrinology</i> , 2021 , 111, 831-839	5.6	3
6	Sildenafil as a therapeutic option for digital ischemic ulceration: case report. <i>Journal of Hand Surgery</i> , 2015 , 40, 890-3	2.6	2
5	Gadoxetate disodium-enhanced MRI: Assessment of arterial phase artifacts and hepatobiliary uptake in a large series. <i>European Journal of Radiology</i> , 2020 , 132, 109313	4.7	2
4	Abbreviated Magnetic Resonance Imaging for HCC Surveillance. Clinical Liver Disease, 2021, 17, 133-138	3 2.2	2
3	Abbreviated Magnetic Resonance Imaging Protocols in the Abdomen and Pelvis. <i>Magnetic Resonance Imaging Clinics of North America</i> , 2020 , 28, 381-394	1.6	1
2	Radiofrequency Ablation to Control Bleeding from a Percutaneous Intrahepatic Puncture Tract of a Large Diameter Portal Vein: A Simple and Rapid Solution to a Potentially Life-Threatening Bleeding. <i>CardioVascular and Interventional Radiology</i> , 2020 , 43, 1725-1727	2.7	1
1	MRI is the most commonly used imaging modality for HCC screening at a tertiary care transplant center. <i>Abdominal Radiology</i> , 2021 , 46, 5142-5151	3	1