Jian Sun

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/3082016/jian-sun-publications-by-year.pdf

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

59	2,126	25	45
papers	citations	h-index	g-index
62	2,771 ext. citations	5	4.38
ext. papers		avg, IF	L-index

#	Paper	IF	Citations
59	Targeting of SPCSV-RNase3 via CRISPR-Cas13 confers resistance against sweet potato virus disease. <i>Molecular Plant Pathology</i> , 2022 , 23, 104-117	5.7	4
58	Damage Signaling by Extracellular Nucleotides: A Role for Cyclic Nucleotides in Elevating Cytosolic Free Calcium?. <i>Frontiers in Plant Science</i> , 2021 , 12, 788514	6.2	2
57	Identification of Shaker K channel family members in sweetpotato and functional exploration of IbAKT1. <i>Gene</i> , 2021 , 768, 145311	3.8	5
56	NaCl-altered oxygen flux profiles and H+-ATPase activity in roots of two contrasting poplar species. <i>Tree Physiology</i> , 2021 , 41, 756-770	4.2	
55	Annexin 1 Is a Component of eATP-Induced Cytosolic Calcium Elevation in Roots. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	7
54	Robust -Mediated Transient Expression in Two Duckweed Species (Lemnaceae) Directed by Non-replicating, Replicating, and Cell-to-Cell Spreading Vectors. <i>Frontiers in Bioengineering and Biotechnology</i> , 2021 , 9, 5	5.8	4
53	Genome-wide analysis of expression quantitative trait loci (eQTLs) reveals the regulatory architecture of gene expression variation in the storage roots of sweet potato. <i>Horticulture Research</i> , 2020 , 7, 90	7.7	17
52	GaHD1, a candidate gene for the Gossypium arboreum SMA-4 mutant, promotes trichome and fiber initiation by cellular HO and Ca signals. <i>Plant Molecular Biology</i> , 2020 , 103, 409-423	4.6	8
51	SISTE1 promotes abscisic acid-dependent salt stress-responsive pathways via improving ion homeostasis and reactive oxygen species scavenging in tomato. <i>Journal of Integrative Plant Biology</i> , 2020 , 62, 1942-1966	8.3	2
50	Characterization of a Novel Chitinase from Sweet Potato and Its Fungicidal Effect against. <i>Journal of Agricultural and Food Chemistry</i> , 2020 , 68, 7591-7600	5.7	6
49	High-throughput deep sequencing reveals the important role that microRNAs play in the salt response in sweet potato (Ipomoea batatas L.). <i>BMC Genomics</i> , 2020 , 21, 164	4.5	19
48	Overexpression of phosphatidylserine synthase affords cellular Na homeostasis and salt tolerance by activating plasma membrane Na/H antiport activity in sweet potato roots. <i>Horticulture Research</i> , 2020 , 7, 131	7.7	6
47	DORN1/P2K1 and purino-calcium signalling in plants: making waves with extracellular ATP. <i>Annals of Botany</i> , 2020 , 124, 1227-1242	4.1	18
46	Improving CRISPR-Cas-mediated RNA targeting and gene editing using SPLCV replicon-based expression vectors in Nicotiana benthamiana. <i>Plant Biotechnology Journal</i> , 2020 , 18, 1993	11.6	6
45	Genome-wide identification, structural and gene expression analysis of the bZIP transcription factor family in sweet potato wild relative Ipomoea trifida. <i>BMC Genetics</i> , 2019 , 20, 41	2.6	28
44	Root-zone-specific sensitivity of K+-and Ca2+-permeable channels to H2O2 determines ion homeostasis in salinized diploid and hexaploid Ipomoea trifida. <i>Journal of Experimental Botany</i> , 2019 , 70, 1389-1405	7	22
43	Involvement of Phosphatidylserine and Triacylglycerol in the Response of Sweet Potato Leaves to Salt Stress. <i>Frontiers in Plant Science</i> , 2019 , 10, 1086	6.2	14

42	A systematic comparison of eight new plastome sequences from L. <i>PeerJ</i> , 2019 , 7, e6563	3.1	6
41	Genome-wide identification, characterisation and functional evaluation of WRKY genes in the sweet potato wild ancestor Ipomoea trifida (H.B.K.) G. Don. under abiotic stresses. <i>BMC Genetics</i> , 2019 , 20, 90	2.6	11
40	Genome-wide identification and expression analysis of glycine-rich RNA-binding protein family in sweet potato wild relative Ipomoea trifida. <i>Gene</i> , 2019 , 686, 177-186	3.8	21
39	Identification, expression analysis, and function evaluation of 42 tomato DEAD-box RNA helicase genes in growth development and stress response. <i>Acta Physiologiae Plantarum</i> , 2018 , 40, 1	2.6	12
38	Melatonin-Stimulated Triacylglycerol Breakdown and Energy Turnover under Salinity Stress Contributes to the Maintenance of Plasma Membrane H-ATPase Activity and K/Na Homeostasis in Sweet Potato. <i>Frontiers in Plant Science</i> , 2018 , 9, 256	6.2	54
37	Hydrogen Sulfide Mediates K and Na Homeostasis in the Roots of Salt-Resistant and Salt-Sensitive Poplar Species Subjected to NaCl Stress. <i>Frontiers in Plant Science</i> , 2018 , 9, 1366	6.2	22
36	Isolation, Expression Analysis, and Function Evaluation of 12 Novel Stress-Responsive Genes of NAC Transcription Factors in Sweetpotato. <i>Crop Science</i> , 2018 , 58, 1328-1341	2.4	9
35	Extracellular ATP mediates cellular K+/Na+ homeostasis in two contrasting poplar species under NaCl stress. <i>Trees - Structure and Function</i> , 2016 , 30, 825-837	2.6	27
34	NaCl-induced changes of ion homeostasis and nitrogen metabolism in two sweet potato (Ipomoea batatas L.) cultivars exhibit different salt tolerance at adventitious root stage. <i>Environmental and Experimental Botany</i> , 2016 , 129, 23-36	5.9	27
33	Heat shock responses in Populus euphratica cell cultures: important role of crosstalk among hydrogen peroxide, calcium and potassium. <i>Plant Cell, Tissue and Organ Culture</i> , 2016 , 125, 215-230	2.7	6
32	-Facilitated Cd Influx through Plasma Membrane Ca-Permeable Channels Is Stimulated by HO and H-ATPase in Ectomycorrhizal Lunder Cadmium Stress. <i>Frontiers in Plant Science</i> , 2016 , 7, 1975	6.2	11
31	Exogenous Abscisic Acid Alleviates Cadmium Toxicity by Restricting Cd2+ Influx in Populus euphratica Cells. <i>Journal of Plant Growth Regulation</i> , 2016 , 35, 827-837	4.7	43
30	NaCl-elicited, vacuolar Ca(2+) release facilitates prolonged cytosolic Ca(2+) signaling in the salt response of Populus euphratica cells. <i>Cell Calcium</i> , 2015 , 57, 348-65	4	21
29	High rates of virus-induced gene silencing by tobacco rattle virus in Populus. <i>Tree Physiology</i> , 2015 , 35, 1016-29	4.2	22
28	Overexpression of copper/zinc superoxide dismutase from mangrove Kandelia candel in tobacco enhances salinity tolerance by the reduction of reactive oxygen species in chloroplast. <i>Frontiers in Plant Science</i> , 2015 , 6, 23	6.2	38
27	Populus euphratica APYRASE2 Enhances Cold Tolerance by Modulating Vesicular Trafficking and Extracellular ATP in Arabidopsis Plants. <i>Plant Physiology</i> , 2015 , 169, 530-48	6.6	48
26	Populus euphratica HSF binds the promoter of WRKY1 to enhance salt tolerance. <i>Plant Science</i> , 2015 , 235, 89-100	5.3	25
25	1-Butanol triggers programmed cell death in Populus euphratica cell cultures. <i>Plant Growth Regulation</i> , 2014 , 74, 33-45	3.2	O

24	Salt tolerance in Populus: Significance of stress signaling networks, mycorrhization, and soil amendments for cellular and whole-plant nutrition. <i>Environmental and Experimental Botany</i> , 2014 , 107, 113-124	5.9	52
23	Ion fluxes in Paxillus involutus-inoculated roots of PopulusBanescens under saline stress. <i>Environmental and Experimental Botany</i> , 2014 , 108, 99-108	5.9	15
22	Overexpression of Populus euphratica xyloglucan endotransglucosylase/hydrolase gene confers enhanced cadmium tolerance by the restriction of root cadmium uptake in transgenic tobacco. <i>Environmental and Experimental Botany</i> , 2014 , 100, 74-83	5.9	36
21	Overexpression of PeHSF mediates leaf ROS homeostasis in transgenic tobacco lines grown under salt stress conditions. <i>Plant Cell, Tissue and Organ Culture</i> , 2013 , 115, 299-308	2.7	21
20	Overexpression of a poplar two-pore K+ channel enhances salinity tolerance in tobacco cells. <i>Plant Cell, Tissue and Organ Culture</i> , 2013 , 112, 19-31	2.7	21
19	Hydrogen sulfide alleviates cadmium toxicity through regulations of cadmium transport across the plasma and vacuolar membranes in Populus euphratica cells. <i>Plant Physiology and Biochemistry</i> , 2013 , 65, 67-74	5.4	122
18	Non-invasive microelectrode cadmium flux measurements reveal the spatial characteristics and real-time kinetics of cadmium transport in hyperaccumulator and nonhyperaccumulator ecotypes of Sedum alfredii. <i>Journal of Plant Physiology</i> , 2013 , 170, 355-9	3.6	32
17	Overexpression of PeHA1 enhances hydrogen peroxide signaling in salt-stressed Arabidopsis. <i>Plant Physiology and Biochemistry</i> , 2013 , 71, 37-48	5.4	28
16	Populus euphratica XTH overexpression enhances salinity tolerance by the development of leaf succulence in transgenic tobacco plants. <i>Journal of Experimental Botany</i> , 2013 , 64, 4225-38	7	70
15	Exogenous hydrogen peroxide, nitric oxide and calcium mediate root ion fluxes in two non-secretor mangrove species subjected to NaCl stress. <i>Tree Physiology</i> , 2013 , 33, 81-95	4.2	47
14	Non-invasive flux measurements using microsensors: theory, limitations, and systems. <i>Methods in Molecular Biology</i> , 2012 , 913, 101-17	1.4	7
13	An ATP signalling pathway in plant cells: extracellular ATP triggers programmed cell death in Populus euphratica. <i>Plant, Cell and Environment</i> , 2012 , 35, 893-916	8.4	58
12	Extracellular ATP signaling and homeostasis in plant cells. <i>Plant Signaling and Behavior</i> , 2012 , 7, 566-9	2.5	14
11	Paxillus involutus strains MAJ and NAU mediate K(+)/Na(+) homeostasis in ectomycorrhizal Populus x canescens under sodium chloride stress. <i>Plant Physiology</i> , 2012 , 159, 1771-86	6.6	59
10	Arabidopsis fatty acid desaturase FAD2 is required for salt tolerance during seed germination and early seedling growth. <i>PLoS ONE</i> , 2012 , 7, e30355	3.7	110
9	Extracellular ATP signaling is mediated by HDDand cytosolic CaD+ in the salt response of Populus euphratica cells. <i>PLoS ONE</i> , 2012 , 7, e53136	3.7	61
8	H2O2 and cytosolic Ca2+ signals triggered by the PM H-coupled transport system mediate K+/Na+ homeostasis in NaCl-stressed Populus euphratica cells. <i>Plant, Cell and Environment</i> , 2010 , 33, 943-58	8.4	136
7	Effects of Stockosorb and Luquasorb polymers on salt and drought tolerance of Populus popularis. <i>Scientia Horticulturae</i> , 2010 , 124, 268-273	4.1	33

LIST OF PUBLICATIONS

6	Effect of NaCl on leaf H+-ATPase and the relevance to salt tolerance in two contrasting poplar species. <i>Trees - Structure and Function</i> , 2010 , 24, 597-607	2.6	14
5	Hydrogen peroxide and nitric oxide mediate K+/Na+ homeostasis and antioxidant defense in NaCl-stressed callus cells of two contrasting poplars. <i>Plant Cell, Tissue and Organ Culture</i> , 2010 , 103, 20	5 ² 2715	78
4	Salt-induced expression of genes related to Na(+)/K(+) and ROS homeostasis in leaves of salt-resistant and salt-sensitive poplar species. <i>Plant Molecular Biology</i> , 2010 , 73, 251-69	4.6	115
3	Ion flux profiles and plant ion homeostasis control under salt stress. <i>Plant Signaling and Behavior</i> , 2009 , 4, 261-4	2.5	29
2	Calcium mediates root K+/Na+ homeostasis in poplar species differing in salt tolerance. <i>Tree Physiology</i> , 2009 , 29, 1175-86	4.2	136
1	NaCl-induced alternations of cellular and tissue ion fluxes in roots of salt-resistant and salt-sensitive poplar species. <i>Plant Physiology</i> , 2009 , 149, 1141-53	6.6	253