Narikazu Boku

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3084082/publications.pdf

Version: 2024-02-01

76 papers 7,042 citations

172207 29 h-index 72 g-index

78 all docs

78 docs citations

78 times ranked 6907 citing authors

#	Article	IF	CITATIONS
1	Nivolumab in patients with advanced gastric or gastro-oesophageal junction cancer refractory to, or intolerant of, at least two previous chemotherapy regimens (ONO-4538-12, ATTRACTION-2): a randomised, double-blind, placebo-controlled, phase 3 trial. Lancet, The, 2017, 390, 2461-2471.	6.3	1,749
2	Adjuvant chemotherapy of S-1 versus gemcitabine for resected pancreatic cancer: a phase 3, open-label, randomised, non-inferiority trial (JASPAC 01). Lancet, The, 2016, 388, 248-257.	6.3	799
3	Randomized Phase III Study of Gemcitabine Plus S-1, S-1 Alone, or Gemcitabine Alone in Patients With Locally Advanced and Metastatic Pancreatic Cancer in Japan and Taiwan: GEST Study. Journal of Clinical Oncology, 2013, 31, 1640-1648.	0.8	548
4	Fluorouracil versus combination of irinotecan plus cisplatin versus S-1 in metastatic gastric cancer: a randomised phase 3 study. Lancet Oncology, The, 2009, 10, 1063-1069.	5.1	536
5	Randomized, Open-Label, Phase III Study Comparing Irinotecan With Paclitaxel in Patients With Advanced Gastric Cancer Without Severe Peritoneal Metastasis After Failure of Prior Combination Chemotherapy Using Fluoropyrimidine Plus Platinum: WJOG 4007 Trial. Journal of Clinical Oncology, 2013, 31, 4438-4444.	0.8	439
6	Definitive Chemoradiotherapy for T4 and/or M1 Lymph Node Squamous Cell Carcinoma of the Esophagus. Journal of Clinical Oncology, 1999, 17, 2915-2915.	0.8	394
7	HER2-positive gastric cancer. Gastric Cancer, 2014, 17, 1-12.	2.7	272
8	Nivolumab plus chemotherapy versus placebo plus chemotherapy in patients with HER2-negative, untreated, unresectable advanced or recurrent gastric or gastro-oesophageal junction cancer (ATTRACTION-4): a randomised, multicentre, double-blind, placebo-controlled, phase 3 trial. Lancet Oncology, The, 2022, 23, 234-247.	5.1	268
9	A late phase II study of S-1 for metastatic pancreatic cancer. Cancer Chemotherapy and Pharmacology, 2008, 61, 615-621.	1.1	156
10	A phase 3 study of nivolumab in previously treated advanced gastric or gastroesophageal junction cancer (ATTRACTION-2): 2-year update data. Gastric Cancer, 2020, 23, 510-519.	2.7	155
11	Phase III Trial of Avelumab Maintenance After First-Line Induction Chemotherapy Versus Continuation of Chemotherapy in Patients With Gastric Cancers: Results From JAVELIN Gastric 100. Journal of Clinical Oncology, 2021, 39, 966-977.	0.8	122
12	Randomized, Phase II Study of Trastuzumab Beyond Progression in Patients With HER2-Positive Advanced Gastric or Gastroesophageal Junction Cancer: WJOG7112G (T-ACT Study). Journal of Clinical Oncology, 2020, 38, 1919-1927.	0.8	107
13	Correlation between immune-related adverse events and prognosis in patients with gastric cancer treated with nivolumab. BMC Cancer, 2019, 19, 974.	1.1	104
14	Randomized Phase III Trial of Adjuvant Chemotherapy with Gemcitabine versus S-1 in Patients with Resected Pancreatic Cancer: Japan Adjuvant Study Group of Pancreatic Cancer (JASPAC-01). Japanese Journal of Clinical Oncology, 2008, 38, 227-229.	0.6	95
15	Docetaxel plus cisplatin and S-1 versus cisplatin and S-1 in patients with advanced gastric cancer (JCOG1013): an open-label, phase 3, randomised controlled trial. The Lancet Gastroenterology and Hepatology, 2019, 4, 501-510.	3.7	88
16	Evolution of checkpoint inhibitors for the treatment of metastatic gastric cancers: Current status and future perspectives. Cancer Treatment Reviews, 2018, 66, 104-113.	3.4	78
17	Four courses versus eight courses of adjuvant S-1 for patients with stage II gastric cancer (JCOG1104) Tj ETQq1 1 Hepatology, 2019, 4, 208-216.	0.784314 3.7	rgBT /Ov <mark>er</mark> 73
18	S-1 plus leucovorin versus S-1 plus leucovorin and oxaliplatin versus S-1 plus cisplatin in patients with advanced gastric cancer: a randomised, multicentre, open-label, phase 2 trial. Lancet Oncology, The, 2016, 17, 99-108.	5.1	63

#	Article	IF	CITATIONS
19	Nivolumab in previously treated advanced gastric cancer (ATTRACTION-2): 3-year update and outcome of treatment beyond progression with nivolumab. Gastric Cancer, 2021, 24, 946-958.	2.7	61
20	Randomized study of <scp>FOLFIRI</scp> plus either panitumumab or bevacizumab for wildâ€type <scp>KRAS</scp> colorectal cancerâ€ <scp>WJOG</scp> 6210G. Cancer Science, 2016, 107, 1843-1850.	1.7	60
21	A subanalysis of Japanese patients in a randomized, double-blind, placebo-controlled, phase 3 trial of nivolumab for patients with advanced gastric or gastro-esophageal junction cancer refractory to, or intolerant of, at least two previous chemotherapy regimens (ONO-4538-12, ATTRACTION-2). Gastric Cancer. 2019. 22. 344-354.	2.7	60
22	Determination of Prognostic Factors in Japanese Patients With Advanced Gastric Cancer Using the Data From a Randomized Controlled Trial, Japan Clinical Oncology Group 9912. Oncologist, 2014, 19, 358-366.	1.9	54
23	Exploratory subgroup analysis of patients with prior trastuzumab use in the ATTRACTION-2 trial: a randomized phase III clinical trial investigating the efficacy and safety of nivolumab in patients with advanced gastric/gastroesophageal junction cancer. Gastric Cancer, 2020, 23, 143-153.	2.7	45
24	Current status of immunotherapy for advanced gastric cancer. Japanese Journal of Clinical Oncology, 2021, 51, 20-27.	0.6	43
25	Multi-omic profiling of peritoneal metastases in gastric cancer identifies molecular subtypes and therapeutic vulnerabilities. Nature Cancer, 2021, 2, 962-977.	5.7	41
26	Weekly paclitaxel for heavily treated advanced or recurrent gastric cancer refractory to fluorouracil, irinotecan, and cisplatin. Gastric Cancer, 2009, 12, 206-211.	2.7	39
27	Hyperprogressive disease during nivolumab or irinotecan treatment in patients with advanced gastric cancer. ESMO Open, 2019, 4, e000488.	2.0	39
28	S-1 plus leucovorin and oxaliplatin versus S-1 plus cisplatin as first-line therapy in patients with advanced gastric cancer (SOLAR): a randomised, open-label, phase 3 trial. Lancet Oncology, The, 2020, 21, 1045-1056.	5.1	39
29	Phase I/II study of ramucirumab plus nivolumab in patients in second-line treatment for advanced gastric adenocarcinoma (NivoRam study) Journal of Clinical Oncology, 2019, 37, 129-129.	0.8	32
30	A randomized phase II study of weekly paclitaxel $\hat{A}\pm$ trastuzumab in patients with HER2-positive advanced gastric or gastro-esophageal junction cancer refractory to trastuzumab combined with fluoropyrimidine and platinum: WJOG7112G (T-ACT) Journal of Clinical Oncology, 2018, 36, 4011-4011.	0.8	30
31	Survival impact of post-progression chemotherapy in advanced gastric cancer: systematic review and meta-analysis. Cancer Chemotherapy and Pharmacology, 2018, 81, 981-989.	1.1	29
32	Efficacy of Postoperative Chemotherapy After Resection that Leaves No Macroscopically Visible Disease of Gastric Cancer with Positive Peritoneal Lavage Cytology (CY1) or Localized Peritoneum Metastasis (P1a): A Multicenter Retrospective Study. Annals of Surgical Oncology, 2020, 27, 284-292.	0.7	28
33	Regional differences in advanced gastric cancer: exploratory analyses of the AVAGAST placebo arm. Gastric Cancer, 2018, 21, 429-438.	2.7	26
34	IL33 Is a Key Driver of Treatment Resistance of Cancer. Cancer Research, 2020, 80, 1981-1990.	0.4	24
35	Validation of the JCOG prognostic index in advanced gastric cancer using individual patient data from the SPIRITS and G-SOX trials. Gastric Cancer, 2017, 20, 757-763.	2.7	21
36	SOURCE: A Registry-Based Prediction Model for Overall Survival in Patients with Metastatic Oesophageal or Gastric Cancer. Cancers, 2019, 11, 187.	1.7	20

#	Article	IF	Citations
37	Maintenance avelumab versus continuation of first-line chemotherapy in gastric cancer: JAVELIN Gastric 100 study design. Future Oncology, 2019, 15, 567-577.	1.1	20
38	Significance of FGF9 gene in resistance to anti-EGFR therapies targeting colorectal cancer: A subset of colorectal cancer patients with FGF9 upregulation may be resistant to anti-EGFR therapies. Molecular Carcinogenesis, 2017, 56, 106-117.	1.3	19
39	Trastuzumab deruxtecan for the treatment of HER2-positive advanced gastric cancer: a clinical perspective. Gastric Cancer, 2021, 24, 567-576.	2.7	19
40	A randomized phase II study of combination therapy with S-1, oral leucovorin, and oxaliplatin (SOL) and mFOLFOX6 in patients with previously untreated metastatic colorectal cancer. Cancer Chemotherapy and Pharmacology, 2015, 75, 569-577.	1.1	16
41	The current status of multimodality treatment for unresectable locally advanced esophageal squamous cell carcinoma. Asia-Pacific Journal of Clinical Oncology, 2018, 14, 291-299.	0.7	16
42	Multidisciplinary management of stage II-III gastric and gastro-oesophageal junction cancer. European Journal of Cancer, 2020, 124, 67-76.	1.3	16
43	Health-related quality of life of adjuvant chemotherapy with S-1 versus gemcitabine for resected pancreatic cancer: Results from a randomised phase III trial (JASPAC 01). European Journal of Cancer, 2018, 93, 79-88.	1.3	14
44	Association between UGT1A1 gene polymorphism and safety and efficacy of irinotecan monotherapy as the third-line treatment for advanced gastric cancer. Gastric Cancer, 2019, 22, 778-784.	2.7	13
45	Treatment Pattern for Advanced Gastric Cancer in Japan and Factors Associated with Sequential Treatment: A Retrospective Administrative Claims Database Study. Advances in Therapy, 2022, 39, 296-313.	1.3	13
46	Human equilibrative nucleoside transporterâ€1 expression is a predictor in patients with resected pancreatic cancer treated with adjuvant Sâ€1 chemotherapy. Cancer Science, 2020, 111, 548-560.	1.7	12
47	Japan Society of Clinical Oncology Clinical Practice Guidelines 2017 for fertility preservation in childhood, adolescent, and young adult cancer patients: part 2. International Journal of Clinical Oncology, 2022, 27, 281-300.	1.0	11
48	Japan Society of Clinical Oncology Clinical Practice Guidelines 2017 for fertility preservation in childhood, adolescent, and young adult cancer patients: part 1. International Journal of Clinical Oncology, 2022, 27, 265-280.	1.0	11
49	Development of an S-1 dosage formula based on renal function by a prospective pharmacokinetic study. Gastric Cancer, 2016, 19, 876-886.	2.7	10
50	Assessment of hyperprogression versus the natural course of disease development with nivolumab with or without ipilimumab versus placebo in phase III, randomized, controlled trials., 2022, 10, e004273.		10
51	Exploration of predictors of benefit from nivolumab monotherapy for patients with pretreated advanced gastric and gastroesophageal junction cancer: post hoc subanalysis from the ATTRACTION-2 study. Gastric Cancer, 2022, 25, 207-217.	2.7	9
52	Efficacy and safety of taxane monotherapy in advanced gastric cancer refractory to triplet chemotherapy with docetaxel, cisplatin, and S-1: a multicenter retrospective study. Cancer Chemotherapy and Pharmacology, 2017, 80, 575-582.	1.1	6
53	Study protocol for a multi-institutional randomized phase III study comparing combined everolimus plus lanreotide therapy and everolimus monotherapy in patients with unresectable or recurrent gastroenteropancreatic neuroendocrine tumors; Japan Clinical Oncology Group Study JCOG1901 (STARTER-NET study), Pancreatology, 2020, 20, 1183-1188.	0.5	6
54	Diagnosis of invasion depth in resectable advanced gastric cancer for neoadjuvant chemotherapy: An exploratory analysis of Japan clinical oncology group study: JCOG1302A. European Journal of Surgical Oncology, 2020, 46, 1074-1079.	0.5	6

#	Article	IF	CITATIONS
55	Retrospective observational study of salvage line ramucirumab monotherapy for patients with advanced gastric cancer. BMC Cancer, 2020, 20, 338.	1.1	6
56	Surgical and perioperative treatment strategy for resectable esophagogastric junction cancer. Japanese Journal of Clinical Oncology, 2022, 52, 417-424.	0.6	6
57	S-1 Monotherapy After Failure of Platinum Plus 5-Fluorouracil Chemotherapy in Recurrent or Metastatic Esophageal Carcinoma. Anticancer Research, 2019, 39, 3931-3936.	0.5	5
58	Impact of peripheral neuropathy induced by platinum in first-line chemotherapy on second-line chemotherapy with paclitaxel for advanced gastric cancer. International Journal of Clinical Oncology, 2020, 25, 595-601.	1.0	5
59	Prospective evaluation and refinement of an S†dosage formula based on renal function for clinical application. Cancer Science, 2021, 112, 751-759.	1.7	5
60	Second-line chemotherapy after early disease progression during first-line chemotherapy containing bevacizumab for patients with metastatic colorectal cancer. BMC Cancer, 2021, 21, 1159.	1.1	5
61	Safety and tolerability of andecaliximab as monotherapy and in combination with an anti-PD-1 antibody in Japanese patients with gastric or gastroesophageal junction adenocarcinoma: a phase $1b$ study., 2022, 10 , e003518.		5
62	Second gastric cancer after curative endoscopic resection of differentiated-type early gastric cancer: post-hoc analysis of a single-arm confirmatory trial. Gastrointestinal Endoscopy, 2022, 95, 650-659.	0.5	5
63	Phase I clinical and pharmacokinetic study of S-1 plus oral leucovorin in patients with metastatic colorectal cancer. Cancer Chemotherapy and Pharmacology, 2017, 79, 107-116.	1.1	4
64	A Phase I study of pevonedistat plus capecitabine plus oxaliplatin in patients with advanced gastric cancer refractory to platinum (NCCH-1811). Future Science OA, 2021, 7, FSO721.	0.9	4
65	Real-world safety and effectiveness of nivolumab in Japanese patients with unresectable advanced or recurrent gastric/gastroesophageal junction cancer that has progressed after chemotherapy: a postmarketing surveillance study. Gastric Cancer, 2021, , 1.	2.7	4
66	Temporal dynamics from phosphoproteomics using endoscopic biopsy specimens provides new therapeutic targets in stage IV gastric cancer. Scientific Reports, 2022, 12, 4419.	1.6	4
67	Consensus Statement on Mandatory Measurements for Pancreatic Cancer Trials for Patients With Resectable or Borderline Resectable Disease (COMM-PACT-RB). JAMA Oncology, 2022, 8, 929.	3.4	4
68	Selection of Second-line Anti-angiogenic Agents After Failure of Bevacizumab-containing First-line Chemotherapy in Metastatic Colorectal Cancer. Clinical Colorectal Cancer, 2018, 17, 251-254.	1.0	3
69	Concordance of human equilibrative nucleoside transporter†expressions between murine (10D7G2) and rabbit (SP120) antibodies and association with clinical outcomes of adjuvant chemotherapy for pancreatic cancer: A collaborative study from the JASPAC 01 trial. Cancer Reports, 2021, , e1507.	0.6	3
70	Trastuzumab Emtansine (T-DM1) Plus S-1 in Patients with Trastuzumab-Pretreated HER2-Positive Advanced or Metastatic Breast Cancer: A Phase Ib Study. Oncology, 2019, 96, 309-317.	0.9	2
71	Primary Tumor-Related Complications Among Patients With Unresectable Stage IV Colorectal Cancer in the Era of Targeted Therapy: A Competing Risk Regression Analysis. Diseases of the Colon and Rectum, 2021, 64, 1074-1082.	0.7	2
72	Influence of precedent drug on the subsequent therapy in the sequence of trifluridine/tipiracil with/out bevacizumab and regorafenib for unresectable or recurrent colorectal cancer. PLoS ONE, 2022, 17, e0269115.	1.1	2

#	Article	IF	CITATIONS
73	A phase $1b$ study of andecaliximab in combination with S-1 plus platinum in Japanese patients with gastric adenocarcinoma. Scientific Reports, 2022, 12, .	1.6	2
74	Prediction of the peritoneal recurrence via the macroscopic diagnosis of the serosal invasion in patients with gastric cancer: Supplementary analysis of JCOG0110. European Journal of Surgical Oncology, 2022, , .	0.5	1
75	Evaluation of clinical validity of an S-1 dosage formula based on renal function using data of the SPIRITS and the G-SOX trials. Gastric Cancer, 2022, , 1.	2.7	1
76	Usefulness of an S-1 dosage formula: an exploratory analysis of randomized clinical trial (JCOG1001). Gastric Cancer, 0, , .	2.7	O