Peter H Gæde

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3093052/publications.pdf

Version: 2024-02-01

50 9,787 20 48
papers citations h-index g-index

52 52 52 7641 all docs docs citations times ranked citing authors

#	Article	IF	Citations
1	Multifactorial Intervention and Cardiovascular Disease in Patients with Type 2 Diabetes. New England Journal of Medicine, 2003, 348, 383-393.	27.0	3,894
2	Effect of a Multifactorial Intervention on Mortality in Type 2 Diabetes. New England Journal of Medicine, 2008, 358, 580-591.	27.0	3,037
3	Intensified multifactorial intervention in patients with type 2 diabetes mellitus and microalbuminuria: the Steno type 2 randomised study. Lancet, The, 1999, 353, 617-622.	13.7	911
4	Years of life gained by multifactorial intervention in patients with type 2 diabetes mellitus and microalbuminuria: 21Âyears follow-up on the Steno-2 randomised trial. Diabetologia, 2016, 59, 2298-2307.	6.3	378
5	Mutation analysis of peroxisome proliferator-activated receptor-Î ³ coactivator-1 (PGC-1) and relationships of identified amino acid polymorphisms to Type II diabetes mellitus. Diabetologia, 2001, 44, 2220-2226.	6.3	288
6	Remission to normoalbuminuria during multifactorial treatment preserves kidney function in patients with type 2 diabetes and microalbuminuria. Nephrology Dialysis Transplantation, 2004, 19, 2784-2788.	0.7	188
7	Cost-Effectiveness of Intensified Versus Conventional Multifactorial Intervention in Type 2 Diabetes. Diabetes Care, 2008, 31, 1510-1515.	8.6	130
8	Double-blind, randomised study of the effect of combined treatment with vitamin C and E on albuminuria in Type 2 diabetic patients. Diabetic Medicine, 2001, 18, 756-760.	2.3	107
9	Plasma N-terminal pro-brain natriuretic peptide as a major risk marker for cardiovascular disease in patients with type 2 diabetes and microalbuminuria. Diabetologia, 2005, 48, 156-163.	6.3	89
10	A Single Nucleotide Polymorphism within the Acetyl-Coenzyme A Carboxylase Beta Gene Is Associated with Proteinuria in Patients with Type 2 Diabetes. PLoS Genetics, 2010, 6, e1000842.	3.5	81
11	Intensified multifactorial intervention in type 2 diabetics with microalbuminuria leads to long-term renal benefits. Kidney International, 2017, 91, 982-988.	5.2	80
12	Reduced risk of heart failure with intensified multifactorial intervention in individuals with type 2 diabetes and microalbuminuria: 21Âyears of follow-up in the randomised Steno-2 study. Diabetologia, 2018, 61, 1724-1733.	6.3	66
13	Intensified multifactorial intervention and cardiovascular outcome in type 2 diabetes: the Steno-2 study. Metabolism: Clinical and Experimental, 2003, 52, 19-23.	3.4	65
14	Polymorphisms in the 3′ UTR in the neurocalcin δgene affect mRNA stability, and confer susceptibility to diabetic nephropathy. Human Genetics, 2007, 122, 397-407.	3.8	57
15	Limited impact of lifestyle education in patients with Type 2 diabetes mellitus and microalbuminuria: results from a randomized intervention study. Diabetic Medicine, 2001, 18, 104-108.	2.3	36
16	Cardiac perfusion, structure, and function in type 2 diabetes mellitus with and without diabetic complications. European Heart Journal Cardiovascular Imaging, 2020, 21, 887-895.	1.2	28
17	Reduced Myocardial Perfusion Reserve in Type 2 Diabetes Is Caused by Increased Perfusion at Rest and Decreased Maximal Perfusion During Stress. Diabetes Care, 2020, 43, 1285-1292.	8.6	25
18	Management of Patients with Type 2 Diabetes with Once-Weekly Semaglutide Versus Dulaglutide, Exenatide ER, Liraglutide and Lixisenatide: A Cost-Effectiveness Analysis in the Danish Setting. Diabetes Therapy, 2019, 10, 1297-1317.	2.5	23

#	Article	IF	CITATIONS
19	Lack of impact of low-dose acetylsalicylic acid on kidney function in type 1 diabetic patients with microalbuminuria. Diabetes Care, 2000, 23, 1742-1745.	8.6	21
20	Glycaemic variability and hypoglycaemia are associated with C-peptide levels in insulin-treated type 2 diabetes. Diabetes and Metabolism, 2020, 46, 61-65.	2.9	21
21	Distinct non-ischemic myocardial late gadolinium enhancement lesions in patients with type 2 diabetes. Cardiovascular Diabetology, 2020, 19, 184.	6.8	21
22	Plasma trimethylamine N-oxide and its metabolic precursors and risk of mortality, cardiovascular and renal disease in individuals with type 2-diabetes and albuminuria. PLoS ONE, 2021, 16, e0244402.	2.5	20
23	Beneficial impact of intensified multifactorial intervention on risk of stroke: outcome of 21Âyears of follow-up in the randomised Steno-2 Study. Diabetologia, 2019, 62, 1575-1580.	6.3	19
24	Impact of low-dose acetylsalicylic acid on kidney function in type 2 diabetic patients with elevated urinary albumin excretion rate. Nephrology Dialysis Transplantation, 2003, 18, 539-542.	0.7	16
25	A cost analysis of intensified vs conventional multifactorial therapy in individuals with type 2 diabetes: a post hoc analysis of the Steno-2 study. Diabetologia, 2019, 62, 147-155.	6.3	16
26	Target intervention against multipleâ€risk markers to reduce cardiovascular disease in patients with type 2 diabetes. Annals of Medicine, 2004, 36, 355-366.	3.8	15
27	Elevated levels of plasma von Willebrand factor and the risk of macro―and microvascular disease in type 2 diabetic patients with microalbuminuria. Nephrology Dialysis Transplantation, 2001, 16, 2028-2033.	0.7	14
28	Fibroblast growth factor-23 is associated with imaging markers of diabetic cardiomyopathy and anti-diabetic therapeutics. Cardiovascular Diabetology, 2020, 19, 158.	6.8	14
29	PROactive study. Lancet, The, 2006, 367, 23-24.	13.7	13
30	Effects of Empagliflozin on Myocardial Flow Reserve in Patients With Type 2 Diabetes Mellitus: The SIMPLE Trial. Journal of the American Heart Association, 2021, 10, e020418.	3.7	12
31	Efficacy of Bolus Calculation and Advanced Carbohydrate Counting in Type 2 Diabetes: A Randomized Clinical Trial. Diabetes Technology and Therapeutics, 2021, 23, 95-103.	4.4	10
32	Prevalence of urinary incontinence among women with diabetes in the Lollandâ€Falster Health Study, Denmark. Neurourology and Urodynamics, 2021, 40, 855-867.	1.5	10
33	Indicator of RNA oxidation in urine for the prediction of mortality in patients with type 2 diabetes and microalbuminuria: A post-hoc analysis of the Steno-2 trial. Free Radical Biology and Medicine, 2018, 129, 247-255.	2.9	9
34	Application of urinary proteomics as possible risk predictor of renal and cardiovascular complications in patients with type 2-diabetes and microalbuminuria. Journal of Diabetes and Its Complications, 2018, 32, 1133-1140.	2.3	9
35	Metabolic improvement with shortâ€term, glucagonâ€like peptideâ€1 receptor agonist treatment does not improve cardiac diastolic dysfunction in patients with type 2 diabetes: A randomized, doubleâ€blind, placeboâ€controlled trial. Diabetes, Obesity and Metabolism, 2021, 23, 2374-2384.	4.4	9
36	Studies of the variability of the hepatocyte nuclear factor- $1\hat{l}^2$ (HNF- $1\hat{l}^2$ / TCF2) and the dimerization cofactor of HNF-1 (DcoH / PCBD) genes in relation to type 2 diabetes mellitus and \hat{l}^2 -cell function. Human Mutation, 2001, 18, 356-357.	2.5	8

#	Article	IF	CITATIONS
37	Hypoglycaemia and its management in primary care setting. Diabetes/Metabolism Research and Reviews, 2020, 36, e3332.	4.0	8
38	The economic burden of poor glycemic control associated with therapeutic inertia in patients with type 2 diabetes in Denmark. Current Medical Research and Opinion, 2021, 37, 949-956.	1.9	8
39	Effect of empagliflozin on myocardial structure and function in patients with type 2 diabetes at high cardiovascular risk: the SIMPLE randomized clinical trial. International Journal of Cardiovascular Imaging, 2021, , 1.	1.5	6
40	Short-term societal economic burden of first-incident type 2 diabetes-related complications – a nationwide cohort study. Expert Review of Pharmacoeconomics and Outcomes Research, 2020, 20, 577-586.	1.4	5
41	Subclinical hypothyroidism: A common finding in adult patients with cyanotic congenital heart disease. Congenital Heart Disease, 2018, 13, 263-270.	0.2	4
42	Interventions involving own treatment choice for people living with coexisting severe mental illness and type 1 or 2 diabetes: A scoping review. Diabetic Medicine, 2021, 38, e14626.	2.3	4
43	Intervención sobre múltiples factores de riesgo para prevenir la enfermedad cardiovascular. Un enfoque basado en la evidencia. Revista Espanola De Cardiologia, 2011, 64, 173-174.	1.2	3
44	Mitochondrial energetics and contents evaluated by flow cytometry in human maternal and umbilical cord blood. Scandinavian Journal of Clinical and Laboratory Investigation, 2020, 80, 351-359.	1.2	3
45	Monitoring patients with acute dyspnoea with a serial focused ultrasound of the heart and the lungs (MODUS): a protocol for a multicentre, randomised, open-label, pragmatic and controlled trial. BMJ Open, 2020, 10, e034373.	1.9	2
46	Intensive glucose control and cardiovascular disease in type 2 diabetes – should we change the recommended target for glycated hemoglobin? Commentary to ACCORD and ADVANCE trials. Polish Archives of Internal Medicine, 2008, 118, 619-621.	0.4	2
47	Multiple Risk Factor Intervention to Prevent Cardiovascular Disease. A High Powered and Evidence Based Approach. Revista Espanola De Cardiologia (English Ed), 2011, 64, 173-174.	0.6	1
48	A Cost Analysis of Intensified vs. Conventional Multifactorial Therapy of Patients with Type 2 Diabetes—The Steno 2 Study. Diabetes, 2018, 67, 162-OR.	0.6	1
49	Rosiglitazone monotherapy for type 2 diabetes mellitusâ€"too soon to ADOPT?. Nature Clinical Practice Endocrinology and Metabolism, 2007, 3, 456-457.	2.8	0
50	Hypothyroidism and urinary incontinence: Prevalence and association in a Danish, female sample from the Lolland-Falster Health study. European Journal of Obstetrics, Gynecology and Reproductive Biology, 2021, 264, 232-240.	1.1	0