

Leonardo Fainboim

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3101494/publications.pdf>

Version: 2024-02-01

44
papers

1,486
citations

279798

23
h-index

315739

38
g-index

45
all docs

45
docs citations

45
times ranked

1865
citing authors

#	ARTICLE	IF	CITATIONS
1	Intrahepatic and peripheral blood phenotypes of natural killer and T cells: differential surface expression of killer cell immunoglobulin-like receptors. <i>Immunology</i> , 2018, 154, 261-273.	4.4	6
2	Historical records under the genetic evidence: "Chiriguano" tribe genesis as a test case. <i>Molecular Biology Reports</i> , 2018, 45, 987-1000.	2.3	7
3	IL28B SNPs rs12979860 and rs8099917 Are Associated with Inflammatory Response in Argentine Chronic HCV Patients. <i>International Journal of Clinical Medicine</i> , 2018, 09, 79-91.	0.2	4
4	The Clinical Features of Patients with Chronic Hepatitis C Virus Infections Are Associated with Killer Cell Immunoglobulin-Like Receptor Genes and Their Expression on the Surface of Natural Killer Cells. <i>Frontiers in Immunology</i> , 2017, 8, 1912.	4.8	19
5	ATP-Induced Inflammation Drives Tissue-Resident Th17 Cells in Metabolically Unhealthy Obesity. <i>Journal of Immunology</i> , 2016, 196, 3287-3296.	0.8	88
6	The tumor antigen N-glycolyl-GM3 is a human CD1d ligand capable of mediating B cell and natural killer T cell interaction. <i>Cancer Immunology, Immunotherapy</i> , 2016, 65, 551-562.	4.2	17
7	Immune dysfunction in cirrhosis: Distinct cytokines phenotypes according to cirrhosis severity. <i>Cytokine</i> , 2016, 77, 14-25.	3.2	84
8	Overexpression of CD85j in TNBC patients inhibits Cetuximab-mediated NK cell ADCC but can be restored with CD85j functional blockade. <i>European Journal of Immunology</i> , 2015, 45, 1560-1569.	2.9	39
9	Identification and Clinical Relevance of Naturally Occurring Human CD8+HLA-DR+ Regulatory T Cells. <i>Journal of Immunology</i> , 2014, 193, 4469-4476.	0.8	65
10	Human memory B cells isolated from blood and tonsils are functionally distinctive. <i>Immunology and Cell Biology</i> , 2014, 92, 882-887.	2.3	24
11	Regulatory and effector T-cells are differentially modulated by Dexamethasone. <i>Clinical Immunology</i> , 2013, 149, 400-410.	3.2	29
12	Analysis of Suppressor and Non-Suppressor FOXP3+ T Cells in HIV-1-Infected Patients. <i>PLoS ONE</i> , 2012, 7, e52580.	2.5	8
13	Anti-ganglioside antibodies induced in chickens by an alum-adsorbed anti-idiotypic antibody targeting NeuGcGM3. <i>Frontiers in Immunology</i> , 2012, 3, 422.	4.8	3
14	The most severe forms of type I autoimmune hepatitis are associated with genetically determined levels of TGF- β 1. <i>Clinical Immunology</i> , 2010, 134, 305-312.	3.2	25
15	A physiological role for inducible FOXP3+ TREG cells. <i>Clinical Immunology</i> , 2010, 136, 432-441.	3.2	38
16	Cytokines and chronic liver disease. <i>Cytokine and Growth Factor Reviews</i> , 2007, 18, 143-157.	7.2	28
17	Gender Susceptibility to Chronic Hepatitis C Virus Infection Associated with Interleukin 10 Promoter Polymorphism. <i>Journal of Virology</i> , 2006, 80, 9144-9150.	3.4	82
18	Cellular and Humoral Immune Response to N-Glycolyl-GM3 Elicited by Prolonged Immunotherapy With an Anti-Idiotypic Vaccine in High-Risk and Metastatic Breast Cancer Patients. <i>Journal of Immunotherapy</i> , 2006, 29, 215-223.	2.4	58

#	ARTICLE	IF	CITATIONS
19	Induction of Maternal Tolerance to Fetal Alloantigens by RANTES Production. American Journal of Reproductive Immunology, 2006, 56, 302-311.	1.2	24
20	New Actors for the Immunological Mechanisms Involved in the Materno- Fetal Tolerance. Current Women's Health Reviews, 2005, 1, 15-20.	0.2	2
21	Identification of RANTES as a novel immunomodulator of the maternal allogeneic response. Clinical Immunology, 2004, 110, 71-80.	3.2	26
22	Simultaneous expression of th1 cytokines and IL-4 confers severe characteristics to type I autoimmune hepatitis in children. Human Immunology, 2004, 65, 683-691.	2.4	42
23	Evidences of the Involvement of Bak, a member of the Bcl-2 Family of Proteins, in Active Coeliac Disease. Autoimmunity, 2002, 35, 29-37.	2.6	11
24	HLA Haplotypes and Class II Molecular Alleles in Argentinian Patients with Pemphigus Vulgaris. Journal of Cutaneous Medicine and Surgery, 2002, 6, 422-426.	1.2	9
25	HLA Haplotypes and Class II Molecular Alleles in Argentinian Patients with Pemphigus Vulgaris. Journal of Cutaneous Medicine and Surgery, 2002, 6, 422-426.	1.2	13
26	Protracted, but not acute, hepatitis A virus infection is strongly associated with HLA-DRB1*1301, a marker for pediatric autoimmune hepatitis. Hepatology, 2001, 33, 1512-1517.	7.3	127
27	Modulation of CD44 in acute lymphoblastic leukemia identifies functional and phenotypic differences of human B cell precursors. European Journal of Haematology, 2001, 66, 377-382.	2.2	5
28	Single-shot plasmid DNA intrasplenic immunization for the production of monoclonal antibodies. Journal of Immunological Methods, 2000, 244, 1-7.	1.4	30
29	Pediatric and adult forms of type I autoimmune hepatitis in argentina: Evidence for differential genetic predisposition. Hepatology, 1999, 30, 1374-1380.	7.3	192
30	Two-locus involvement in the association of human leukocyte antigen with the extrahepatic manifestations of autoimmune chronic active hepatitis. Hepatology, 1994, 19, 1371-1374.	7.3	32
31	Molecular characterization of HLA class II genes in celiac disease patients of Latin American caucasian origin. Tissue Antigens, 1994, 43, 83-87.	1.0	29
32	Chronic active autoimmune hepatitis in children. Human Immunology, 1994, 41, 146-150.	2.4	119
33	Hepatitis B Virus Antigens in Peripheral Blood Mononuclear Cells during the Course of Viral Infection. Clinical Immunology and Immunopathology, 1994, 70, 99-103.	2.0	11
34	Assessment by Planimetry of Langerhans's Cell Density in Penile Epithelium with Human Papillomavirus Infection: Changes Observed After Topical Treatment. Journal of Urology, 1992, 147, 1268-1273.	0.4	14
35	Intracellular expression of CD1 molecules on PHA-activated normal T lymphocytes. Immunology Letters, 1992, 34, 91.	2.5	0
36	Cytoplasmic expression of a CD24-related epitope in human PHA activated normal T lymphocytes. Immunology Letters, 1992, 34, 109-113.	2.5	4

#	ARTICLE	IF	CITATIONS
37	Intracellular expression of CD1 molecules on PHA-activated normal T lymphocytes. Immunology Letters, 1992, 33, 61-65.	2.5	5
38	Complete nucleotide sequence of a genomic clone encoding HLA-B35 isolated from a Caucasian individual of hispanic origin identification of a new variant of HLA-B35. Human Immunology, 1991, 31, 153-158.	2.4	25
39	Expression of CD1 antigens by peripheral blood mononuclear cells from hepatitis B patients. Immunology Letters, 1989, 21, 139-143.	2.5	2
40	Restriction fragment length polymorphism in HLA class II genes of Latin-American caucasian celiac disease patients. Human Immunology, 1989, 26, 272-280.	2.4	27
41	Differential expression of CD25 and HC2 antigens on subtypes of acute myeloid leukemias. European Journal of Haematology, 1989, 42, 436-440.	2.2	11
42	DNA polymorphism of the HLA-B35 gene associated to different HLA-C locus alleles. Human Immunology, 1988, 23, 241-253.	2.4	11
43	Genetic and functional relationship of the HLA-DR and HLA-DQ antigens. Immunogenetics, 1985, 21, 97-101.	2.4	26
44	Differential expression of HLA-DR and DR-linked determinants on human leukemias and lymphoid cells. European Journal of Immunology, 1983, 13, 172-176.	2.9	64