

John D Kennedy

List of Publications by Citations

Source: <https://exaly.com/author-pdf/3104086/john-d-kennedy-publications-by-citations.pdf>

Version: 2024-04-29

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

385
papers

6,939
citations

35
h-index

47
g-index

413
ext. papers

7,298
ext. citations

3.7
avg, IF

5.17
L-index

| # | Paper | IF | Citations |
|-----|---|-----|-----------|
| 385 | The Polyhedral Metallaboranes Part II. Metallaborane Clusters with Eight Vertices and More. <i>Progress in Inorganic Chemistry</i> , 211-434 | | 120 |
| 384 | Ten-vertex metallaborane chemistry. Aspects of the iridadecaborane closo-isonido-isocloso structural continuum. <i>Journal of the Chemical Society Dalton Transactions</i> , 1992 , 563-576 | | 85 |
| 383 | The Polyhedral Metallaboranes Part I. Metallaborane Clusters with Seven Vertices and Fewer. <i>Progress in Inorganic Chemistry</i> , 519-679 | | 84 |
| 382 | Polyhedral Monocarbaborane Chemistry. A Review of Recent Developments Among C-Aryl Monocarbaborane Systems. <i>Collection of Czechoslovak Chemical Communications</i> , 2002 , 67, 869-912 | | 81 |
| 381 | Identification of the endo,exo isomer of 6,9-(PMe2Ph)2-arachno-B10H12 by nuclear magnetic resonance spectroscopy. <i>Journal of the Chemical Society Dalton Transactions</i> , 1987 , 1573 | | 79 |
| 380 | Indirect nuclear spin-spin coupling of lead-207 to other magnetic nuclei. <i>Inorganic Chemistry</i> , 1976 , 15, 1299-1302 | 5.1 | 77 |
| 379 | Structure and bonding in recently isolated metallaboranes. <i>Inorganic Chemistry</i> , 1986 , 25, 111-112 | 5.1 | 73 |
| 378 | Nine-vertex polyhedral iridamonocarbaborane chemistry. Products of thermolysis of [(CO)(PPh3)2IrCB7H8] and emerging alternative cluster-geometry patterns. <i>Journal of the Chemical Society Dalton Transactions</i> , 1994 , 229-236 | | 70 |
| 377 | One-Dimensional Coordination Polymers with Phenyl-carbaborane Anions: Ag(I)/4,4'-Bipyridine and 2,3-Bis-(2-pyridyl)pyrazine Complexes. <i>Crystal Growth and Design</i> , 2007 , 7, 658-667 | 3.5 | 61 |
| 376 | Polyhedral metallacarborane chemistry: preparation, molecular structure, and nuclear magnetic resonance investigation of [3-(η -C5Me5)-closo-3,1,2-MC2B9H11] (M = Rh or Ir). <i>Journal of the Chemical Society Dalton Transactions</i> , 1990 , 681-689 | | 56 |
| 375 | Novel rhodathiaborane complexes derived from [(PPh3)2RhSB9H10]. <i>Journal of the Chemical Society Chemical Communications</i> , 1990 , 891-894 | | 55 |
| 374 | Selective dismantling of closo-1,2-C2B10H12 to lower-cage dicarbaborane systems. <i>Collection of Czechoslovak Chemical Communications</i> , 1991 , 56, 1618-1635 | | 54 |
| 373 | Nuclear spin-spin coupling between tin and other directly bound elements. <i>Journal of the Chemical Society Dalton Transactions</i> , 1975 , 386-390 | | 52 |
| 372 | Magnetic double-resonance studies of platinum-195 chemical shifts in organoplatinum compounds. <i>Journal of the Chemical Society Dalton Transactions</i> , 1976 , 874 | | 52 |
| 371 | Magnetic double resonance studies of 119Sn chemical shifts in organotin compounds. <i>Journal of the Chemical Society C, Organic</i> , 1969 , 1136 | | 52 |
| 370 | Monocarbaborane anion chemistry. The elusive C-arylated [PhCB11H11]-, [PhCB9H9]- and [PhCB8H8]- anions. <i>Chemical Communications</i> , 2001 , 1790-1 | 5.8 | 51 |
| 369 | Metallaheteroborane chemistry. Part 3. Synthesis of [2,2-(PR3)2-1,2-TePtB10H10](R3= Et3, Bun3, or Me2Ph), their characterisation by nuclear magnetic resonance spectroscopy, and the crystal and molecular structure of [2,2-(PET3)2-1,2-TePtB10H10]. <i>Journal of the Chemical Society Dalton Transactions</i> , 1988 , 2555 | | 49 |

- 368 Open cluster configurations of [1,1-(PPh₃)₂-1-H-1,2,4-IrC₂B₈H₁₀] and other formally Wadlan 24-electron 11-vertex species. *Inorganic Chemistry*, **1989**, 28, 2219-2221 5.1 44
- 367 The molecular basis for sound velocity in n-alkanes, 1-alcohols and dimethylsiloxanes. *Physical Chemistry Chemical Physics*, **2003**, 5, 73-78 3.6 43
- 366 Polyhedral boron-containing cluster chemistry: Aspects of architecture beyond the icosahedron. *Pure and Applied Chemistry*, **2003**, 75, 1239-1248 2.1 42
- 365 Assigned cluster 11B and 1H NMR properties of [3-(η -C₆Me₆)-closo-3,1,2-RuC₂B₉H₁₁]. *Magnetic Resonance in Chemistry*, **1989**, 27, 947-949 2.1 42
- 364 An approach to megalo-boranes. Mixed and multiple cluster fusions involving iridaborane and platinaborane cluster compounds. Crystal structure determinations by conventional and synchrotron methods. *Inorganica Chimica Acta*, **1999**, 289, 95-124 2.7 41
- 363 Two-dimensional 1H-1H COSY n.m.r. spectroscopy in polyhedral boron hydride chemistry. *Journal of the Chemical Society Chemical Communications*, **1986**, 779-781 4.1
- 362 Quantitative ortho-cycloboronation of P-phenyl groups in metallaborane chemistry and the crystal and molecular structure of the novel iso-closo-ten-vertex metallaborane [1,1,1-H(PPh₃)(Ph₂P-ortho-C₆H₄)-iso-closo-(1-IrB₉H₈-2-)]. *Journal of the Chemical Society Chemical Communications*, **1982**, 465 4.0
- 361 Lead-207 shielding in organolead compounds. *Journal of the Chemical Society Dalton Transactions*, **1977**, 2332 4.0
- 360 Processing and Sol Chemistry of a Triol-Based Sol-Gel Route for Preparing Lead Zirconate Titanate Thin Films. *Journal of the American Ceramic Society*, **2004**, 83, 1914-1920 3.8 38
- 359 Group 1 coordination chains and hexagonal networks of host cyclotrimeratrylene with halogenated monocarbaborane anions. *Chemistry - A European Journal*, **2004**, 10, 2190-8 4.8 38
- 358 Structural chemistry of halogenated monocarbaboranes: the extended structures of Cs[1-HCB₉H₄Br₅], Cs[1-HCB₁₁H₅Cl₆] and Cs[1-HCB₁₁H₅Br₆]. *New Journal of Chemistry*, **2004**, 28, 1499-1505 3.6 38
- 357 Carbaborane salts of [ZnCl(HpztBu)₃]⁺, a host for inorganic anions (HpztBu = 5-tert-butylpyrazole). *New Journal of Chemistry*, **2002**, 26, 1634-1637 3.6 37
- 356 Metallheteroborane chemistry. Part 8. NMR study of some arsena- and stibaboranes and of the rhodadiarsenaboranes [3,3-(PPh₃)₂-3-(H)-closo-3,1,2-RhAs₂B₉H₉] and [3-(η -C₅Me₅)-closo-3,1,2-RhAs₂B₉H₉]. *Magnetic Resonance in Chemistry*, **1991**, 29, 711-720 2.1 36
- 355 An unusual direct conversion of metallaboranes to metallacarboranes and the isolation of a novel isoarachno twelve-vertex cluster compound. *Journal of the Chemical Society Chemical Communications*, **1990**, 1741-1743 3.6
- 354 The unexpected open cluster structure of [1-(η -6-MeC₆H₄CHMe₂-4)-2,4-Me₂-1,2,4-RuC₂B₈H₈]. *Organometallics*, **1987**, 6, 2254-2255 3.8 36
- 353 The Effect of Interbond Angles at Tin Upon 119Sn Chemical Shifts in Organotin Alkane- η -EDithiolates and Some Related Compounds. *Bulletin Des Sociétés Chimiques Belges*, **2010**, 84, 289-298 3.5
- 352 Monocarbaborane chemistry. Preparation and characterisation of [4-CB₈H₉]⁻, the 'missing' closo-carbaborane anion. *Chemical Communications*, **2001**, 1756-7 5.8 35
- 351 The first osmaboranes and a new iridatetraborane. *Journal of Organometallic Chemistry*, **1983**, 249, 11-21. 3.3 35

- 350 Monocarbaborane Anion Chemistry. Syntheses and Structures Within the closo Nine-Vertex System. *Collection of Czechoslovak Chemical Communications*, **2002**, 67, 1035-1050 34
- 349 Eleven-vertex polyhedral metalladecaborane chemistry. Reactions of neutral nido-5,6-C₂B₈H₁₂ and the [nido-6,9-C₂B₈H₁₀]²⁻ anion to give formally closo isomeric 1-(arene) and 1-(cyclopentadiene)-1,2,4- and 1,2,3-metalladecaboranes, and some substituted 2.3 34
- 348 Boron-11 and proton nuclear magnetic resonance study of anti-B₁₈H₂₂ and its anions, anti-[B₁₈H₂₁]⁻ and anti-[B₁₈H₂₀]²⁻. The crystal and molecular structure of [NMe₄]⁺[anti-B₁₈H₂₀]²⁻. *Journal of the Chemical Society Dalton Transactions*, **1988**, 1785-1793 34
- 347 Magnetic double resonance studies of some methyltin alkane- and benzene-thiolates. *Journal of the Chemical Society Perkin Transactions II*, **1974**, 146 34
- 346 Magnetic double resonance studies of tin-119 chemical shifts in compounds with tin-sulphur bonds and related species. *Journal of the Chemical Society Perkin Transactions II*, **1975**, 1234-1239 34
- 345 Thermochromic Fluorescence from B₁₈H₂₀(NC₅H₅)₂: An Inorganic/Organic Composite Luminescent Compound with an Unusual Molecular Geometry. *Advanced Optical Materials*, **2017**, 5, 1600694 8, 1 33
- 344 Polyhedral monocarbaborane chemistry. Some C-phenylated seven, eight, nine, ten, eleven and twelve-vertex species. *Dalton Transactions*, **2006**, 5753-69 4-3 33
- 343 Ten-Vertex Monocarbaborane Chemistry. A Convenient New Preparation of the [closo-1-CB₉H₁₀]⁻ Anion and the Crystal and Molecular Structure of [(C₅-C₅Me₅)₂Ir₂Cl₃]⁺[closo-1-CB₉H₁₀]⁻. *Collection of Czechoslovak Chemical Communications*, **1992**, 57, 1262-1268 33
- 342 Oxarhodaborane chemistry: the formation of [η-9,9?-O-{5-(C₅-C₅Me₅)-nido-5-RhB₉H₁₂}]₂ and [7-(C₅-C₅Me₅)-10-(NEt₃)-nido-7, 12-RhOB₁₀H₁₀] from [nido-(C₅-C₅Me₅)RhB₁₀H₁₃Cl]. *Journal of the Chemical Society Chemical Communications*, **1990**, 1692-1694 33
- 341 NMR Studies on Halogeno-cis-bis(phosphine)-nido-pentaboranyl Derivatives of Nickel, Palladium, and Platinum. *Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences*, **1979**, 34, 808-813 1 33
- 340 Facile thermally-induced cluster oxidations in metallaborane chemistry: arachno-nido-closo reaction sequences exhibited by iridanonaboranes and iridadecaboranes, and the stabilization of the iridium(V) oxidation state. *Journal of the Chemical Society Chemical Communications*, **1982**, 346-348 33
- 339 Solvent and counterion control of stereochemistry in a formal nucleophilic displacement reaction. Mechanisms of reaction of trimethyltin alkalis with alkyl halides. *Journal of the American Chemical Society*, **1972**, 94, 7206-7208 16.4 33
- 338 Polyhedral monocarbaborane chemistry. Routes to neutral, monoanionic and dianionic carbo-carbaborane rods. *Chemical Communications*, **2003**, 1684 5.8 32
- 337 Organoruthenaborane chemistry. Part 7. Preparation, molecular structure, and nuclear magnetic resonance properties of eleven-vertex closo-type clusters: [2,4-Me₂-1-(pcym)-1,2,4-RuC₂B₈H₈], [2,3-Me₂-1-(pcym)-1,2,3-RuC₂B₈H₈], [1-(pcym)-1-RuB₁₀H₁₀] and some related species. *Journal of the Chemical Society Dalton Transactions*, **1990**, 3039 31
- 336 Indirect nuclear spin spin coupling constants of tin-119 and lead-207 to nitrogen and phosphorus in organometallic compounds. *Journal of Organometallic Chemistry*, **1980**, 195, 285-290 2.3 31
- 335 Studies of ¹⁹⁹Hg nuclear shielding anisotropies and their relation to isotropic chemical shifts. *Journal of the Chemical Society, Faraday Transactions 2*, **1976**, 72, 1653 31
- 334 Nuclear magnetic double resonance studies of organotin selenides. *Journal of the Chemical Society Dalton Transactions*, **1973**, 2134 31
- 333 Macropolyhedral Boron-Containing Cluster Chemistry: Isolation and Characterization of the First Macropolyhedral Thiaborane, the arachno-Type [9,9?-S₂B₁₇H₁₈]³⁻ ion. *Angewandte Chemie International Edition in English*, **1994**, 33, 1599-1601 30

- 332 Metallaheteroborane chemistry. Part 5. Synthesis of closo-[2,2-(PR₃)₂-1,2-SePtB₁₀H₁₀](R₃= Et₃, Bun₃ or Me₂Ph) and arachno-[9,9-(PMe₂Ph)₂-6,9-SePtB₈H₁₀] and their characterisation by nuclear magnetic resonance techniques: the crystal and molecular structures of closo-[2,2-(PEt₃)₂-1,2-SePtB₁₀H₁₀] and arachno-[9,9-(PMe₂Ph)₂-6,9-SePtB₈H₁₀]. *Journal of the Chemical Society Dalton Transactions*, **1988**, 1657 30
- 331 Preparation, structure, and nuclear magnetic resonance properties of the nine-vertex nido-rhenaborane [(PMe₂Ph)₃H₂ReB₈H₁₁] and some related chemistry. *Journal of the Chemical Society Dalton Transactions*, **1988**, 1969 30
- 330 Two unusual closo-type ruthenaboranes: preparation, molecular structure, and nuclear magnetic resonance properties of [1,1,1-(PPh₃)HCl-1-RuB₉H₇-3,5-(PPh₃)₂] and [1,1-(PPh₃)₂-1-RuB₁₀H₈-2,5-(OEt)₂]. *Journal of the Chemical Society Dalton Transactions*, **1985**, 2407 30
- 329 The chemistry of isomers of icosaborane(26), B₂₀H₂₆: synthesis and nuclear magnetic resonance study of various isomers of platinahenicosaboranes and diplatinadocosaboranes, and the X-ray crystal and molecular structures of 7,7-bis(dimethylphenylphosphine)-nido-7-platinaundecaborane [Pt₂(C₆H₅)₂(PMe₂Ph)₂]-7-PtB₁₁H₁₂. *Journal of the Chemical Society Dalton Transactions*, **1985**, 1969 30
- 328 A novel oxidative insertion of a metal centre into a degraded closo-borane cluster; crystal and molecular structure of the ten-vertex iso-nido cluster [IrC(OH)B₈H₆(OMe)}(C₆H₄PPH₂)(PPh₃)]. *Journal of the Chemical Society Chemical Communications*, **1981**, 933 30
- 327 New modes of bonding in some platinum derivatives of B₁₈H₂₂: X-ray crystal structures of [(Pt₂B₁₈H₁₆)(PMe₂Ph)₄] and of three structural isomers of [(PtB₁₈H₂₀)(PMe₂Ph)₂]. *Journal of the Chemical Society Chemical Communications*, **1982**, 80 30
- 326 Eleven-Vertex Polyhedral Dicarbaplatinaborane Chemistry. Aspects of the Chemistry of Some closo-Structured {1,2,3-PtC₂B₈} Species and Some Related Compounds. *Collection of Czechoslovak Chemical Communications*, **1993**, 58, 2090-2120 30
- 325 Crystal-packing motifs of [Ag₄L₄]⁴⁺ star-burst tetrahedra. *New Journal of Chemistry*, **2006**, 30, 1390 3.6 29
- 324 Polyhedral monocarbaborane chemistry. *Journal of Organometallic Chemistry*, **2002**, 657, 176-179 2.3 29
- 323 Phosphacarborane chemistry. New cluster isomers in the eleven-vertex nido-phosphadecaborane series: synthesis of the nido phosphadecaboranes 7,8,11-PC₂B₈H₁₁, [7,8,11-PC₂B₈H₁₀] and 7-Ph-7,8,10-PC₂B₈H₁₀. *Inorganic Chemistry Communication*, **2000**, 3, 178-181 3.1 29
- 322 Reactions of 6,6'-bis(nido-decaboranyl) oxide and 6-hydroxy-nido-decaborane with dihalogenobis(phosphine) complexes of nickel, palladium, and platinum, and some related chemistry; nuclear magnetic resonance investigations and the crystal and molecular structures of bis(dimethylphosphine)-di-μ-(2,3,4-B-nido-hexaboranyl)-diplatinum(Pt₂B₁₀H₁₂) and bis(dimethylphosphine)-di-μ-(2,3,4-B-nido-hexaboranyl)-diplatinum(Pt₂B₁₀H₁₂). *Journal of the Chemical Society Dalton Transactions*, **1985**, 953-972 29
- 321 An unusual open twelve-vertex oxametallaborane cluster compound: synthesis and structure of [7-(B-C₅Me₅)-8-Cl-11-(PMe₂Ph)-nido-7,12-RhOB₁₀H₉]. *Journal of the Chemical Society Chemical Communications*, **1985**, 1722-1723 29
- 320 Silver-Dabco Coordination Networks with Distinct Carbaborane Anions: Investigating Ag^{III}HB and Ag^{III}B Interactions. *Crystal Growth and Design*, **2013**, 13, 3162-3170 3.5 28
- 319 Monocarbaborane anion chemistry. An interesting encapsulation of the Pd₂I₂(P(C₆H₄(4-Me)₃)₄)₂⁺ cation by a pair of [PhCB₉H₄I(C₆H₄Me)₄]⁻ anions. *Chemical Communications*, **2002**, 2048-9 5.8 28
- 318 Metallaborane Heteroatom Incorporation Reactions: Enyne Insertion into arachno-[(CO)(PMe₃)₂HrB₈H₁₂]. *Organometallics*, **1998**, 17, 902-907 3.8 28
- 317 Synthesis, molecular structures, and nuclear magnetic resonance properties of the macropolyhedral metallaboranes [(PMe₂Ph)₄Pt₃B₁₄H₁₆] and [(PMe₂Ph)PtB₁₆H₁₈(PMe₂Ph)], and a discussion of the bonding at platinum in these and some related platinaborane clusters. *Journal of the Chemical Society Dalton Transactions*, **1991**, 1079 28
- 316 Macropolyhedral boron-containing cluster chemistry. Nineteen-vertex [S₂B₁₇H₁₇(SMe₂)]. An unusual apical boron atom of cluster connectivity six that introduces a new polyhedral borane building block. *Chemical Communications*, **1996**, 273-275 5.8 27
- 315 POLYHEDRAL OXABORANE CHEMISTRY. SOME COMPARATIVE ¹¹B NMR SHIELDING PATTERNS WITHIN THE TWELVE-VERTEX NIDO-TYPE GEOMETRY. *Main Group Metal Chemistry*, **1996**, 19, 1-10 1.6 27

- 314 Ten-vertex metallaborane chemistry: facile, thermally induced, nido-isocloso cluster-closure oxidation reactions in iridadecaborane clusters. *Journal of the Chemical Society Dalton Transactions*, **1990**, 1451-1458 27
- 313 Hexamethylbenzene-ruthenaborane and -osmaborane chemistry. *Journal of Organometallic Chemistry*, **1986**, 309, C67-C69 2.3 27
- 312 Organo-ruthenaborane and -osmaborane cluster compounds: preparation and NMR studies of [6-(η -C₆Me₆)-nido-6-MB₉H₁₃] (M = Ru, Os) and the rearranged cluster [5-(η -C₆Me₆)-nido-5-RuB₉H₁₁-7-(PMe₂Ph)]. *Journal of Organometallic Chemistry*, **1987**, 325, 233-246 2.3 27
- 311 Polyhedral thiaborane chemistry: a comparative nuclear magnetic resonance study of arachno-[6-SB₉H₁₂] and nido-6-SB₉H₁₁. *Journal of the Chemical Society Dalton Transactions*, **1988**, 1467 27
- 310 Polyhedral monocarbaborane chemistry: The closo-[PhCB₇H₇] and closo-[PhCB₁₀H₁₀] anions: the two missing species in the closo-[PhCB_nH_n] sequence. *Journal of Organometallic Chemistry*, **2002**, 657, 180-186 2.3 26
- 309 Eight-vertex polyhedral monocarbaborane chemistry. Three closo anions, [CB₇H₈], [CB₇H₇] and [CB₇H₆I₂]. Preparation and structural studies. *Journal of the Chemical Society Dalton Transactions*, **1995**, 431-437 26
- 308 Metallaheteroborane chemistry. Part 11. Selective syntheses of the palladium heteroborane complexes [2,2-(PR₃)₂-closo-2,1-PdEB₁₀H₁₀](R₃ = Ph₃, MePh₂ or Me₂Ph; E = Se or Te) and [2-X-2-(PPh₃)-closo-2,1-PdTeB₁₀H₉(PPh₃)](X = Cl, Br, I, CN, SCN or O₂CMe). *Journal of the Chemical Society Dalton Transactions*, **1993**, 27 26
- 307 A novel closo-type metallaundecaborane, [1,1-(PPh₃)₂-2,5-(OEt)₂-1-RuB₁₀H₈], and its relationship to the η -ocloso series of metallaborane clusters. *Polyhedron*, **1984**, 3, 901-904 2.7 26
- 306 Macropolyhedral boron-containing cluster chemistry. Assessment of the possibilities of thermolytic mixed-cluster fusion, and of the use of synchrotron X-radiation for the examination of small single crystals of metallaboranes. Isolation and structure of [18-vertex-7-(CO)-7,7-(PMe₃)₂-syn-7-IrB₁₇H₂₀]. *Journal of the Chemical Society Dalton Transactions*, **1997**, 2405-2413 25
- 305 Macropolyhedral boron-containing cluster chemistry. Triple cluster fusion and the molecular structure of [(PMe₃)₂IrB₂₆H₂₄Ir(CO)(PMe₃)₂]. A 28-vertex metallaborane cluster with a polyboron core. *Chemical Communications*, **1997**, 2405-2406 5.8 24
- 304 Polyhedral azadiborane chemistry. The η -inverse synthesis of a small family of 6,9-(NHR)-bridged-arachno-5,10-dicarbododecaboranes. *Inorganica Chimica Acta*, **2000**, 304, 268-273 2.7 24
- 303 Dicarbaheteroborane Chemistry. Representatives of Two Eleven-Vertex Dicarbaundecaborane Families: nido-10,7,8-NC(2)B(8)H(11), Its N-Substituted Derivatives, and arachno-1,8,11-NC(2)B(8)H(13). *Inorganic Chemistry*, **1998**, 37, 3902-3909 5.1 24
- 302 Macropolyhedral boron-containing cluster chemistry: nineteen-vertex [(PPh₃)NiS₂B₁₆H₁₂(PPh₃)] and eighteen-vertex S₂B₁₆H₁₄(PPh₃). *Journal of the Chemical Society Dalton Transactions*, **1996**, 4155 24
- 301 Metallaborane reaction chemistry. Part 1. Two interesting closed cluster compounds from the reaction of acetylene with an open nido-6-iridadecaborane. *Journal of the Chemical Society Dalton Transactions*, **1993**, 2335 24
- 300 Preparation of Smaller Nine- and Ten-Vertex Monocarbaborane Cages by the Selective Dismantling of Twelve-Vertex Dicarboranes via the [nido-7,9-C₂B₁₀H₁₃]⁻ Anion. Synthesis of the Monocarbaborane Base Adducts endo-9-Me-8-(NMe₃)-arachno-6-CB₉H₁₂, 9-Me-8-L-nido-6-CB₉H₁₀ (l = NMe₃, OH⁻) and exo-6-1-arachno-4-CB₈H₁₂ (l = SMe₂, NMe₃, Urotropine, Pyridine, Quinoline, Me₃N⁺, Ph₃Me₃⁺). *Inorganic Chemistry*, **1994**, 33, 3994-3997 5.1 24
- 299 Macropolyhedral boron-containing cluster chemistry. Cluster fusion to give the novel nido-18-vertex dithiooctadecaborane (anti)-[9,9'-S₂B₁₆H₁₆]. *Journal of the Chemical Society Chemical Communications*, **1994**, 1415-1416 24
- 298 The isolation and characterisation of the 1- and 2-isomers of nido-[(η -C₆Me₆)RuB₉H₁₃] but is the 1-isomer nido or arachno?. *Journal of the Chemical Society Chemical Communications*, **1987**, 817-818 24
- 297 Synthesis, molecular structure, and nuclear magnetic resonance investigation of the platinumaborane and diplatinadecaborane cluster compounds, [4,4-(PMe₂Ph)₂-arachno-4-PtB₈H₁₂] and [6,6,9,9-(PMe₂-Ph)₄-arachno-6,9-Pt₂B₈H₁₀]. *Journal of the Chemical Society Dalton Transactions*, **1991**, 1415 24

- 296 Metallaheteroborane chemistry. *Journal of Organometallic Chemistry*, **1998**, 550, 151-164 2.3 23
- 295 Polyhedral oxaplatinaborane chemistry: characterisation of [9,9-(PMe₂Ph)₂-arachno-9,6-PtOB8H₁₀] and its metallaborane non-oxa cognate [6,6-(PMe₂Ph)₂-arachno-6-PtB9H₁₁-9-(PMe₂Ph)]. *Inorganic Chemistry Communication*, **1998**, 1, 19-22 3.1 23
- 294 Metallaborane reaction chemistry. Macropolyhedral metallaheteroborane synthesis by direct heteroatom insertion. Formation of the twenty-vertex [(B-C5Me5)IrB18H19S]⁻ anion from nineteen-vertex syn-[(B-C5Me5)IrB18H20]. *Inorganic Chemistry Communication*, **1998**, 1, 97-100 3.1 23
- 293 Macropolyhedral boron-containing cluster chemistry. The [S2B18H19]⁻ anion, and the reversible dismantling and regeneration of an apical boron cluster site with cluster connectivity six. *Journal of the Chemical Society Dalton Transactions*, **1998**, 2965-2968 23
- 292 Metallaheteroborane chemistry. Part 10. Synthesis and characterisation of closo-structured rhodathiaborane complexes [1-(CO)-1-L-3-L?-1,2-RhSB9H8](L = L? = PPh₃; L = PMe₂Ph, L? = PMe₂Ph or PPh₃). *Journal of the Chemical Society Dalton Transactions*, **1992**, 2865-2871 23
- 291 Iridanonaborane chemistry. Some reactions of arachno-4-irida-nonaboranes: nuclear magnetic resonance investigations and the molecular structure of the nido nine-vertex species [2,2,2-(CO)(PMe₃)₂-nido-2-IrB8H11]. *Journal of the Chemical Society Dalton Transactions*, **1984**, 2477 23
- 290 A proton and boron-11 NMR study of icosaborane oxide, B₂₀H₂₆O. *Inorganica Chimica Acta*, **1980**, 38, 93-96 2.7 23
- 289 Sign and magnitude of ¹K(Sn^{IV}) in trimethylstannyl-lithium: the first example of a negative reduced nuclear spin-spin coupling constant between two directly bound Group IV elements. *Journal of the Chemical Society Chemical Communications*, **1974**, 983-984 23
- 288 Auto-association in organometallic compounds: a nuclear magnetic resonance study of methylcadmium alkoxides. *Journal of the Chemical Society Perkin Transactions II*, **1977**, 1187 23
- 287 Magnetic double-resonance measurements of rhodium, phosphorus, fluorine, and proton nuclear magnetic resonance parameters in some rhodium(I) and rhodium(III) compounds. *Journal of the Chemical Society Dalton Transactions*, **1977**, 1571 23
- 286 Isolation and Structure of [(PPh₃)₃(PPh₂)₂Pd₄B₂₀H₁₆]. A Possible Prognostic for New Globular Borane-Based Cluster Architectures. *Collection of Czechoslovak Chemical Communications*, **1999**, 64, 927-937 23
- 285 Monocarbaborane anion chemistry. [COOH], [CH₂OH] and [CHO] units as functional groups on ten-vertex monocarbaborane anionic compounds. *Dalton Transactions*, **2004**, 3552-61 4.3 22
- 284 Polyhedral monocarbaborane chemistry: positional isomerism within the 10-vertex monocarbaborane cluster: synthesis and characterisation of the [2-Ph-closo-2-CB9H9]⁻ anion. *Inorganic Chemistry Communication*, **2002**, 5, 581-584 3.1 22
- 283 Conformational polymorphism and fluxional behaviour of M(PR₃)₂ units in closo-twelve-atom metallaheteroboranes with MX₂B₉(X = C or As) and MZB₁₀ cages (Z = S, Se or Te). *Journal of the Chemical Society Dalton Transactions*, **1996**, 3323-3333 22
- 282 A unique acetylene-induced reductive PPh₃-iPH₃ stripping of phenyl groups from triphenylphosphine ligands bound to a metallaborane B-frame matrix. *Journal of the Chemical Society Chemical Communications*, **1989**, 1763-1765 22
- 281 Ten-vertex metallaborane chemistry: synthesis and characterization of some ortho-cycloboronated nido-5- and -6-iridadecaboranes; crystal structures of [5-H-5-(PPh₃)-5-(PPh₂-o-C₆H₄)-nido-5-IrB9H12-2] and [5-H-5,7-(PPh₃)₂-5-(PPh₂-o-C₆H₄)-nido-5-IrB9H12-2]. *Journal of the Chemical Society Dalton Transactions*, **1990**, 1441 22
- 280 Silicon, Germanium, Tin, and Lead **1987**, 305-333 22
- 279 Ruthenium(II) complexes of closo-dodecaboranyl anions and the molecular structure of the pileo thirteen-vertex ruthenaborane [(PPh₃)₂ClRuB₁₂H₁₁(NEt₃)]. *Journal of the Chemical Society Dalton Transactions*, **1987**, 451 22

- 278 Preparation of [(C5Me5)2Rh2B17H19] via a degradative insertion from anti-B18H22, and a possible mechanism for anti-syn macropolyhedral interconversion. *Journal of the Chemical Society Chemical Communications*, **1986**, 1111-1113 22
- 277 Auto-association in organometallic compounds: a nuclear magnetic double resonance study of methyl- and n-butyl-tin alkoxides. *Journal of the Chemical Society Perkin Transactions II*, **1977**, 242 22
- 276 Auto-association in organotin compounds: butyltin trialkoxides. *Journal of the Chemical Society Perkin Transactions II*, **1973**, 1785 22
- 275 Borane reaction chemistry. Alkyne insertion reactions into boron-containing clusters. Products from the thermolysis of [6,9-(2-HC[triple bond]C-C5H4N)2-arachno-B10H12]. *Dalton Transactions*, **2008**, 1552-63 4.3 21
- 274 Pentahapto-bonded gold heteroborane clusters [3-(R3P)-closo-2,1-AuTeB10H10]- and [3-(R3P)-closo-3,1,2-AuAs2B9H9]-. *Dalton Transactions*, **2006**, 2133-9 4.3 21
- 273 Macropolyhedral boron-containing cluster chemistry—the nineteen-vertex oxaborane anion, [OB18H21]. *Journal of the Chemical Society Chemical Communications*, **1995**, 1665-1666 21
- 272 Organoruthenaborane Chemistry. VIII. Reactions of [(η -C6Me6)RuCl2]2 and [(η -MeC6HPr)RuCl2]2 with Cs[arachno-6-SB9H12]: Isolation of ten-, eleven-, and twelve-vertex ruthenathaboranes and their characterization by N.M.R. spectroscopy. *Zeitschrift Fur Anorganische Und Allgemeine Chemie*, **1991**, 602, 17-29 1.3 21
- 271 An interesting low barrier to metal-ligand contrarotational fluxionality in closed 11-vertex [1,1-(PMe2Ph)2-1,2,3-PtC2B8H9X] compounds that is characterized by arachno .dblharw. nido geometric changes within the 10-vertex .eta.6-{C2B8H9X} ligand. *Inorganic Chemistry*, **1991**, 30, 4481-4484 5.1 21
- 270 Comparative metallaborane chemistry: preparation and nuclear magnetic resonance studies of some nido-5-metalladecaboranes of rhodium, iridium, ruthenium, and osmium. *Journal of the Chemical Society Dalton Transactions*, **1988**, 2597 21
- 269 Molecular structure of the 17-vertex conjuncto-platinaheptadecaborane [(PMe2Ph){PtB16H18(PMe2Ph)}]: a 16-vertex η -hexadecaboranyl ligand. *Journal of the Chemical Society Chemical Communications*, **1982**, 552-553 21
- 268 Magnetic double resonance studies of trimethyltin chalcogenides. *Journal of Organometallic Chemistry*, **1975**, 94, 7-14 2.3 21
- 267 Substitution of the laser borane anti-BH with pyridine: a structural and photophysical study of some unusually structured macropolyhedral boron hydrides. *Dalton Transactions*, **2018**, 47, 1709-1725 4.3 20
- 266 Metallaborane reaction chemistry. A facile and reversible dioxygen capture by a B-frame-supported bimetallic: structure of [(PMe2Ph)4(O2)Pt2B10H10]. *Chemical Communications*, **2004**, 2380-1 5.8 20
- 265 Metallaborane reaction chemistry. Part 12. Some interactions of acetylenes and isocyanides with selected metallaboranes. *Journal of Organometallic Chemistry*, **2005**, 690, 2701-2720 2.3 20
- 264 NMR Assignments of [6-R-nido-5,6-C2B8H10]- Anions (where R = H, Me, and n-C6H13). An Irreversible 5- \rightarrow 6 Alkyl Migration via a B9 Vertex-Swing Mechanism. *Collection of Czechoslovak Chemical Communications*, **1999**, 64, 986-992 20
- 263 Macropolyhedral boron-containing cluster chemistry. syn- and anti-[(C5Me5)2Rh2S2B15H14(OH)]; an interesting nineteen-vertex isomeric pair. *Journal of the Chemical Society Dalton Transactions*, **1996**, 1775-1777 20
- 262 Metallaheteroborane chemistry. Part 12. Synthesis of cationic metallaheteroboranes [2-L-2-(PPh3)-closo-2,1-PdTeB10H9(PPh3)][BF4]; molecular structures of the compounds with L = H2O or CO. *Journal of the Chemical Society Dalton Transactions*, **1993**, 35 20
- 261 A novel exo-bicyclic closo-six-vertex dimetalla-hexaborane : [1,1,2-(CO)3-1-(PPh3)-2,2-(Ph2PC6H4)2-closo-(1,2-Ir2B4H2-3,5-)]. CH2Cl2. *Journal of the Chemical Society Chemical Communications*, **1982**, 383 20

- 260 The magnitude and sign of $1J(207\text{Pb}-207\text{Pb})$ in hexamethyldecylidene. *Journal of Organometallic Chemistry*, **1974**, 80, C47-C49 2.3 20
- 259 A Return to the Plešek Reaction and Some Useful Variations. Carbon-Substituted Methyl and Phenyl Derivatives of 5,6-Dicarbonyl-nido-decaborane(12), nido-5,6-C₂B₈H₁₂. *Collection of Czechoslovak Chemical Communications*, **1997**, 62, 1229-1238 19
- 258 Ten-vertex monocarbaborane chemistry. Synthesis of carbon-substituted ligand derivatives 1-L-closo-1-CB₉H₉ and crystal and molecular structure of 1-(Me₃N)-closo-1-CB₉H₉. *Journal of the Chemical Society Dalton Transactions*, **1997**, 4231-4236 19
- 257 Polyhedral metallaheteroborane chemistry. Synthesis, spectroscopy, structure and dynamics of eleven-vertex {RhNB(9)} and {PtCB(9)} metallaheteroboranes. *Dalton Transactions*, **2007**, 2885-97 4.3 19
- 256 Dimethylsulfide-dicarbaborane chemistry. Isolation and characterisation of isomers [9-(SMe₂)-nido-7,8-C₂B₉H₁₀-X-Me] (where X = 1, 2, 3 and 4) and some related compounds. An unusual skeletal rearrangement. *Dalton Transactions*, **2007**, 4859-65 4.3 19
- 255 Polyhedral monocarbaborane chemistry. Carboxylic acid derivatives of the [closo-2-CB₉H₁₀]- anion. *Chemical Communications*, **2004**, 328-9 5.8 19
- 254 B-frame-supported bimetallics. Isoelectronic arachno-structured [(PMe₂Ph)₄Pd₂B₈H₁₀] and closo-structured [(PMe₃)₄(CO)₂Ir₂B₈H₈]. *Inorganica Chimica Acta*, **1999**, 285, 290-295 2.7 19
- 253 The Parent Hexacarbaborane arachno-C₆B₆H₆ and a Methylated Pentacarbaborane arachno-CH₂C₅B₅H₅: Domains of Incipient Hydrocarbon Behavior within Borane Clusters. *Angewandte Chemie - International Edition*, **1999**, 38, 1806-1808 16.4 19
- 252 Macropolyhedral boron-containing cluster chemistry. The [SB₁₇H₁₉]⁻ anion: a nido-ten-vertex : arachno-ten-vertex cluster architecture and the first single-sulfur macropolyhedral thiaborane. *Chemical Communications*, **1999**, 1905-1906 5.8 19
- 251 Macropolyhedral boron-containing cluster chemistry. Isolation and characterisation of the eighteen-vertex nido-5⁺-iridaoctaborano[3⁺,8⁺:1⁺,2⁺]-closo-4-iridadodecaborane, [(CO)(PMe₃)₂IrB₁₆H₁₄Ir(CO)(PMe₃)₂]. *Journal of the Chemical Society Dalton Transactions*, **1996**, 3145-3149 19
- 250 Some thought-provoking polyhedral borane chemistry. *Pure and Applied Chemistry*, **1991**, 63, 317-326 2.1 19
- 249 Organoruthenaborane chemistry. Part 5. Products of the reaction between closo-[B₁₀H₁₀]₂ and [(C₆Me₆)ClRuB₃H₈]. Nuclear magnetic resonance studies and the crystal and molecular structure of [(C₆Me₆)₂Ru₂H₄]RuB₁₀H₈(OEt)₂. *Journal of the Chemical Society Dalton Transactions*, **1997**, 2761-2767 19
- 248 Iridanonaborane chemistry: preparation and characterization of some arachno-4-iridanonaboranes by nuclear magnetic resonance spectroscopy and single-crystal X-ray diffraction analysis. *Journal of the Chemical Society Dalton Transactions*, **1984**, 1903 19
- 247 The crystal and molecular structure of [1,1,1,1-(CO)H(PPh₃)₂-arachno-1-IrB₃H₇] and some bonding considerations in arachno-type four-vertex metal allyl and borallyl clusters. *Journal of the Chemical Society Dalton Transactions*, **1985**, 1843-1847 19
- 246 Cage expansion in metallo-pentaborane chemistry: the preparation and structure of 2-{carbonylbis(triphenylphosphine)irida}-nido-hexaborane, [(IrB₅H₈)(CO)(PPh₃)₂]. *Journal of the Chemical Society Dalton Transactions*, **1979**, 117 19
- 245 Magnetic double resonance studies of carbon-13 and nitrogen-15 chemical shift anisotropies in methyl cyanide partially oriented in a nematic liquid crystal. *Molecular Physics*, **1975**, 29, 593-597 1.7 19
- 244 Dicarbonyl(η⁵-cyclopentadienyl)iron(II) derivatives of pentaborane(9). *Journal of the Chemical Society Dalton Transactions*, **1978**, 237-244 19
- 243 Addition of trimethyltin hydride and methyl halotin hydrides to norbornadiene. *Journal of Organic Chemistry*, **1971**, 36, 2083-2088 4.2 19

- 242 Macropolyhedral boron-containing cluster chemistry: isolation and characterization of twenty-one-vertex $[(PMe_2Ph)_3HR_2B_{20}H_{15}Ph(PHMe_2)]$. *Chemical Communications*, **1997**, 217-218 5.8 18
- 241 The capture of dioxygen, carbon monoxide and sulfur dioxide by $[(PMe_2Ph)_4Pt_2B_{10}H_{10}]$. *Dalton Transactions*, **2005**, 1574-82 4.3 18
- 240 Structural chemistry of arachno-nonaboranes. *Journal of the American Chemical Society*, **2002**, 124, 7429-34 18
- 239 Polyhedral carbathaborane chemistry. Ten-vertex nido and arachno, eleven-vertex nido, and eight-vertex hypho families of carbathaboranes. *Journal of the Chemical Society Dalton Transactions*, **1994**, 1317-1323 18
- 238 The high-yield preparation of arachno-thiaboranes 4-SB $8H_{12}$ and 4-SB $8H_{11}$. *Polyhedron*, **1989**, 8, 2089-2090 18
- 237 Heterobimetallic B-frame complexes: novel isonido eleven-vertex rhodairidaboranes with $Rh_{11}Ir$ and $B_{11}B$ bridges. *Journal of the Chemical Society Chemical Communications*, **1989**, 455-456 18
- 236 Arachno-6,9-diazadecaborane(12), $[6,9-N_2B_8H_{12}]$, an unsubstituted contiguous cluster diazaborane. *Journal of the Chemical Society Chemical Communications*, **1990**, 1309-1310 18
- 235 Pentamethylcyclopentadienylrhodaborane chemistry. Part 3. The reaction of $[(Rh(\eta-C_5Me_5)Cl)_2]$ with closo- $[B_{10}H_{10}]_2$ and arachno- $B_{10}H_{12}-6,9-(PMe_2Ph)_2$, and the characterisation of some novel rhodaboranes by X-ray diffraction analysis and nuclear magnetic resonance spectroscopy. *Journal of the Chemical Society Dalton Transactions*, **1987**, 2117-2120 18
- 234 Facile pathway-defined fluxional cluster isomerization in ten-vertex closo-2,1,6-metalladecaboranes of ruthenium and rhodium. *Journal of the Chemical Society Chemical Communications*, **1988**, 974-975 18
- 233 closo-Nido-Cluster isomerism in metallacarbaboranes via a change of metal valency state: molecular structure of $[\eta-1,2-acetato-2-hydrido-2,10-bis(triphenylphosphine)-closo-1,2-carbairidadecaborane]$. *Journal of the Chemical Society Chemical Communications*, **1989**, 93-94 18
- 232 Preparation, crystal and molecular structure of, and NMR parameters for, the exopolyhedral heterocyclic platinaundecaborane $[Pt_2,7-(SCSNET_2)-7-(PMe_2Ph)-NIDO-7-PtB_{10}H_{11}]$. *Polyhedron*, **1985**, 4, 505-511 2.7 18
- 231 Reactions of the octahydrotriborate(1-) anion, $[B_3H_8]^-$ with some complexes of cobalt(I), cobalt(II), rhodium(I), and iridium(I), and the characterization of the borallyl complex $[IrIII(\eta-B_3H_7)(CO)H(PPh_3)_2]$. *Journal of the Chemical Society Dalton Transactions*, **1980**, 196-200 18
- 230 Macropolyhedral boron-containing cluster chemistry. $[PtMe_2(PMe_2Ph)_2]$ as a cluster metallating agent. Isolation and characterisation of nineteen-vertex $[(PMe_2Ph)HPT_4-syn-B_{18}H_{19}(PMe_2Ph)]$ and eighteen-vertex $[(PMe_2Ph)_2Pt_2S_2B_{15}H_{14}(NHCOMe)]$. *Journal of Organometallic Chemistry*, **1998** 557, 181-185 2.3 17
- 229 Macropolyhedral boron-containing cluster chemistry. An interesting angular change in relative subcluster orientation in the oxidative protonation of $[S_2B_{17}H_{18}]^-$ to $[S_2B_{17}H_{17}]$. *Inorganic Chemistry Communication*, **1998**, 1, 179-181 3.1 17
- 228 Monocarbaborane anion chemistry: use of an anilinyll residue as an anchor group for the construction of rod-like monocarbaboranyl diazo and imino compounds. *Inorganic Chemistry Communication*, **2003**, 6, 1104-1108 3.1 17
- 227 Polyhedral Oxaruthenaborane Chemistry. Characterisation of a $[(\eta-C_6Me_6)RuOB_9H_{13}]$ Species of arachno Eleven-Vertex Cluster Character and Other Aspects of Oxaborane Chemistry. *Collection of Czechoslovak Chemical Communications*, **2005**, 70, 410-429 17
- 226 Macropolyhedral boron-containing cluster chemistry. Isolation and structure of the twenty-one-vertex globular cluster compound $[(\eta-C_5Me_5)_3Ir_3B_{18}H_{15}(OH)]$. *Inorganic Chemistry Communication*, **2000**, 3, 169-172 3.1 17
- 225 Two new ten-vertex arachno mixed-atom diheteroboranes, 6,9-CNB $8H_{13}$ and 6,9-CSB $8H_{12}$. *Journal of the Chemical Society Chemical Communications*, **1991**, 1389-1390 17

- 224 A macropolyhedral polymetallaborane cluster: molecular structure of the seventeen-vertex triplatinaheptadecaborane [(PhMe₂P)₄Pt₃B₁₄H₁₆]. *Journal of the Chemical Society Chemical Communications*, **1983**, 1228 17
- 223 Pentamethylcyclopentadienylrhodaborane chemistry. Part 1. High-yield planned synthesis, molecular structure, and nuclear magnetic resonance properties of the ten-vertex nido-6-rhodadecaborane [(η -C₅Me₅)RhB₉H₁₃]. *Journal of the Chemical Society Dalton Transactions*, **1986**, 547-552 17
- 222 Nuclear spin-spin coupling between directly bound tin atoms studied by a novel form of INDOOR spectroscopy. *Journal of the Chemical Society Dalton Transactions*, **1976**, 1219-1223 17
- 221 Reversible capture of small molecules on bimetallaborane clusters: synthesis, structural characterization, and photophysical aspects. *Inorganic Chemistry*, **2011**, 50, 7511-23 5.1 16
- 220 An experimental solution to the "missing hydrogens" question surrounding the macropolyhedral 19-vertex boron hydride monoanion [B₁₉H₂₂]⁻, a simplification of its synthesis, and its use as an intermediate in the first example of syn-B₁₈H₂₂ to anti-B₁₈H₂₂ isomer conversion. *Inorganic Chemistry*, **2010**, 49, 4092-8 5.1 16
- 219 Polyhedral ruthenaborane chemistry. The ten-vertex isocloso cluster geometries of [(pcym)RuB₉H₉] and [(pcym)Ru₂B₈H₈]. *Inorganic Chemistry Communication*, **1998**, 1, 375-378 3.1 16
- 218 Metallaborane reaction chemistry. A predicted and found tailored facile and reversible capture of SO₂ by a B-frame-supported bimetallic: structures of [(PMe₂Ph)₂PtPd(phen)B₁₀H₁₀] and [(PMe₂Ph)₂Pt(SO₂)Pd(phen)B₁₀H₁₀]. *Chemical Communications*, **2008**, 2447-9 5.8 16
- 217 Reactions of carbon disulfide with open nido-6-iridadecaboranes. The formation of closed ten-vertex cluster compounds with boron-to-metal dithioformate bridges and a novel isocloso-closo cluster conversion. *Journal of the Chemical Society Dalton Transactions*, **1996**, 3819-3824 16
- 216 Eleven-vertex polyhedral monocarbaplatinaborane chemistry. Structure and fluxionality of [8,8-(PMe₂Ph)₂-nido-8,7-PtCB₉H₁₁]. *Journal of the Chemical Society Dalton Transactions*, **1993**, 1261-1267 16
- 215 Synthesis, molecular structure and NMR behaviour of [9,9-(PMe₂Ph)₂-arachno-9,6,8-PtS₂B₇H₇]: Direct evidence for the ten-vertex borane vertex-flip mechanism of isomerization. *Journal of Organometallic Chemistry*, **1993**, 445, C15-C16 2.3 16
- 214 Pentamethylcyclopentadienylrhodaborane chemistry. Part 2. The reaction of [6-(η -C₅Me₅)-nido-6-RhB₉H₁₃] with dimethylphenylphosphine, and the characterization of [5-(η -C₅Me₅)-nido-5-RhB₉H₁₁-7-(PMe₂Ph)], [2-(η -C₅Me₅)-closo-2-RhB₉H₇-3,10-(PMe₂Ph)₂],* and [2-(η -C₅Me₅)-nido-2-RhB₈H₁₀-5-(PMe₂Ph)] by X-ray diffraction analysis and nuclear magnetic resonance spectroscopy. *Journal of the Chemical Society Dalton Transactions*, **1987**, 1431-1443 16
- 213 B-frame supported heterobimetallic species; molecular structure of [(Me₃P)₂Pt(Ph₃P)(Ph₂PC₆H₄)HfB₉H₁₀]. *Journal of the Chemical Society Chemical Communications*, **1983**, 949 16
- 212 Eleven-vertex isocloso type rhodaundecaboranes: crystal structures and nuclear magnetic resonance properties of [(PMe₂Ph)₂RhHB₁₀H₈(OMe)₂] and [(PMe₂Ph)₂RhHB₁₀H₈Cl(OMe)]. *Journal of the Chemical Society Dalton Transactions*, **1986**, 517 16
- 211 A new route to isomers of icosaborane(26), B₂₀H₂₆. The use of 115.5-MHz ¹¹B and ¹¹B-{¹H} nuclear magnetic resonance spectroscopy for the comparison and characterisation of separated isomers and the identification of three further icosaboranes as 1,2?-, 2,5?-, and 5,5?(or 5,7?)-icosaborane. *Journal of the Chemical Society Dalton Transactions*, **1991**, 1489-1497 16
- 210 Preparation and properties of some trimethyltin-substituted bicyclo[2.2.1] carbocycles, 3-nortricyclyltrimethyltin and norborn-2-en-7-yltrimethylsilanes. *Journal of Organometallic Chemistry*, **1973**, 61, 167-177 2.3 16
- 209 Effects of metal-centre orbital control on cluster character and electron distribution between borane and hydrocarbon ligands; significance of the structures of [η -9,10-(SMe)-8,8-(PPh₃)₂-nido-8,7-IrSB₉H₉] and [η -9,10-(SMe)-8-(η -C₅Me₅H)-nido-8,7-RhSB₉H₉]. *Journal of the Chemical Society Dalton Transactions*, **1997**, 149-152 15
- 208 The use of Kitaigorodskii's Aufbau principle in the solid-state study of crystalline borane compounds. A preliminary account. *Journal of Organometallic Chemistry*, **2002**, 657, 20-39 2.3 15
- 207 Monocarbaborane anion chemistry. The substantiation of the C-arylated [PhCB₆H₆]⁻ seven-vertex closo cluster anion by single-crystal synchrotron X-ray diffraction analysis. *Dalton Transactions RSC*, **2002**, 2807-2808 15

- 206 Macropolyhedral boron-containing cluster chemistry. The isolation and characterisation of the eighteen-vertex azaborane anti-[9-NB17H20]. *Journal of the Chemical Society Chemical Communications*, **1994**, 677-678 15
- 205 11-Vertex polyhedral rhodathiocarborane chemistry. Part 21. Synthesis and characterization of nido-structured [8,8-(PPh₃)₂-8-H-8,7,9-RhCSB8H10] and closo-structured [1,1-(PPh₃)₂-1,2,3-RhCSB8H9]. *Inorganic Chemistry*, **1992**, 31, 3339-3341 5.1 15
- 204 Steric and metal-centre control of two-, three- and four-orbital carborane-metal interaction in tin and platinum complexes of the {C₂B₈H₉X} ligand, where X = H and Ph. *Journal of Organometallic Chemistry*, **1992**, 437, C1-C6 2.3 15
- 203 Homobimetallic B-frame species: [(PPh₃)₂Ru]-(μ-MeCO₂)₃-(μ-H)₂-[RuB₁₀H₇], a unique diruthenaborane in which the metal atoms are linked by one-, two-, three-, and four-membered bridges. *Journal of the Chemical Society Chemical Communications*, **1987**, 1717-1718 15
- 202 Heterobimetallic B-frame compounds: systematic synthesis and molecular structure of the seven-vertex μ-HOs,Pt-nido-osmaplatinaborane [(Ph₃P)₂(CO)(Os)(PhMe₂P)ClHPtB₅H₇]. *Journal of the Chemical Society Chemical Communications*, **1983**, 951-952 15
- 201 Identification and molecular structure of the eighteen-vertex macropolyhedral diplatinoctadecaborane [(PMe₂Ph)₂Pt₂B₁₆H₁₅(C₆H₄Me)(PMe₂Ph)]. *Journal of the Chemical Society Chemical Communications*, **1986**, 556 15
- 200 Isomers of icosaborane(26): some synthetic routes and preliminary characterisations in the bis(nido-decaboranyl) system. *Journal of the Chemical Society Dalton Transactions*, **1979**, 840 15
- 199 The characterisation of 1,1-diphenylstannacyclopentane and 1,1,6,6-tetraphenyl-1,6-distannacyclodecane. An X-ray diffraction study of the distannacyclodecane. *Journal of the Chemical Society Perkin Transactions II*, **1981**, 369 15
- 198 Proton and boron-11 nuclear spin relaxation and the molecular tumbling of nido-decaborane in perdeuterotoluene solution. An interesting transition in solute-solvent interaction behaviour. *Journal of the Chemical Society, Faraday Transactions 2*, **1982**, 78, 525-536 15
- 197 Thermal isomerizations of monothiolated carboranes (HS)C₂B₁₀H₁₁ and the solid-state investigation of 9-(HS)-1,2-C₂B₁₀H₁₁ and 9-(HS)-1,7-C₂B₁₀H₁₁. *Journal of Organometallic Chemistry*, **2015**, 798, 132-140 2.3 14
- 196 Decaborane thiols as building blocks for self-assembled monolayers on metal surfaces. *Inorganic Chemistry*, **2012**, 51, 1685-94 5.1 14
- 195 New iridathiaboranes with reversible isonido nido cluster flexibility. *Inorganic Chemistry*, **2010**, 49, 7353-7361 14
- 194 Macropolyhedral boron-containing cluster chemistry. Mixed and multiple cluster fusion in platinaborane chemistry and the structure of [(PMe₂Ph)₂PtB₁₆H₁₇PtB₁₀H₁₁(PMe₂Ph)] as determined by synchrotron X-ray diffraction analysis. *Journal of the Chemical Society Dalton Transactions*, **1998**, 2777-2778 14
- 193 Vibrational Spectrum and Electronic Structure of the [B₁₁H₁₁]²⁻Dianion. *European Journal of Inorganic Chemistry*, **2007**, 2007, 4911-4918 2.3 14
- 192 Macropolyhedral boron-containing cluster chemistry. The eighteen-vertex monocarbaborane [(Me₃CNH₂)CB₁₇H₁₈(CN)]. *Journal of the Chemical Society Chemical Communications*, **1995**, 2407-2408 14
- 191 Eleven- and twelve-vertex polyhedral metalladithiaborane chemistry. Novel compounds from the arachno-[S₂B₉H₁₀]⁻anion: [(PPh₃)₃H₂IrS₂B₉H₁₀], [(PPh₃)₂H₂IrS₂B₉H₉] and [(PPh₃)₂HRhS₂B₈H₈]. *Journal of the Chemical Society Chemical Communications*, **1994**, 2265-2266 14
- 190 Macropolyhedral boron-containing cluster chemistry. A reductive trimerisation of MeNC to give an imidazole-based carbene stabilized by coordination to boron in an eighteen-vertex cluster compound. *Journal of the Chemical Society Chemical Communications*, **1994**, 1999-2000 14
- 189 Stepwise reduction of MeNC to Me₂NH on a metallaborane substrate: an interesting reaction sequence and the molecular structure of [μ-6,9-(NMe₂)-10-(PMe₂Ph)-5-(μ-C₆Me₆)-arachno-5-RuB₉H₁₀]. *Journal of the Chemical Society Chemical Communications*, **1989**, 1762-1763 14

| | | | | |
|-----|---|-----|--|----|
| 188 | Polyhedral iridamonocarbaborane chemistry. Two ten-vertex arachno-6,9-iridacarbadeboranes and some related ten-vertex carbaborane chemistry. Comparative nuclear magnetic resonance studies and the molecular structure of [asym-9-(CO)-9,9-(PPh ₃) ₂ -9-H-arachno-9,6-IrCB ₈ H ₁₂]. <i>Journal of the Chemical Society Dalton Transactions</i> , 1990 , 2887-2894 | | | 14 |
| 187 | Metallaheteroborane chemistry. Part 9. Syntheses and spectroscopy of platinum and palladium phosphine complexes containing B-(As ₂ B ₉)-based cluster ligands. Crystal structures of [3,3-L ₂ -closo-3,1,2-PtAs ₂ B ₉ H ₉](L = PPh ₃ or PMe ₂ Ph) and [3-Cl-3,8-(PPh ₃) ₂ -closo-3,1,2-PdAs ₂ B ₉ H ₈]. <i>Journal of the Chemical Society Dalton Transactions</i> , 1991 , 2895-2898 | | | 14 |
| 186 | Synthesis, molecular structure, and nuclear magnetic resonance spectra of the anionic monocarbametallaborane [NEt ₄][(CO) ₃ (PPh ₃)(OH)-closo-MoCB ₁₀ H ₁₀][CH ₂ Cl ₂]. <i>Journal of the Chemical Society Dalton Transactions</i> , 1987 , 2385-2389 | | | 14 |
| 185 | Homo- and hetero-bimetallic B-frame compounds: some novel twelve-vertex Ru/Ru, Os/Os, Ru/Os, and Os/Ru metallaboranes. <i>Journal of the Chemical Society Chemical Communications</i> , 1984 , 1398 | | | 14 |
| 184 | Preparation and nuclear magnetic resonance properties of eleven-vertex closo-type osmaundecaboranes and the X-ray crystal structure of the ortho-cycloboronated compound [2,5-(OEt) ₂ -1-(PPh ₃)-1-(o-Ph ₂ PC ₆ H ₄)-closo-1-OsB ₁₀ H ₇ -3]. <i>Journal of the Chemical Society Dalton Transactions</i> , 1986 , 2277 | | | 14 |
| 183 | Some palladium(II) and platinum(II) derivatives of pentaborane(9). <i>Journal of the Chemical Society Dalton Transactions</i> , 1978 , 1146 | | | 14 |
| 182 | Polyhedral Iridamonocarbaborane Chemistry. The Isolation and Characterisation of the Nine-Vertex closo-Type Species [(CO)(PPh ₃) ₂ IrCB ₇ H ₈] and the Partially Open Structures of Formally closo Nine-Vertex Clusters. <i>Collection of Czechoslovak Chemical Communications</i> , 1992 , 57, 1439-1450 | | | 14 |
| 181 | Chemistry of 11-vertex rhodathiaboranes: reactions with monodentate phosphines. <i>Dalton Transactions</i> , 2011 , 40, 6555-64 | 4.3 | | 13 |
| 180 | Macropolyhedral boron-containing cluster chemistry. Isolation and characterisation of the 27-vertex contiguous triple-cluster species [(PMe ₂ Ph)PtB ₂₆ H ₂₆ (PMe ₂ Ph)]. <i>Inorganic Chemistry Communication</i> , 1998 , 1, 365-367 | 3.1 | | 13 |
| 179 | Macropolyhedral boron-containing cluster chemistry. Ligand-induced two-electron variations of intercluster bonding intimacy. Structures of nineteen-vertex [(eta ⁵ -C ₅ Me ₅)HfB ₁₈ H ₁₉ (PMe ₂ Ph)] and the related carbene complex [(eta ⁵ -C ₅ Me ₅)HfB ₁₈ H ₁₉ [C(NHMe) ₂]]. <i>Dalton Transactions</i> , 2004 , 1521-3 | 4.3 | | 13 |
| 178 | Macropolyhedral boron-containing cluster chemistry. Aspects of the S ₂ B ₁₆ H ₁₆ system. Preparation, structure, NMR spectroscopy and isomerism. <i>Journal of Organometallic Chemistry</i> , 2003 , 680, 312-322 | 2.3 | | 13 |
| 177 | Metallaheteroborane chemistry: Part 16. Contrasting metal to heteroborane bonding modes in isoelectronic {MC ₂ B ₉ } and {MAS ₂ B ₉ } clusters. Synthesis and characterisation of [9-{Fe(CO) ₂ (B-C ₅ H ₅)}-nido-7,8-C ₂ B ₉ H ₁₂], [7-{Fe(CO) ₂ (B-C ₅ H ₅)}-nido-7,8-As ₂ B ₉ H ₁₀] and [7-{Fe(CO) ₂ (B-C ₅ H ₅)}-nido-7,8-As ₂ B ₉ H ₁₀]. <i>Journal of the Chemical Society Dalton Transactions</i> , 2003 , 1557-1564 | 4.3 | | 13 |
| 176 | A rearrangement of the 10-boron nido/arachno decaboranyl cluster. <i>Inorganic Chemistry Communication</i> , 2001 , 4, 544-546 | 3.1 | | 13 |
| 175 | B-frame supported bimetallics. Composite cluster compounds and the structures of [2,7-(B-C ₅ Me ₅) ₂ -nido-2,7,8,6-Ir ₂ CSB ₆ H ₈] and its 9-chloro derivative. Synchrotron and conventional X-ray studies. <i>Journal of Organometallic Chemistry</i> , 2000 , 614-615, 57-60 | 2.3 | | 13 |
| 174 | Macropolyhedral boron-containing cluster chemistry. An X-ray diffraction and nuclear magnetic resonance study of the double cluster 11-vertex: 11-vertex tetracarbaborane C ₄ B ₁₈ H ₂₂ . <i>Journal of the Chemical Society Dalton Transactions</i> , 1996 , 3813-3818 | | | 13 |
| 173 | Oxidative and cluster Aufbau reactions of nido-6-metalladeboranes induced by Lewis bases such as SMe ₂ : high-yield formation of [1-(B-C ₆ Me ₆)-isocloso-1-RuB ₉ H ₉] and the molecular structure of the Rh-commo-(isonido-eleven-vertex)-(closo-ten-vertex)conjuncto species [Rh ₂ (CO) ₂ (PPh ₃) ₂ (B-C ₆ Me ₆)(C ₁₀ H ₁₂ B ₁₀ H ₁₂)]. <i>Journal of the Chemical Society Dalton Transactions</i> , 1991 , 185 | | | 13 |
| 172 | Polyhedral osmaborane chemistry: the platination of [4,4,4-(CO)(PPh ₃) ₂ -nido-4-OsB ₅ H ₉] and the isolation, nuclear magnetic resonance properties, and molecular structures of two seven-vertex nido-type dimetallaheptaboranes [(CO)(PPh ₃) ₂ HOs(PMe ₂ Ph)ClPtB ₅ H ₇] and [(CO)(PPh ₃)(PPh ₂ O)(PPh ₃)PtB ₅ H ₇ Ph]. <i>Journal of the Chemical Society Dalton Transactions</i> , 1991 , 185 | | | 13 |
| 171 | Quantitative thermal rearrangement of an eleven-vertex metalladecaborane to give the isomeric cluster nido-[2-(B-C ₆ Me ₆)-8,10-Me ₂ -2,8,10-OsC ₂ B ₈ H ₈] with an unexpected BCBCB open face. <i>Journal of the Chemical Society Chemical Communications</i> , 1987 , 1650-1651 | | | 13 |

- 170 Metallaheteroborane chemistry. Part 4. The synthesis of closo-[2,2-(PPh₃)₂-2-H-1,2-XMB10H₁₀](X = Se or Te, M = Rh or Ir) compounds, their characterisation by nuclear magnetic resonance techniques, and the crystal and molecular structure of the X = Te, M = Rh complex. *Journal of the Chemical Society Dalton Transactions*, **1988**, 2739 13
- 169 Synthesis and nuclear magnetic resonance study of the auranoborane and diauradecaborane arachno-type compounds, [4-(S₂CNEt₂)-arachno-4-AuB₈H₁₂] and [6,9-(S₂CNEt₂)₂-arachno-6,9-Au₂B₈H₁₀]; crystal and molecular structure of [6,9-(S₂CNEt₂)₂-arachno-6,9-Au₂B₈H₁₀]. *Journal of the Chemical Society Dalton Transactions*, **1991**, 1427 13
- 168 Polyhedral rhenaborane chemistry: crystal and molecular structures of the nido-6-rhenadecaborane cluster compounds [6,6,6,6-(PMe₂Ph)₃H-nido-6-ReB₉H₁₃] and [2-(PMe₂Ph)-6,6,6,6-(PMe₂Ph)₂ClH-nido-6-ReB₉H₁₂]; nuclear magnetic resonance parameters of these and other related nido-undecaborane cluster species. *Journal of the Chemical Society Dalton Transactions*, **1985**, 1119 13
- 167 Comparative ten-vertex metallaborane chemistry: some nido-6-metalladecaboranes of tungsten, rhenium, ruthenium, osmium, and iridium; including the crystal and molecular structures of [6,6,6,6,6-(PMe₂Ph)₃H₂-nido-6-WB₉H₁₃] and [6,6,6-(PMe₂Ph)₃-nido-6-OsB₉H₁₃]. *Journal of the Chemical Society Dalton Transactions*, **1994**, 521 13
- 166 The chemistry of isomeric icosaboranes, B₂₀H₂₆. Molecular structures and physical characterization of 2,2?-bi(nido-decaboranyl) and 2,6?-bi(nido-decaboranyl). *Journal of the Chemical Society Dalton Transactions*, **1980**, 790-796 13
- 165 Some ten-vertex nido-metallaboranes: oxidative insertion of iridium(I) and rhodium(I) into the arachno-nonaborate anion, [B₉H₁₄]⁻ and the crystal and molecular structure of 6-hydrido-6,6-bis(triphenylphosphine)-nido-6-iridadecaborane, [(HirIII₂B₉H₁₃)(PPh₃)₂]. *Journal of the Chemical Society Dalton Transactions*, **1992**, 749-758 13
- 164 Phosphorus-31 chemical shift anisotropies in organophosphorus compounds. *Journal of the Chemical Society Chemical Communications*, **1976**, 666 13
- 163 Polyhedral azadirhodaborane chemistry. Reaction of [RhCl₂(C₅Me₅)₂] with [EtH₂NB₈H₁₁NHET] to give contiguousten-vertex [1-Et-6,7-(C₅Me₅)₂-closo-6,7,1-Rh₂NB₇H₇]. *Journal of the Chemical Society Dalton Transactions*, **1997**, 707-708 12
- 162 Diphosphacarborane analogues of ferrocene: the synthesis of two isomeric twelve-vertex closo-[(eta⁵-C₅H₅)FeP₂CB₈H₉] complexes. *Dalton Transactions*, **2005**, 909-13 4.3 12
- 161 Macropolyhedral boron-containing cluster chemistry. Metallathiaboranes from S₂B₁₇H₁₇: isolation and characterisation of [(eta⁶-MeC₆H₄isoPr)RuS₂B₁₆H₁₆] and [(eta⁶-MeC₆H₄isoPr)RuS₂B₁₅H₁₅]. *Dalton Transactions*, **2006**, 3624-6 4.3 12
- 160 Azanonaborane-pyridine derivatives [(R²C₅H₄N)B₈H₁₁NHR¹]: synthesis, structure and some molecular-orbital calculations. *Journal of Organometallic Chemistry*, **2002**, 657, 205-216 2.3 12
- 159 Ten-vertex osmamonocarbaboranes via arachno and nido {CB₉} monocarbaboranes. Polyhedral contraction promoted by [OsCl₂(PPh₃)₃] in MeOH and the crystal and molecular structure of [1-H-1,1-(PPh₃)₂-2-Ph-3-(OMe)-isocloso-1,2-OsCB₈H₇]. *Inorganic Chemistry Communication*, **2003**, 6, 1454-1458 3.1 12
- 158 Metallaborane reaction chemistry. *Inorganic Chemistry Communication*, **1999**, 2, 315-318 3.1 12
- 157 Nine-Vertex Polyhedral Monothiaaborane Chemistry: Synthesis, Molecular Structures, and NMR Characterization of the New Thianonaboranes exo-6-L-arachno-4-SB₈H₁₀. *Inorganic Chemistry*, **1994**, 33, 4545-4552 5.1 12
- 156 Metallaheteroborane chemistry. Part 7. Synthesis, crystal structure, and characterisation of two dinuclear rhodatelluraboranes, [(PPh₃)₂RhTeB₁₀H₁₀]₂ and [(PPh₃)(CO)Rh₂Te₂B₂₀H₂₀]. *Journal of the Chemical Society Dalton Transactions*, **1990**, 1831 12
- 155 arachno[(C₆Me₆)RuB₈H₁₄]; A nine-vertex structural analogue of n-B₉H₁₅ rather than of iso-B₉H₁₅. *Journal of Organometallic Chemistry*, **1986**, 315, C1-C4 2.3 12
- 154 A novel eleven-vertex closo-type azametallaborane: [1-(C₆Me₆H₄-4-Pri)-closo-1,2-RuNB₉H₁₀]. *Journal of the Chemical Society Chemical Communications*, **1988**, 1240-1241 12
- 153 An n.m.r. study of the arachno-[B₁₀H₁₂PPh₂]⁻ anion: and interesting arachno-arachno topological transition induced by deprotonation. *Journal of the Chemical Society Chemical Communications*, **1983**, 575-576 12

- 152 The reactions of arachno-decaboranyl complexes L2B10H12 (L = two-electron donor ligand) with some platinum(II) compounds; nuclear magnetic resonance studies and the crystal and molecular structure of [8-Cl-7,7-(PMe2Ph)2-nido-7-PtB10H11]. *Journal of the Chemical Society Dalton Transactions*, **1984**, 2487 12
- 151 The signs of nuclear spin coupling constants in some trifluoromethyl derivatives of platinum and of gold. *Journal of the Chemical Society Dalton Transactions*, **1976**, 745 12
- 150 Metallaborane Reaction Chemistry. Part 4. Polyhedral Thiametallaborane Synthesis via Direct Incorporation of Sulfur Into Metallaboranes. *Collection of Czechoslovak Chemical Communications*, **1997**, 62, 1289-1298 11
- 149 Polyhedral azaborane chemistry. The establishment of members of the hpho-type family [(RH2N)B8H11NHR], where groups R are now other than ethyl. *Journal of the Chemical Society Dalton Transactions*, **1997**, 2547-2550 11
- 148 B-frame supported bimetallics. [(PMe2ph)2PtB9H11Ru(η-isoPrC6H4Me)] and [(PMe2ph)2PtB9H9Ru(η-isoPrC6H4Me)]; an interesting pair of electron-deficient nido and closo geometries. *Journal of Organometallic Chemistry*, **1998**, 550, 341-345 2.3 11
- 147 Polyhedral iridaborane chemistry: Elements of the 10-vertex closo/nido/closo continuum: Molecular structures of [(PPh3)2HirB9H9(PPh3)], [(PPh3)(Ph2PC6H4)IrB9H7(PPh3)], [(PPh3)(Ph2PC6H4)HirB9H6Cl(PPh3)], [(PPh3)(Ph2PC6H4)HirB9H6(PPh3)2] and [(PPh3)(Ph2PC6H4)HirB9H6(PPh3)2]. *Inorganica Chimica Acta*, **2001**, 256, 2703-2705 2.7 11
- 146 Twelve-vertex polyhedral carbaborane chemistry. Isostructural cations and anions: The globule/globule salt [H3NCH2C2B10H11][H3CCH2CB11H11]. *Polyhedron*, **2006**, 25, 1069-1075 2.7 11
- 145 Macropolyhedral boron-containing cluster chemistry. *Journal of Organometallic Chemistry*, **2002**, 657, 256-261 2.3 11
- 144 Nine-vertex polyhedral monocarbaplatinaborane chemistry. Isolation, molecular structure and nuclear magnetic resonance properties of [6,6-(PPh3)2-arachno-6,4-PtCB7H11]. *Journal of the Chemical Society Dalton Transactions*, **1993**, 1269-1274 11
- 143 An alternative route to cationic metallaheteroboranes. *Journal of the Chemical Society Dalton Transactions*, **1993**, 3147-3148 11
- 142 Ten-vertex polyhedral dicarbaborane chemistry: new positional isomers of cluster constituents in the ten-vertex arachnodicarbaborane family: the patent [arachno-5,10-C2B8H13] anion and the isomeric ligand derivatives exo-9-L-arachno-5,6-C2B8H12 and exo-6-L-arachno-5,10-C2B8H12. *Inorganica Chimica Acta*, **1994**, 221, 5-7 2.7 11
- 141 Ten-vertex polyhedral azametallaborane chemistry. An arachno-6,9-azaplatinadecaborane and a nido-6,8-azaruthenadecaborane. *Inorganica Chimica Acta*, **1994**, 227, 163-166 2.7 11
- 140 Polyhedral organoiridazaborane chemistry. The preparation of the eleven-vertex nido compound [9-Cl-8-(η-C5Me5)-nido-8,7-IrNHB9H10] and its characterization by single-crystal X-ray diffraction analysis and nuclear magnetic resonance spectroscopy. *Collection of Czechoslovak Chemical Communications*, **1991**, 56, 1607-1617 11
- 139 Arene/ruthenaborane clusters I. Products of the reaction between [(η-C6Me6)2Ru2H4]RuB10H8(OEt)2 and carbon monoxide. Nuclear magnetic resonance studies and the crystal structure of [(η-C6Me6)2Ru2H2(CO)2]RuB10H8(OEt)2. *Journal of Organometallic Chemistry*, **1993**, 566, 273-287 2.3 11
- 138 Polyhedral iridathiaborane chemistry. Reactions of nido-6-SB9H11 and of [arachno-6-SB9H12] with [(Ir(η-C5Me5)Cl2)2] to give nido-8,7-iridathiaundecaboranes. A nuclear magnetic resonance and structural study. *Journal of the Chemical Society Dalton Transactions*, **1991**, 2657-2667 11
- 137 A double cluster of ruthenium and boron; [1-η-((η-C6Me6)2Ru2H4)-isocloso-1-RuB10H8-2,3-(OEt)2]. *Journal of the Chemical Society Chemical Communications*, **1987**, 442-443 11
- 136 Polyhedral ruthenaborane chemistry: characterization of several new ruthenaboranes by nuclear magnetic resonance spectroscopy, and the crystal and molecular structure of [5,6,6-(PPh3)3-6-H-nido-6-RuB9H12]. *Journal of the Chemical Society Dalton Transactions*, **1985**, 2397 11
- 135 Isomers of B20H26: structural characterisation by X-ray diffraction of 2,2'-Bi(nido-decaboranyl). *Journal of the Chemical Society Chemical Communications*, **1979**, 17-18 11

- 134 Polyhedral Azaborane Chemistry: NMR Parameters for the Unique Eight-Boron Polyhedral Species [(C₂H₅NH₂)B₈H₁₁NHC₂H₅]. *Collection of Czechoslovak Chemical Communications*, **1996**, 61, 1773-1782 11
- 133 Macropolyhedral Boron-Containing Cluster Chemistry. A Metallathiaborane from S₂B₁₇H₁₇: Isolation and Characterisation of [(PMe₂Ph)₂PtS₂B₁₆H₁₆]; A neo-arachno Ten-Vertex Cluster Shape, and the Constitution of the [arachno-B₁₀H₁₅]- Anion. *Collection of Czechoslovak Chemical Communications*, **2005**, 70, 430-440 11
- 132 Macropolyhedral metallaboranes [Aspects of preparation, constitution and structure. *Coordination Chemistry Reviews*, **2016**, 323, 71-86 23.2 10
- 131 Polyhedral nonaborane chemistry: an unsuspected new isomeric type in the nine-vertex arachno-B₉H₁₃L system. *Journal of Organometallic Chemistry*, **1998**, 550, 441-444 2.3 10
- 130 Macropolyhedral boron-containing cluster chemistry. Synchrotron X-ray structural analysis of [(PMe₂Ph)₂Pd₂B₁₆H₂₀(PMe₂Ph)₂] and [(PMe₂Ph)₃Pt₂B₁₆H₁₈(PMe₂Ph)]: models of intermediates to more condensed metallaboranes from the [(PMe₂Ph)₂PtB₈H₁₂] thermolysis system. *Chemical Communications*, **2005**, 1584-6 5.8 10
- 129 Azacarbaborane chemistry. Butyl nitrite synthesis of the new eight-, nine- and ten-vertex azacarbaboranes exo- and endo-7-CH₃-hypho-7,8-NCB₆H₁₁, nido-6,8,9-NC₂B₇H₁₀, arachno-6,5,9-NC₂B₇H₁₂ and arachno-6,5,10-C₂NB₇H₁₂. *Dalton Transactions*, **2003**, 1326 4.3 10
- 128 Synthesis of N-substituted derivatives of the hypho-(amine)(amino)B₈H₁₁ system. *Journal of Organometallic Chemistry*, **2000**, 614-615, 215-222 2.3 10
- 127 Nine- and Eight-Vertex Polyhedral Dicarborane Chemistry: New arachno and hypho Dicarboranes from arachno-4,5-C₂B₇H₁₃: Isolation and Characterization of the [arachno-4,5-C₂B₇H₁₂]- and [hypho-7,8-C₂B₆H₁₃]- Anions, and of the Neutral Ligand Derivatives [(4,5-C₂B₇H₁₂)(L)B₂], [(7,8-C₂B₆H₁₃)(L)B], [(4,5-C₂B₇H₁₂)(L)B₂], [(7,8-C₂B₆H₁₃)(L)B]. *Journal of Organometallic Chemistry*, **2005**, 694, 153-160 10
- 126 Polyhedral metallaphosphaborane chemistry: the preparation, molecular structure, and nuclear magnetic resonance study of [2-Ph-1,1-(PMe₂Ph)₂-closo-1,2-PtPB₁₀H₁₀], and some related chemistry. *Journal of the Chemical Society Dalton Transactions*, **1990**, 73 10
- 125 The preparation of [Ru(NCMe)₆][7-(η -C₆Me₆)-nido-7-RuB₁₀H₁₃]₂ and its characterization by single-crystal X-ray diffraction analysis and nuclear magnetic resonance spectroscopy: a triply hydrogen-bridged eleven-vertex nido-7-metallaundecaborane. *Journal of the Chemical Society Dalton Transactions*, **1997**, 1163-1171 10
- 124 Pentamethylcyclopentadienylrhodaborane chemistry. Part 5. Preparation, molecular structure, and nuclear magnetic resonance spectra of the arachno ten-vertex rhodadecaboranes [6-(η -C₅Me₅)-6,9-(RNC)₂-arachno-6-RhB₉H₁₁](R = Me or p-MeC₆H₄). *Journal of the Chemical Society Dalton Transactions*, **1988**, 2809-2815 10
- 123 Isomers of B₂₀H₂₆: elucidation of the structure of 6,6?-Bi(nido-decaboranyl) by ¹¹B-{1H} and ¹H-{¹¹B} n.m.r. spectroscopy. *Journal of the Chemical Society Chemical Communications*, **1979**, 16-17 10
- 122 Dynamic n.m.r. behaviour of {6,7,8,9- η -nido-decaborato(2)}bis(dimethylphenylphosphine)platinum(II), [Pt(η -B₁₀H₁₂)(PMe₂Ph)₂]; a novel type of metalloborane fluxionality. *Journal of the Chemical Society Chemical Communications*, **1980**, 305-306 10
- 121 N.M.R. studies of dimethyl cadmium partially oriented in the nematic phase of EBBA. *Molecular Physics*, **1977**, 34, 215-222 1.7 10
- 120 An assessment of the intercarbon stretching phenomenon in C-substituted β pseudocloso \square {3,1,2-RuC₂B₉} metalladicarbaboranes. *Journal of Organometallic Chemistry*, **2014**, 749, 163-173 2.3 9
- 119 Nine-vertex metallaborane chemistry. Preparation and characterisation of [1,1,1-(PMe₃)₂H-isocloso-IrB₈H₇-8-X], where X = H or Cl. *Journal of Organometallic Chemistry*, **2012**, 721-722, 155-163 2.3 9
- 118 Macropolyhedral boron-containing cluster chemistry. Characterisation of rigid 77-atom [Pt(B₁₈H₂₀)₂]₂-dianion isomers and an unusual metallaborane homophilic interaction. *Journal of the Chemical Society Dalton Transactions*, **1997**, 2543-2546 9
- 117 Macropolyhedral boron-containing cluster chemistry: The reaction of syn-B₁₈H₂₂ with SMe₂ and I₂ in monoglyme: Structure of [7-(SMe₂)-syn-B₁₈H₂₀]. *Inorganic Chemistry Communication*, **2007**, 10, 125-128¹ 9

| | | | |
|-----|--|-----|---|
| 116 | Macropolyhedral boron-containing cluster chemistry. The reaction of B ₁₆ H ₂₀ and B ₁₄ H ₁₈ with [PtMe ₂ (PMe ₂ Ph) ₂] to give [(PMe ₂ Ph) ₂ PtB ₁₆ H ₁₇ Me] and [(PMe ₂ Ph) ₂ PtB ₁₄ H ₁₆]. <i>Dalton Transactions</i> , 2005 , 1499-503 | 4.3 | 9 |
| 115 | Macropolyhedral boron-containing cluster chemistry: two-electron variations in intercluster bonding intimacy. Contrasting structures of 19-vertex [(B-C ₅ Me ₅)HrB ₁₈ H ₁₉ (PPh ₂)] and [(B-C ₅ Me ₅)IrB ₁₈ H ₁₈ (PPh ₂)]. <i>Inorganica Chimica Acta</i> , 2004 , 357, 3119-3123 | 2.7 | 9 |
| 114 | Macropolyhedral boron-containing cluster chemistry. Cluster assembly about a molybdenum centre. Formation of the 19-vertex [(CO) ₂ MoB ₁₆ H ₁₅ C ₂ Ph ₂]- anion. <i>Dalton Transactions</i> , 2004 , 2612-3 | 4.3 | 9 |
| 113 | Macropolyhedral boron-containing cluster chemistry. An unusual neonido ten-vertex subcluster configuration in a [(PPh ₃) ₂ RuB ₁₆ H ₂₀] species. <i>Journal of Organometallic Chemistry</i> , 2005 , 690, 2857-2859 ^{2,3} | | 9 |
| 112 | Polyhedral ruthenaborane chemistry. Approaches to encapsulated boron cores. The isolation and characterisation of the partially encapsulated isocloso 10-vertex {RuB ₉ } cluster compound [(PPh ₃)RuB ₉ H ₉ {RuCl ₂ (PPh ₃) ₂ } ₂]. <i>Inorganic Chemistry Communication</i> , 2005 , 8, 147-150 | 3.1 | 9 |
| 111 | An air-stable, cationic metallacarborane without a charge-compensated carborane ligand. <i>Chemical Communications</i> , 1996 , 679-681 | 5.8 | 9 |
| 110 | The first cluster azathiaboranes: isolation and characterisation of eight-vertex hypho-7,8-NSB ₆ H ₁₁ and ten-vertex arachno-6,9-NSB ₈ H ₁₁ . <i>Journal of the Chemical Society Chemical Communications</i> , 1993 , 1628-1629 | | 9 |
| 109 | Eleven-vertex polyhedral stannadecaborane chemistry. Naked tin[closo-1,2,3-Sn ₂ B ₈]-cluster compounds. <i>Journal of the Chemical Society Dalton Transactions</i> , 1993 , 1661-1663 | | 9 |
| 108 | Ten-Vertex Polyhedral Rhodadecaborane Chemistry. Molecular Structure, NMR Properties, and Cluster Dynamic Behaviour of [2,2,2-(PPh ₃) ₂ H-closo-2,1,6-RhC ₂ B ₇ H ₉]. <i>Collection of Czechoslovak Chemical Communications</i> , 1993 , 58, 1555-1568 | | 9 |
| 107 | Novel phosphonium-betaine ligands [SC(H)PPh ₃ -C,S] and [S(H)CC(PPh ₃)S-S,S] stabilised in rhodacarborane complexes. <i>Journal of the Chemical Society Chemical Communications</i> , 1994 , 1595-1596 | | 9 |
| 106 | Direct evidence for N-B donation into a polyhedral borane cluster. <i>Journal of the Chemical Society Chemical Communications</i> , 1989 , 1115-1116 | | 9 |
| 105 | Metallaheteroborane chemistry. Part 6. Synthesis of closo-[2-(ligand)-1,2-TeMB ₁₀ H ₁₀] complexes with M(ligand)= Rh(B-C ₅ Me ₅)(1), Ru(B-p-MeC ₆ H ₄ Pri)(2), Ru(B-C ₆ Me ₆)(3), and of nido-[6-(B-C ₆ Me ₆)-8-(OEt)-6-RuB ₉ H ₁₂](4), their characterisation by nuclear magnetic resonance spectroscopy. <i>Journal of the Chemical Society Dalton Transactions</i> , 1991 , 1211-1214 | | 9 |
| 104 | Determination of the relative signs of 2J(31P _B 1P) in complexes of tungsten(0) and molybdenum(0) using two-dimensional [31P _B 1P]-COSY-45 nuclear magnetic resonance chemical shift correlation. <i>Journal of the Chemical Society Dalton Transactions</i> , 1987 , 2401-2405 | | 9 |
| 103 | Homobimetallic B-frame species: in interesting (2 → 1) → (10 → 2) cluster disproportionation of an Eleven-vertex nido-Platinaundecaborane and the molecular structure of the Twelve-vertex closo-diplatinadodecaborane [1-Cl-1,2,2,4-(PMe ₂ Ph) ₄ -closo-1,2-Pt ₂ B ₁₀ H ₉]. <i>Inorganica Chimica Acta</i> , 1997 , 258, 123-144 | 2.7 | 9 |
| 102 | Correlation of thallium boron nuclear spin states by two-dimensional n.m.r. spectroscopy: magnitudes and relative signs of couplings nJ(205Tl → 1B) in the nido-thallaundecaborane anion [Me ₂ TlB ₁₀ H ₁₂]. <i>Journal of the Chemical Society Chemical Communications</i> , 1985 , 855-856 | | 9 |
| 101 | Synthesis, reactions, and nuclear magnetic resonance studies of some substituted arachno-decaborane and arachno-nonaborane derivatives, and the isolation of novel polyhedral diplatinaboranes. Crystal and molecular structure of [Pt ₂ (PMe ₂ Ph) ₂ (B-B ₂ H ₅)(B-B ₆ H ₉)]. <i>Journal of the Chemical Society Dalton Transactions</i> , 1991 , 3123-3128 | | 9 |
| 100 | Polyhedral phosphaborane chemistry: crystal and molecular structure of the diphenylphosphido-bridged arachno-decaboranyl cluster compound [PMePh ₃][6,9-μ-(PPh ₂)B ₁₀ H ₁₂]. <i>Journal of the Chemical Society Dalton Transactions</i> , 1986 , 303-308 | | 9 |
| 99 | A novel diplatinadecaborane with a PtPt bond: [(PMe ₂ Ph) ₂ (Pt ₂ B ₈ H ₁₄)]. <i>Journal of the Chemical Society Chemical Communications</i> , 1982 , 1019-1020 | | 9 |

- 98 Concerning tin-119 NMR evidence for the polymeric structure of methyltin alkoxides $\text{Me}_4\text{-n Sn(OR)}_n$. *Journal of Molecular Structure*, **1976**, 31, 207-209 3.4 9
- 97 Organometallic reactions. Part XVII. Some exchange and addition reactions of aminodimethyltin compounds. *Journal of the Chemical Society C, Organic*, **1970**, 759 9
- 96 Nomenclature for boranes and related species (IUPAC Recommendations 2019). *Pure and Applied Chemistry*, **2020**, 92, 355-381 2.1 8
- 95 A DFT and crystallographic reinvestigation of the $[\text{L}_2\text{RuC}_2\text{B}_7\text{H}_9]$ and $[\text{L}_3\text{RuC}_2\text{B}_7\text{H}_9]$ hypercloso and closo systems. *Polyhedron*, **2011**, 30, 2140-2145 2.7 8
- 94 Macropolyhedral boron-containing cluster chemistry. The unique nido-five-vertex--nido-ten-vertex conjuncto structure of $[(\text{eta}^5\text{-C}_5\text{Me}_5)_2\text{Rh}_2\text{B}_{11}\text{H}_{15}]$ via an unexpected cluster-dismantling. *Chemical Communications*, **2007**, 3559-61 5.8 8
- 93 Polyhedral Dipalladaborane Chemistry. The Molecular Structure and Cluster Electron Count of $[\text{7,8-(PPh}_3)_2\text{-7,8-(PPh}_2)_9,11\text{-(OEt)}_2\text{-nido-7,8-Pd}_2\text{B}_9\text{H}_8]$. *Collection of Czechoslovak Chemical Communications*, **2007**, 72, 1631-1638 8
- 92 Macropolyhedral boron-containing cluster chemistry. *Inorganica Chimica Acta*, **2005**, 358, 1709-1714 2.7 8
- 91 Metallaborane reaction chemistry. Part 10. Phenylacetylene incorporation via $[\text{4,4-(PMe}_2\text{Ph)}_2\text{-arachno-4-PtCB}_8\text{H}_{12}]$ in a β -reverse-metalladiborane synthesis of $[\text{7,7-(PMe}_2\text{Ph)}_2\text{-isonido-7,6,8-PtC}_2\text{B}_6\text{H}_7\text{-6-Ph}]$. *Inorganic Chemistry Communication*, **2005**, 8, 143-146 3.1 8
- 90 Polyhedral platinaborane chemistry. Some structural and NMR studies on the four-vertex arachno-1-platinatetraborane system. *Inorganica Chimica Acta*, **1992**, 196, 201-208 2.7 8
- 89 Ten-vertex metallaborane clusters: action as a B-frame support for heterobimetallic species: $[\text{7,7-(PMe}_3)_2\text{-9,9,9-H(PPh}_3)(\text{Ph}_2\text{P-ortho-})]$ and related compounds. *Inorganica Chimica Acta*, **1993**, 203, 193-200 2.7 8
- 88 Novel bonding mode and reaction of a phosphonium-betaine ligand $\text{S}_2\text{C(H)(PPh}_3)_2$ dimers (X = Se, Te) and X-ray diffraction study of the te-compound. *Journal of the Chemical Society Chemical Communications*, **1989**, 906 8
- 87 Organoruthenaborane chemistry. Part 6. The reaction of $[\{\text{Ru}(\text{C}_6\text{Me}_6)\text{Cl}_2\}_2]$ with the closo- $[\text{B}_{10}\text{H}_{10}]^{2-}$ anion: characterisation of some closo-type eleven-vertex 1-ruthenaundecaboranes and the molecular structure of $[\text{5-(C}_6\text{Me}_6\text{-6-(OMe)-nido-5-RuB}_9\text{H}_{12}]$. *Journal of the Chemical Society Dalton Transactions*, **1990**, 225-230 8
- 86 The signs of nuclear spin-spin coupling constants in some organoplatinum compounds. *Journal of Organometallic Chemistry*, **1979**, 172, 479-489 2.3 8
- 85 Ten-Vertex Polyhedral Monocarbaborane Chemistry. The Novel High-Yield Formation of the Ten-Vertex closo Compound $[\text{6-(PMe}_2\text{Ph)-closo-1-CB}_9\text{H}_9]$ from the Thermolysis of $[\text{8,8-(PMe}_2\text{Ph)}_2\text{-nido-8,7-PtCB}_9\text{H}_{11}]$. *Collection of Czechoslovak Chemical Communications*, **1993**, 58, 2924-2935 8
- 84 Nine-Vertex Polyhedral Monoazaborane Chemistry: Synthesis and NMR Characterization of exo-6-Ligand-arachno-4-azanoboranes(11), 6-L-4-NB $_8\text{H}_{11}$. *Collection of Czechoslovak Chemical Communications*, **1994**, 59, 2244-2252 8
- 83 Polymethylated $[\text{Fe}(\text{C}_6\text{-arene})_2]^{2+}$ dications: methyl-group rearrangements and application of the EINS mechanism. *Dalton Transactions*, **2011**, 40, 5916-20 4.3 7
- 82 The β -globule- β -globule-hybrid dicarbaborane- β -polyoxometallate salt, $[\text{C}_2\text{B}_{10}\text{H}_{11}\text{CH}_2\text{NHCH}(\text{CH}_3)_2]_4[\text{W}_{10}\text{O}_{32}][\text{H}_2\text{O}]_2[(\text{CH}_3)_2\text{CO}]_4$. *CrystEngComm*, **2003**, 5, 93-95 3.3 7
- 81 Polyhedral dicarbaborane chemistry: a convenient synthesis of the $[\text{nido-7,8-C}_2\text{B}_9\text{H}_{12}]^{2-}$ dicarbollide anion via the Brellocks reaction. *Inorganic Chemistry Communication*, **2005**, 8, 52-54 3.1 7

- 80 Polyhedral palladaborane chemistry: isolation and structural characterization of ten-vertex [(PMe₂Ph)₂PdB₉H₁₂(PMe₂Ph)] and eleven-vertex [(PMe₂Ph)₂PdB₁₀H₁₂]. *Journal of Chemical Crystallography*, **2000**, 30, 283-289 0.5 7
- 79 Polyhedral Azaplatinaborane Chemistry. Reaction of Members of the hypso-Type Family R'²H₂NB₈H₁₁NHR with [PtCl₂(PMe₂Ph)₂] and [PtCl₂(PPh₃)₂]. Formation of Members of the Family of Open Seven-Boron Species [3,3-(PMe₂Ph)₂-3-PtB₇H₁₀-B₆-(NHR)]. *Collection of Czechoslovak Chemical Communications*, **1999**, 64, 938-946 7
- 78 The relationship between the molecular structure of [3,3-(PMe₂Ph)₂-closo-3,1,2-PtC₂B₉H₁₁] and the mechanism of the fluxional behaviour of M(PR₃)₂-units in closo-twelve-atom metallaheteroboranes. *Journal of Organometallic Chemistry*, **1995**, 503, C12-C15 2.3 7
- 77 Ten-vertex polyhedral monoheteroborane chemistry: syntheses, molecular structures, and nuclear magnetic resonance characterization of some 9-substituted 6-aza- and 6-thia-arachno-decaboranes. *Journal of the Chemical Society Dalton Transactions*, **1996**, 1741-1751 7
- 76 Synthesis and crystal structure of [Pd(PPh₃)₂S₂B₆H₈]₂: a dimeric metallaheteroborane with heteroatoms acting both as cluster atoms and two-electron donor sites. *Journal of the Chemical Society Chemical Communications*, **1994**, 2403 7
- 75 Polyhedral organoiridaborane chemistry. The reaction of MeNC with [6-(B-C₅Me₅)-nido-6-IrB₉H₁₃]. Nuclear magnetic resonance studies and the crystal and molecular structure of [4,4-(B-C₅Me₅)(MeNC)-arachno-4-IrB₈H₁₂]. *Journal of the Chemical Society Dalton Transactions*, **1992**, 1415-1419 7
- 74 A fluxional arachno-1 -metallapentaborane, [1,1,1 -(CO)(PMe₃)₂(1-IrB₄H₉)]. *Journal of the Chemical Society Dalton Transactions*, **1982**, 481 7
- 73 Formation and X-ray crystal structure of 1,1,6,6-tetraphenyl-1,6-distannacyclodecane. *Journal of the Chemical Society Chemical Communications*, **1978**, 791 7
- 72 Through-space spin coupling between phosphorus and hydrogen in .pi.-C₅H₅W(CO)₂PR₃X (R = methyl, phenyl, methoxy, etc. X = iodine, trimethyltin). *Inorganic Chemistry*, **1973**, 12, 2742-2742 5.1 7
- 71 Metallaborane Reaction Chemistry. Part 3. Reaction of Carbon Monoxide with [6-H-6-(PPh₃)₆P,5C-[2-Ph₂PC₆H₄]-nido-6-IrB₉H₁₂] and the Isolation and Characterisation of Two arachno-6-Monoiridadecaboranes [6-(CO)-6-H-6-(PPh₃)₆P,5C-[2-Ph₂PC₆H₄]-nido-6-IrB₉H₁₂] and [6-(CO)-6-H-6-(PPh₃)₆P,5C-[2-Ph₂PC₆H₄]-nido-6-IrB₉H₁₂]. *Journal of the Chemical Society Dalton Transactions*, **1999**, 1415-1419 7
- 70 Metallaborane Reaction Chemistry. Part 7. B-Frame Supported Bimetallics: Ligand-to-Metal Organometallic Interaction in Dimetallaboranes and an Interesting Ligand Displacement Cascade. *Collection of Czechoslovak Chemical Communications*, **1999**, 64, 938-946 7
- 69 Polyhedral Platinaborane Chemistry. Interaction of PMe₂Ph with [(PMe₂Ph)₂PtB₁₀H₁₂]. *Organometallics*, **2012**, 31, 2691-2696 3.8 6
- 68 Borane Polyhedra as Building Blocks for Unknown but Potentially Isolatable New Molecules □ Extensions based on Computations of the Known B₁₈H₂₂ Isomers. *Croatica Chemica Acta*, **2013**, 86, 485-494 0.8 6
- 67 Polyhedral metallathaborane chemistry: Synthesis and characterisation of metallathaboranes based on the twelve-vertex icosahedral closo-{MSB₁₀H₁₀} unit, where M is Rh or Ir. *Journal of Organometallic Chemistry*, **2008**, 693, 435-445 2.3 6
- 66 Macropolyhedral boron-containing cluster chemistry. *Journal of Organometallic Chemistry*, **2002**, 657, 262-266 2.3 6
- 65 Polyhedral monocarbaborane chemistry. *Journal of Organometallic Chemistry*, **2005**, 690, 2815-2828 2.3 6
- 64 Macropolyhedral borane reaction chemistry: Reductive oligomerisation of terBuNC by anti-B₁₈H₂₂ to give the boron-coordinated {(terBuNHCH){terBuNHC(CN)}CH₂:} carbene residue. *Inorganic Chemistry Communication*, **2005**, 8, 491-494 3.1 6
- 63 Triple linking of the decaboranyl cluster. Structure of [(SMe₂)₂B₁₀H₁₀(B₁₀H₁₃)₂] as determined by synchrotron X-ray diffraction analysis. *Chemical Communications*, **2001**, 1788-9 5.8 6

- 62 Ten-vertex polyhedral mixed diheteromonometallaborane chemistry. Directed synthesis of arachno-9,6,8-platinathiacarbadecaboranes. *Inorganica Chimica Acta*, **1994**, 218, 1-3 2.7 6
- 61 Organoruthenaborane Chemistry. X. The SM2-catalysed formation of [l-(η -C6Me6)-isocloso-RuB9H9], and some B-phenylamino derivatives. NMR and X-ray diffraction studies. *Zeitschrift Fur Anorganische Und Allgemeine Chemie*, **1992**, 616, 79-85 1.3 6
- 60 Polyhedral metallacarbathiaborane chemistry. Designed synthetic routes to nine-vertex mixed diheteromonometallaborane cluster compounds. *Inorganica Chimica Acta*, **1993**, 214, 17-19 2.7 6
- 59 Preparation and structure of [7-(η -C5Me5)-8-Cl-11-(PMe2Ph)-nido-7-RhB10H11], a metallaborane analogue of the fluxional borane anion nido-B11H14. *Journal of the Chemical Society Chemical Communications*, **1985**, 1165-1166 6
- 58 An NMR investigation of the [B10H12]2 π ligand in [Pt(B10H12)(PMe2Ph)2]. *Journal of Magnetic Resonance*, **1980**, 38, 529-535 6
- 57 Bis-(η -2,3,4- η -nido-hexaboranyl)-bis(dimethylphenylphosphine)diplatinum(PtPt), [Pt2(η -B6H9)2(PMe2Ph)2], an unexpected product from the reaction of (B10H13)2O with [PtCl2(PMe2Ph)2]; crystal and molecular structure. *Journal of the Chemical Society Chemical Communications*, **1980**, 375-38 6
- 56 Mass spectroscopic evidence for icosaborane(26). *The Journal of Physical Chemistry*, **1978**, 82, 623-625 6
- 55 Organometallic reactions. Part XIX. Some reactions of aldehydes with aminotin compounds. *Journal of the Chemical Society C, Organic*, **1971**, 68 6
- 54 Polyhedral platinadicarbaborane chemistry. The molecular structure and some NMR properties of [9,9-(PPH3)2-arachno-9,5,6-PtC2B7H11]. *Collection of Czechoslovak Chemical Communications*, **1991**, 56, 646-656 6
- 53 Nine-Vertex Polyhedral Monothiaaborane Chemistry. The First nido Thianonaborane: Isolation and Characterization by NMR of the [nido-9-SB8H9]- Anion, and its Conversion to Compounds of the 6-L-arachno-4-SB8H10 Series (where L = SME2 and Pyridine). *Collection of Czechoslovak Chemical Communications*, **1994**, 59, 367-373 6
- 52 Macropolyhedral Nickelaboranes from the Metal-Assisted Fusion of KBH. *Inorganic Chemistry*, **2019**, 58, 13258-13267 5.1 5
- 51 Isonitrile ligand effects on small-molecule-sequestering in bimetalladodecaborane clusters. *Journal of Organometallic Chemistry*, **2013**, 747, 76-84 2.3 5
- 50 Polyhedral azaborane chemistry. Nitrogen-vertex incorporation in metallaazaborane formation from 4-(NHET2)B9H13. Two isomers of [(η -C5Me5)RhB9H12(NET2)]. *Journal of the Chemical Society Dalton Transactions*, **1998**, 2353-2358 5
- 49 Macropolyhedral boron-containing cluster chemistry. Cluster opening and B-frame rearrangement in the reaction of B(16)H(20) with [(η -C5Me5)IrCl(2)(η -C(5)Me(5))2]. Synchrotron X-ray structures of [(η -C(5)Me(5))2Ir(2)B(16)H(17)Cl] and [(η -C(5)Me(5))2Ir(2)B(16)H(15)Cl]. *Dalton Transactions*, **2001**, 5281-5284 4.3 5
- 48 Bond rotamers and calculated 11B-NMR chemical shifts in boron-containing cluster chemistry. Some effects in the {nido-7,8,10-PC2B8} system. *Journal of Organometallic Chemistry*, **2000**, 614-615, 61-65 2.3 5
- 47 Chemie makropolyedrischer Borcluster: Isolierung und Charakterisierung des ersten makropolyedrischen Thiaborans, des Anions [9,9?-S2B17H18] vom arachno-Typ. *Angewandte Chemie*, **1994**, 106, 1663-1665 3.6 5
- 46 Signs of interphosphorus nuclear spin coupling constants in octahedral transition-metal complexes containing chelated phosphine ligands. *Inorganic Chemistry*, **1988**, 27, 1111-1112 5.1 5
- 45 Pentamethylcyclopentadienylrhodaborane chemistry. Part 4. The isolation, molecular structure, and nuclear magnetic resonance study of an interesting phosphino-bridged arachno-type nine-vertex rhodaborane [2,5-(η -PPh2)-1-(PHPh2)-2-(η -C5Me5)-2-RhB8H9]. *Journal of the Chemical Society Dalton Transactions*, **1988**, 2059-2061 5

| | | | |
|----|---|-----|---|
| 44 | The chemistry of isomers of icosaborane(26): Crystal and molecular structure of 1,2-bi(nido-decaboranyl). <i>Polyhedron</i> , 1985 , 4, 1981-1984 | 2.7 | 5 |
| 43 | Mercury-199 chemical shift anisotropy in methyl mercuric bromide. <i>Journal of the Chemical Society Chemical Communications</i> , 1974 , 595 | | 5 |
| 42 | Preparation and structure of 1,4-dichloro-1,1,3,3-tetraphenyl-catenadi-(boraphosphane), BH ₂ Cl[PPH ₂]BH ₂ [PPH ₂]Cl. <i>Journal of the Chemical Society Dalton Transactions</i> , 1978 , 40-43 | | 5 |
| 41 | Macropolyhedral boron-containing cluster chemistry. Further progress beyond the icosahedron. July 1999. <i>Special Publication - Royal Society of Chemistry</i> , 2007 , 171-174 | 0.1 | 5 |
| 40 | exo,endo and exo,exo isomers of 6,9-(PMe ₂ Ph) ₂ -arachno-B ₁₀ H ₁₂ and its halogenated derivatives. Molecular structures of exo,endo- and exo,exo-6,9-(PMe ₂ Ph) ₂ -arachno-B ₁₀ H ₁₂ and exo-6,endo-9-(PMe ₂ Ph) ₂ -2-Br-arachno-B ₁₀ H ₁₁ . <i>Journal of the Chemical Society Dalton Transactions</i> , 1997 , 4739-4746 | | 4 |
| 39 | Ten-vertex polyhedral azametallaborane chemistry: a unique nido-6,9 to nido-6,8-cluster isomerization. <i>Dalton Transactions</i> , 2008 , 4776-83 | 4.3 | 4 |
| 38 | Polyhedral Monocarbaborane Chemistry. Functionality and Isomerism: Reactions of the [6-Ph-nido-6-CB ₉ H ₁₁]- Anion with Aminopyridines NC ₅ H ₄ NH ₂ to Yield Neutral arachno and closo Ten-Vertex Monocarbaborane Derivatives. <i>Collection of Czechoslovak Chemical Communications</i> , 2005 , 70, 1873-1890 | | 4 |
| 37 | Macropolyhedral boron-containing cluster chemistry. A synthetic approach via the auto-fusion of [6,9-(SMe ₂) ₂ -arachno-B ₁₀ H ₁₂]. <i>Dalton Transactions</i> , 2006 , 3752-65 | 4.3 | 4 |
| 36 | The nido-osmaboranes [2,2,2-(CO)(PPh ₃)(2)-nido-2-OsB(5)H(9)] and [6,6,6-(CO)(PPh ₃)(2)-nido-6-OsB(9)H(13)]. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2001 , 57, 1245-7 | | 4 |
| 35 | Two potential pyridine-borane oligomer and polymer building blocks. Structural characterisation of [NC ₅ H ₄ .C ₅ H ₄ N.B ₁₀ H ₁₂ .NC ₅ H ₄ .C ₅ H ₄ N] and [Me ₂ S.B ₁₀ H ₁₂ .NC ₄ H ₄ N.B ₁₀ H ₁₂ .SMe ₂] by conventional and synchrotron X-ray methods. <i>Inorganic Chemistry Communication</i> , 1999 , 2, 298-300 | 3.1 | 4 |
| 34 | An interesting arachno eight-vertex tungstaborane: isolation and characterisation of the fluxional anion [3,3,3,3-(CO)4-arachno-WB ₇ H ₁₂]. <i>Journal of the Chemical Society Chemical Communications</i> , 1987 , 476 | | 4 |
| 33 | Transition-Metal Derivatives of Nido-Boranes and Some Related Species 1982 , 43-118 | | 4 |
| 32 | Determination of the sign and magnitude of J(XX) in [AnX] M _p nuclear spin systems. <i>Journal of the Chemical Society Chemical Communications</i> , 1975 , 638 | | 4 |
| 31 | Tin-119 magnetic shielding in norbornenyltrimethyltins. <i>Journal of Organometallic Chemistry</i> , 1976 , 104, 311-315 | 2.3 | 4 |
| 30 | Macropolyhedral boron-containing cluster chemistry. The reversible disassembly and reassembly of the hexagonal pyramidal {B ₇ } feature in the [S ₂ B ₁₈ H ₁₉]- anion. <i>Dalton Transactions</i> , 2008 , 1625-34 | 4.3 | 3 |
| 29 | Macropolyhedral boron-containing cluster chemistry. Novel intercluster linkages from the reaction of [Pt(cod)Cl ₂] and [PtMe ₂ (PMe ₂ Ph) ₂] with 6,6'-(B ₁₀ H ₁₃) ₂ O. <i>Chemical Communications</i> , 2007 , 5084-6 | 5.8 | 3 |
| 28 | An interesting metallocarborane cage closure and dismantling reaction including the facile halogenation of a cluster carbon atom: Structure of [1,2-(B-C ₅ Me ₅) ₂ -Et,2-H-closo-1,2,3-Rh ₂ CB ₄ H ₄ -3-I]. <i>Journal of Organometallic Chemistry</i> , 2005 , 690, 4967-4970 | 2.3 | 3 |
| 27 | Polyhedral azaborane chemistry, 6-(C ₅ H ₅ N)-arachno-4-NB ₈ H ₁₁ . <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2000 , 56 (Pt 5), 600-1 | | 3 |

- 26 Ten-vertex metallaborane chemistry. Structure of 3-(OC₂H₅)-4,9-(PPh₃)₂-7,7-Cl(Ph₂P-ortho-C₆H₄)-isonido-7-IrB₉H₅-10-.CH₂Cl₂. *Acta Crystallographica Section C: Crystal Structure Communications*, **1990**, 46, 1010-1013 3
- 25 The geometry of triethynyl phosphine partially oriented in a nematic liquid crystal. *Journal of Molecular Structure*, **1976**, 30, 125-127 3-4 3
- 24 Organometallic reactions. Part XIII. The reaction of trialkyltin alkoxides and of bistralkyltin oxides with sulphur dioxide, sulphodi-imides, and sulphinylamines. *Journal of the Chemical Society C, Organic*, **1968**, 2630 3
- 23 Macropolyhedral Boron-Containing Cluster Chemistry. Intermolecular Coordination via Hydrogen-Metal Interaction. The Solvent-Free Synthesis and Dimeric Constitution of [Pd₂B₃₆H₄₀(PMe₂Ph)₄]. *Collection of Czechoslovak Chemical Communications*, **2007**, 72, 1639-1645 3
- 22 Macropolyhedral boron-containing cluster chemistry [S₂B₁₆H₁₇]□A new eighteen-vertex thiaborane anion. *Collection of Czechoslovak Chemical Communications*, **2010**, 75, 807-812 3
- 21 The contrarotational fluxionality of [3,3-(PMe₂Ph)₂-closo-3,1,2-PtC₂B₉H₁₁] and related species. *Dalton Transactions*, **2015**, 44, 9620-9 4-3 2
- 20 Monocarbaborane Anions with 10 or 12 Vertices. *Inorganic Syntheses*, **2014**, 174-185 2
- 19 Coordination and cluster compounds of ruthenium with the [hypho-1,2-S₂B₆H₉]-ligand. *Dalton Transactions*, **2005**, 1979-84 4-3 2
- 18 Estimating organic chain length through sound velocity measurements. *Ultrasonics*, **2005**, 43, 219-26 3-5 2
- 17 Ten-vertex rhodadithiaborane chemistry.. *Acta Crystallographica Section C: Crystal Structure Communications*, **2001**, 57, 520-2 2
- 16 Isomeric icosaboranes B₂₀H₂₆: the synchrotron structure of 1,1'-bis(nido-decaboranyl). *Acta Crystallographica Section C: Crystal Structure Communications*, **2001**, 57, 779-80 2
- 15 Two iridanonaborane compounds. *Acta Crystallographica Section C: Crystal Structure Communications*, **2001**, 57, 49-51 2
- 14 Das Hexacarbaboran arachno-C₆B₆H₁₂ und ein methyliertes Pentacarbaboran arachno-CH₃C₅B₇H₁₂: Bereiche mit beginnenden Kohlenwasserstoffeigenschaften in Boranclustern. *Angewandte Chemie*, **1999**, 111, 1922-1924 3,6 2
- 13 Organoruthenaborane Chemistry. IX. The Reactions of [Cp-C₆Me₆RuCl₂]₂ and [Cp-MeC₆H₄-4-iPrRuCl₂]₂ with the [nido-B₉H₁₂]⁻anion. *Zeitschrift Fur Anorganische Und Allgemeine Chemie*, **1991**, 598, 45-52 1,3 2
- 12 Organometallic reactions. Part XVIII. The reactions of N-sulphinyltoluene-p-sulphonamide and of N-(2,2,2-trichloroethylidene)toluene-p-sulphonamide with some tributyltin-oxygen and nitrogen compounds. *Journal of the Chemical Society C, Organic*, **1970**, 1570-1576 2
- 11 Big Borane Assemblies, Macropolyhedral Species and Related Chemistry. *Challenges and Advances in Computational Chemistry and Physics*, **2015**, 139-180 0.7 1
- 10 Synthesis and characterization of dicarborylmethylammonium polyoxometallates. *Collection of Czechoslovak Chemical Communications*, **2010**, 75, 1075-1096 1
- 9 9-(Triphenylphosphine-P)-arachno-6-thiadecaborane(11) Trichloromethane (1/1). *Acta Crystallographica Section C: Crystal Structure Communications*, **1998**, 54, 121-123 1

- 8 Macropolyhedral boron-containing cluster chemistry. Synthesis of the nineteen vertex monocarbaborane [9-(terBuNH₂)-(anti)-9-CB18H₂₀] by direct carbon-atom Aufbau. *Dalton Transactions*, **2007**, 4766-8 4.3 1
- 7 Intramolecular and Supramolecular Cluster Interactions. *ChemInform*, **2004**, 35, no 1
- 6 Twisted [(R(3)P)PdX] groups above dicarbaborane ligands: 4-dimethylsulfido-3-iodo-3-triphenylphosphine-closo-3-pallada-1,2-dicarbadoecaborane and 3-dimethylphenylphosphine-3-chloro-4-dimethylsulfido-closo-3-pallada-1,2-dicarbadoecaborane. *Acta Crystallographica Section C: Crystal Structure Communications*, **2005**, 61, m393-6 1
- 5 Polyhedral Ruthenaborane Chemistry Preparation, Molecular Structure, and N.M.R. Properties of the Mixed-Ligand Eleven-Vertex isocloso-Type Compound [1,1-(PMe₃) (PPh₃)-closo-1-RuB₁₀H₈-2,3-(OEt)₂]. *Zeitschrift Fur Anorganische Und Allgemeine Chemie*, **1987**, 553, 24-34 1.3 1
- 4 The small-scale synthesis of ¹³C-labelled methyltin iodides. *Journal of Labelled Compounds*, **1975**, 11, 285-286 1
- 3 Macropolyhedral Chalcogenaboranes: Insertion of Selenium into the Isomers of BH_n. *Inorganic Chemistry*, **2022**, 5.1 1
- 2 [1,1,2,2-]. *Acta Crystallographica Section C: Crystal Structure Communications*, **2000**, 56 (Pt 12), 1423-4
- 1 [Mu-6,9-Cl-8-(OMe)-6,9-eta⁵-C₅Me₅)₂-arachno-6,9,5-rh²SB7H₇]. *Acta Crystallographica Section C: Crystal Structure Communications*, **2001**, 57, 52-4