

# Ying Cheng

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/311789/publications.pdf>

Version: 2024-02-01

11  
papers

341  
citations

1163117  
8  
h-index

1281871  
11  
g-index

11  
all docs

11  
docs citations

11  
times ranked

486  
citing authors

#	ARTICLE	IF	CITATIONS
1	Empagliflozin Inhibits Hepatic Gluconeogenesis and Increases Glycogen Synthesis by AMPK/CREB/GSK3 $\beta$ Signalling Pathway. <i>Frontiers in Physiology</i> , 2022, 13, 817542.	2.8	6
2	The SGLT2 inhibitor empagliflozin negatively regulates IL-17/IL-23 axis-mediated inflammatory responses in T2DM with NAFLD via the AMPK/mTOR/autophagy pathway. <i>International Immunopharmacology</i> , 2021, 94, 107492.	3.8	50
3	Prostaglandin F $_{2\alpha}$ protects against pericyte apoptosis by inhibiting the PI3K/Akt/GSK3 $\beta$ / $\beta$ -catenin signaling pathway. <i>Annals of Translational Medicine</i> , 2021, 9, 1021-1021.	1.7	10
4	PACS-2 attenuates diabetic kidney disease via the enhancement of mitochondria-associated endoplasmic reticulum membrane formation. <i>Cell Death and Disease</i> , 2021, 12, 1107.	6.3	17
5	An SGLT2 inhibitor modulates SHH expression by activating AMPK to inhibit the migration and induce the apoptosis of cervical carcinoma cells. <i>Cancer Letters</i> , 2020, 495, 200-210.	7.2	30
6	GADD45B Promotes Glucose-Induced Renal Tubular Epithelial-Mesenchymal Transition and Apoptosis via the p38 MAPK and JNK Signaling Pathways. <i>Frontiers in Physiology</i> , 2020, 11, 1074.	2.8	9
7	Urinary miR-3137 and miR-4270 as potential biomarkers for diabetic kidney disease. <i>Journal of Clinical Laboratory Analysis</i> , 2020, 34, e23549.	2.1	3
8	Pancreatic kallikrein protects against diabetic retinopathy in KK Cg-Ay/J and high-fat diet/streptozotocin-induced mouse models of type 2 diabetes. <i>Diabetologia</i> , 2019, 62, 1074-1086.	6.3	54
9	Effects of SGLT2 inhibitors on fractures and bone mineral density in type 2 diabetes: An updated meta-analysis. <i>Diabetes/Metabolism Research and Reviews</i> , 2019, 35, e3170.	4.0	50
10	Triptolide Attenuates Renal Tubular Epithelial-mesenchymal Transition Via the MiR-188-5p-mediated PI3K/AKT Pathway in Diabetic Kidney Disease. <i>International Journal of Biological Sciences</i> , 2018, 14, 1545-1557.	6.4	84
11	Saxagliptin regulates M1/M2 macrophage polarization via CaMKK $\beta$ /AMPK pathway to attenuate NAFLD. <i>Biochemical and Biophysical Research Communications</i> , 2018, 503, 1618-1624.	2.1	28