Rebecca Haddad

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3127310/publications.pdf

Version: 2024-02-01

1478280 1372474 22 138 10 6 citations h-index g-index papers 25 25 25 141 all docs docs citations times ranked citing authors

#	Article	IF	Citations
1	Intermittent Self-catheterization in Older Adults: Predictors of Success for Technique Learning. International Neurourology Journal, 2018, 22, 65-71.	0.5	20
2	Predictive factors of adherence to urinary selfâ€catheterization in older adults. Neurourology and Urodynamics, 2019, 38, 770-778.	0.8	16
3	Nocturia and Nocturnal Polyuria in Neurological Patients: From Epidemiology to Treatment. A Systematic Review of the Literature. European Urology Focus, 2020, 6, 922-934.	1.6	16
4	Efficiency and satisfaction with telephone consultation of followâ€up patients in neuroâ€urology: Experience of the COVIDâ€19 pandemic. Neurourology and Urodynamics, 2021, 40, 929-937.	0.8	15
5	Impact of food and drinks on urine production: A systematic review. International Journal of Clinical Practice, 2020, 74, e13539.	0.8	13
6	Association between nocturia and frailty among elderly males in a veterans administration population. Aging Clinical and Experimental Research, 2020, 32, 1993-2000.	1.4	10
7	Nocturia in patients with cognitive dysfunction: a systematic review of the literature. BMC Geriatrics, 2020, 20, 230.	1.1	6
8	Comparison of clinical and paraclinical characteristics of patients with urge, mixed, and passive fecal incontinence: a systematic literature review. International Journal of Colorectal Disease, 2021, 36, 633-644.	1.0	6
9	Is There a Relationship Between Overactive Bladder and Sexual Dysfunction in Women with Multiple Sclerosis?. Journal of Sexual Medicine, 2022, 19, 729-737.	0.3	6
10	Development of a flowchart reflecting the current attitude and approach towards idiopathic overactive bladder treatment in Belgium: A Delphi study. Neurourology and Urodynamics, 2020, 39, 1781-1795.	0.8	4
11	Prevalence of comorbidities in multiple sclerosis patients with neurogenic bladder. Progres En Urologie, 2021, 31, 732-738.	0.3	4
12	Efficacy of posterior tibial nerve stimulation (PTNS) on overactive bladder in older adults. European Geriatric Medicine, 2018, 9, 249-253.	1.2	3
13	Fecal impaction is associated with postoperative urinary retention after hip fracture surgery. Annals of Physical and Rehabilitation Medicine, 2021, 64, 101464.	1.1	3
14	Diagnosing Nocturnal Polyuria from a Single Nocturnal Urine Sample. European Urology Focus, 2020, 6, 738-744.	1.6	2
15	Long-term functional outcomes of artificial urinary sphincter (AMS 800â,,¢) implantation in women aged over 75Âyears and suffering from stress urinary incontinence caused by intrinsic sphincter deficiency. World Journal of Urology, 2021, 39, 3897-3902.	1.2	2
16	Detrusor contractility in post-menopausal women: Impact of ageing, complaint and urodynamic diagnosis. Progres En Urologie, 2021, 31, 406-413.	0.3	2
17	French version of the short form of neurogenic bladder symptom score: Cross-cultural adaptation and validation. Canadian Urological Association Journal, 2022, 16, .	0.3	2
18	Prioritization of risk situations in neuro-urology: guidelines from Association Fran§aise d'Urologie (AFU), Association Francophone Internationale des Groupes d'Animation de la Paraplégie (A.F.I.G.A.P.), Groupe de Neuro-urologie de Langue Française (GENULF), Société Française de Médecine Physique et de Réadaptation (SOFMER) and Société Interdisciplinaire Francophone d'UroDynamique et de Pelvi-Périnéologie (SIFUD-PP). World Journal of Urology, 2021, , 1.	: 1.2	1

#	Article	IF	CITATIONS
19	Functional independence measure predicts the outcome of clean intermittent catheterization training in patients with multiple sclerosis. Annals of Physical and Rehabilitation Medicine, 2022, 65, 101539.	1.1	1
20	Fecal incontinence subtype assessment (FI-SA): A new tool to distinguish among subtypes of fecal incontinence in a neurogenic population. Clinics and Research in Hepatology and Gastroenterology, 2022, 46, 101900.	0.7	1
21	Persistent need to urinate: A common sensory symptom leading to urinary discomfort. A study of 79 cases., 2022, 2, 100007.		1
22	In multiple sclerosis, a Functional Independence Measure ≥ 107 is the best predictor of outcome after clean intermittent catheterization training. Annals of Physical and Rehabilitation Medicine, 2023, 66, 101636.	1.1	0