Fengqin Li

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3135767/publications.pdf

Version: 2024-02-01

394421 454955 1,038 41 19 30 citations h-index g-index papers 48 48 48 1284 all docs docs citations times ranked citing authors

#	Article	IF	CITATIONS
1	Prevalence and Characterization of Staphylococcus aureus Cultured From Raw Milk Taken From Dairy Cows With Mastitis in Beijing, China. Frontiers in Microbiology, 2018, 9, 1123.	3.5	88
2	Emergence and Diversity of Salmonella enterica Serovar Indiana Isolates with Concurrent Resistance to Ciprofloxacin and Cefotaxime from Patients and Food-Producing Animals in China. Antimicrobial Agents and Chemotherapy, 2016, 60, 3365-3371.	3.2	75
3	Prevalence of Salmonella Isolates from Chicken and Pig Slaughterhouses and Emergence of Ciprofloxacin and Cefotaxime Co-Resistant S. enterica Serovar Indiana in Henan, China. PLoS ONE, 2015, 10, e0144532.	2.5	71
4	Enterotoxigenicity and Antimicrobial Resistance of Staphylococcus aureus Isolated from Retail Food in China. Frontiers in Microbiology, 2017, 8, 2256.	3.5	63
5	Natural Occurrence of Four <i>Alternaria</i> Mycotoxins in Tomato- and Citrus-Based Foods in China. Journal of Agricultural and Food Chemistry, 2015, 63, 343-348.	5.2	58
6	Natural Occurrence of Alternaria Toxins in the 2015 Wheat from Anhui Province, China. Toxins, 2016, 8, 308.	3.4	53
7	Natural Occurrence of Alternaria Toxins in Wheat-Based Products and Their Dietary Exposure in China. PLoS ONE, 2015, 10, e0132019.	2.5	52
8	Prevalence and quantification of Salmonella contamination in raw chicken carcasses at the retail in China. Food Control, 2014, 44, 198-202.	5 . 5	41
9	Co-occurrence of multi-mycotoxins in wheat grains harvested in Anhui province, China. Food Control, 2019, 96, 180-185.	5.5	38
10	Genomic characterization of a large plasmid containing a bla NDM-1 gene carried on Salmonella enterica serovar Indiana C629 isolate from China. BMC Infectious Diseases, 2017, 17, 479.	2.9	29
11	Prevalence and quantification of Campylobacter contamination on raw chicken carcasses for retail sale in China. Food Control, 2017, 75, 196-202.	5.5	28
12	Serovar diversity and antimicrobial resistance of non-typhoidal Salmonella enterica recovered from retail chicken carcasses for sale inÂdifferent regions of China. Food Control, 2017, 81, 46-54.	5.5	26
13	Epidemiological Study on Prevalence, Serovar Diversity, Multidrug Resistance, and CTX-M-Type Extended-Spectrum \hat{I}^2 -Lactamases of <i>Salmonella</i> spp. from Patients with Diarrhea, Food of Animal Origin, and Pets in Several Provinces of China. Antimicrobial Agents and Chemotherapy, 2020, 64, .	3.2	26
14	Genome-Scale Metabolic Models and Machine Learning Reveal Genetic Determinants of Antibiotic Resistance in Escherichia coli and Unravel the Underlying Metabolic Adaptation Mechanisms. MSystems, 2021, 6, e0091320.	3.8	26
15	A risk assessment of salmonellosis linked to chicken meals prepared in households of China. Food Control, 2017, 79, 279-287.	5.5	24
16	Prevalence and Molecular Characteristics of Extended-Spectrum \hat{I}^2 -Lactamase Genes in Escherichia coli Isolated from Diarrheic Patients in China. Frontiers in Microbiology, 2017, 8, 144.	3.5	24
17	Complete Genomic Analysis of a Salmonella enterica Serovar Typhimurium Isolate Cultured From Ready-to-Eat Pork in China Carrying One Large Plasmid Containing mcr-1. Frontiers in Microbiology, 2018, 9, 616.	3.5	24
18	Genomic characterization of an extensively-drug resistance Salmonella enterica serotype Indiana strain harboring blaNDM-1 gene isolated from a chicken carcass in China. Microbiological Research, 2017, 204, 48-54.	5. 3	23

#	Article	IF	Citations
19	Molecular characterization ofblaESBL-producingEscherichia colicultured from pig farms in Ireland. Journal of Antimicrobial Chemotherapy, 2016, 71, 3062-3065.	3.0	22
20	<i>Salmonella</i> harbouring the <i>mcr-1</i> gene isolated from food in China between 2012 and 2016. Journal of Antimicrobial Chemotherapy, 2019, 74, 826-828.	3.0	22
21	Natural co-occurrence of multi-mycotoxins in unprocessed wheat grains from China. Food Control, 2021, 130, 108321.	5.5	22
22	Whole-genome sequencing and gene sharing network analysis powered by machine learning identifies antibiotic resistance sharing between animals, humans and environment in livestock farming. PLoS Computational Biology, 2022, 18, e1010018.	3.2	19
23	A novel disrupted <i>mcr-1</i> gene and a lysogenized phage P1-like sequence detected from a large conjugative plasmid, cultured from a human atypical enteropathogenic <i>Escherichia coli</i> (aEPEC) recovered in China. Journal of Antimicrobial Chemotherapy, 2017, 72, dkw564.	3.0	18
24	Susceptibility (re)-testing of a large collection of Listeria monocytogenes from foods in China from 2012 to 2015 and WGS characterization of resistant isolates. Journal of Antimicrobial Chemotherapy, 2019, 74, 1786-1794.	3.0	18
25	Novel SCC <i>mec</i> type XV (7A) and two pseudo-SCC <i>mec</i> variants in foodborne MRSA in China. Journal of Antimicrobial Chemotherapy, 2022, 77, 903-909.	3.0	18
26	Whole-Genome Sequencing and Machine Learning Analysis of Staphylococcus aureus from Multiple Heterogeneous Sources in China Reveals Common Genetic Traits of Antimicrobial Resistance. MSystems, 2021, 6, e0118520.	3.8	17
27	Natural Occurrence of Beauvericin and Enniatins in Corn- and Wheat-Based Samples Harvested in 2017 Collected from Shandong Province, China. Toxins, 2019, 11, 9.	3.4	16
28	Effects of metal and metalloid pollutants on the microbiota composition of feces obtained from twelve commercial pig farms across China. Science of the Total Environment, 2019, 647, 577-586.	8.0	15
29	Emergence of a <i>Salmonella enterica</i> serovar Typhimurium ST34 isolate, CFSA629, carrying a novel <i>mcr-1.19</i> variant cultured from egg in China. Journal of Antimicrobial Chemotherapy, 2021, 76, 1776-1785.	3.0	14
30	Genomic insights into the pathogenicity and environmental adaptability of <i>Enterococcus hirae</i> R17 isolated from pork offered for retail sale. MicrobiologyOpen, 2017, 6, e00514.	3.0	12
31	Dynamic Fumonisin B2 Production by Aspergillus niger Intented Used in Food Industry in China. Toxins, 2017, 9, 217.	3.4	11
32	Dynamic Ochratoxin A Production by Strains of Aspergillus niger Intended Used in Food Industry of China. Toxins, 2019, 11, 122.	3.4	9
33	Further data on the levels of emerging Fusarium mycotoxins in cereals collected from Tianjin, China. Food Additives and Contaminants: Part B Surveillance, 2021, 14, 74-80.	2.8	6
34	Tetrodotoxin detection and species identification of pufferfish in retail roasted fish fillet by DNA barcoding in China. Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment, 2015, 32, 1-6.	2.3	5
35	Complete Genome and Plasmid Sequences of Seven Isolates of Salmonella enterica subsp. enterica Harboring the mcr-1 Gene Obtained from Food in China. Microbiology Resource Announcements, 2019, 8, .	0.6	5
36	Co-Occurrence of Beauvericin and Enniatins in Edible Vegetable Oil Samples, China. Toxins, 2019, 11, 100.	3.4	5

Fengqin Li

#	Article	IF	CITATIONS
37	Antimicrobial Resistance and Genomic Characterization of Two mcr-1-Harboring Foodborne Salmonella Isolates Recovered in China, 2016. Frontiers in Microbiology, 2021, 12, 636284.	3.5	4
38	Complete Genome Sequence of Enterococcus hirae R17, a Daptomycin-Resistant Bacterium Isolated from Retail Pork in China. Genome Announcements, 2016 , 4 , .	0.8	3
39	Whole Genome Analysis of Three Multi-Drug Resistant Listeria innocua and Genomic Insights Into Their Relatedness With Resistant Listeria monocytogenes. Frontiers in Microbiology, 2021, 12, 694361.	3.5	2
40	Molecular Evolution and Genomic Insights into Community-Acquired Methicillin-Resistant Staphylococcus aureus Sequence Type 88. Microbiology Spectrum, 0, , .	3.0	2
41	Occurrence of CTX-M-123-producing Salmonella Indiana in chicken carcasses: a new challenge for the poultry industry and food safety. Journal of Antimicrobial Chemotherapy, 2019, 74, 3637-3639.	3.0	1