

# Ya-Zhen Qin

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3136650/publications.pdf>

Version: 2024-02-01

76  
papers

1,405  
citations

430874

18  
h-index

395702

33  
g-index

94  
all docs

94  
docs citations

94  
times ranked

1552  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Low EVI1 expression at diagnosis predicted poor outcomes in pediatric Ph-negative B cell precursor acute lymphoblastic leukemia patients. <i>Pediatric Hematology and Oncology</i> , 2022, 39, 97-107.  | 0.8 | 1         |
| 2  | Prognostic significance of TIM-3 expression pattern at diagnosis in patients with t(8;21) acute myeloid leukemia. <i>Leukemia and Lymphoma</i> , 2022, 63, 152-161.   | 1.3 | 4         |
| 3  | Preemptive Interferon- $\gamma$ Therapy Could Protect Against Relapse and Improve Survival of Acute Myeloid Leukemia Patients After Allogeneic Hematopoietic Stem Cell Transplantation: Long-Term Results of Two Registry Studies. <i>Frontiers in Immunology</i> , 2022, 13, 757002.                                     | 4.8 | 13        |
| 4  | Monitoring of post-transplant MLL-PTD as minimal residual disease can predict relapse after allogeneic HSCT in patients with acute myeloid leukemia and myelodysplastic syndrome. <i>BMC Cancer</i> , 2022, 22, 11.   | 2.6 | 2         |
| 5  | Independent prognostic significance of TP53 mutations in adult acute myeloid leukaemia with complex karyotype. <i>International Journal of Laboratory Hematology</i> , 2022, , .  | 1.3 | 4         |
| 6  | Combination of KIT and FLT3-ITD mutation status with minimal residual disease levels guides treatment strategy for adult patients with inv(16) acute myeloid leukemia in first complete remission. <i>Hematological Oncology</i> , 2022, 40, 724-733.   | 1.7 | 2         |
| 7  | Prognostic value of post-transplantation Wilms' tumor gene 1 expression in acute myeloid leukaemia subgroup according to different pre-transplant disease status. <i>International Journal of Laboratory Hematology</i> , 2022, 44, .   | 1.3 | 0         |
| 8  | High PRDM16 expression predicts poor outcomes in adult acute myeloid leukemia patients with intermediate cytogenetic risk: a comprehensive cohort study from a single Chinese center. <i>Leukemia and Lymphoma</i> , 2021, 62, 185-193.   | 1.3 | 3         |
| 9  | Wilms' tumor gene 1 is an independent prognostic factor for pediatric acute myeloid leukemia following allogeneic hematopoietic stem cell transplantation. <i>BMC Cancer</i> , 2021, 21, 292.   | 2.6 | 5         |
| 10 | Development of a poor-prognostic-mutations derived immune prognostic model for acute myeloid leukemia. <i>Scientific Reports</i> , 2021, 11, 4856.  | 3.3 | 7         |
| 11 | The Prognostic Significance of ZNF384 Fusions in Adult Ph-Negative B-Cell Precursor Acute Lymphoblastic Leukemia: A Comprehensive Cohort Study From a Single Chinese Center. <i>Frontiers in Oncology</i> , 2021, 11, 632532.   | 2.8 | 9         |
| 12 | Minimal residual disease monitoring and preemptive immunotherapies for frequent 11q23 rearranged acute leukemia after allogeneic hematopoietic stem cell transplantation. <i>Annals of Hematology</i> , 2021, 100, 1267-1281.   | 1.8 | 3         |
| 13 | The impact of the combination of KIT mutation and minimal residual disease on outcome in t(8;21) acute myeloid leukemia. <i>Blood Cancer Journal</i> , 2021, 11, 67.  | 6.2 | 9         |
| 14 | Profiles of NK cell subsets are associated with successful tyrosine kinase inhibitor discontinuation in chronic myeloid leukemia and changes following interferon treatment. <i>Annals of Hematology</i> , 2021, 100, 2557-2566.  | 1.8 | 4         |
| 15 | Interferon- $\gamma$ as maintenance therapy can significantly reduce relapse in patients with favorable-risk acute myeloid leukemia. <i>Leukemia and Lymphoma</i> , 2021, 62, 2949-2956.  | 1.3 | 14        |
| 16 | PML-RARA transcript levels at the end of induction therapy are associated with prognosis in non-high-risk acute promyelocytic leukaemia with all-trans retinoic acid plus arsenic in front-line therapy: long-term follow-up of a single-centre cohort study. <i>British Journal of Haematology</i> , 2021, 195, 722-730. | 2.5 | 3         |
| 17 | Prognostic value of RASD1 transcript levels in adult Philadelphia-negative B-cell acute lymphoblastic leukemia. <i>Hematology</i> , 2021, 26, 9-15.   | 1.5 | 0         |
| 18 | Preemptive Immunotherapy for Minimal Residual Disease in Patients With t(8;21) Acute Myeloid Leukemia After Allogeneic Hematopoietic Stem Cell Transplantation. <i>Frontiers in Oncology</i> , 2021, 11, 773394.  | 2.8 | 8         |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Overexpressed WT1 exhibits a specific immunophenotype in intermediate and poor cytogenetic risk acute myeloid leukemia. <i>Annals of Hematology</i> , 2020, 99, 215-221.   | 1.8 | 3         |
| 20 | Subgroup Analysis Can Optimize the Relapse-Prediction Cutoff Value for WT1 Expression After Allogeneic Hematologic Stem Cell Transplantation in Acute Myeloid Leukemia. <i>Journal of Molecular Diagnostics</i> , 2020, 22, 188-195.   | 2.8 | 4         |
| 21 | The predictive value of minimal residual disease when facing the inconsistent results detected by real-time quantitative PCR and flow cytometry in NPM1-mutated acute myeloid leukemia. <i>Annals of Hematology</i> , 2020, 99, 73-82.   | 1.8 | 15        |
| 22 | &lt;p&gt;Both Methylation and Copy Number Variation Participated in the Varied Expression of PRAME in Multiple Myeloma&lt;/p&gt;. <i>OncoTargets and Therapy</i> , 2020, Volume 13, 7545-7553.   | 2.0 | 2         |
| 23 | Characterization of somatic mutation-associated microenvironment signatures in acute myeloid leukemia patients based on TCGA analysis. <i>Scientific Reports</i> , 2020, 10, 19037.  | 3.3 | 4         |
| 24 | DPEP1 expression promotes proliferation and survival of leukaemia cells and correlates with relapse in adults with common B cell acute lymphoblastic leukaemia. <i>British Journal of Haematology</i> , 2020, 190, 67-78.  | 2.5 | 11        |
| 25 | Prognostic significance of SET-NUP214 fusion gene in acute leukemia after allogeneic hematopoietic stem cell transplantation. <i>Medicine (United States)</i> , 2020, 99, e23569.  | 1.0 | 6         |
| 26 | High aldehyde dehydrogenase activity at diagnosis predicts relapse in patients with t(8;21) acute myeloid leukemia. <i>Cancer Medicine</i> , 2019, 8, 5459-5467.   | 2.8 | 7         |
| 27 | The prognostic significance of Wilms&#x2013;tumor gene 1 (WT1) expression at diagnosis in adults with Ph-negative B cell precursor acute lymphoblastic leukemia. <i>Annals of Hematology</i> , 2019, 98, 2551-2559.  | 1.8 | 8         |
| 28 | Incidence, risk factors and outcomes of sinusoidal obstruction syndrome after haploidentical allogeneic stem cell transplantation. <i>Annals of Hematology</i> , 2019, 98, 1733-1742.  | 1.8 | 6         |
| 29 | &lt;i>S100A16</i> suppresses the growth and survival of leukaemia cells&Aand correlates with relapse and relapse free survival in adults with Philadelphia chromosome&#x2013;negative B&#x2013;cell acute lymphoblastic leukaemia. <i>British Journal of Haematology</i> , 2019, 185, 836-851. | 2.5 | 7         |
| 30 | Overexpression of WT1 and PRAME predicts poor outcomes of patients with myelodysplastic syndromes with thrombocytopenia. <i>Blood Advances</i> , 2019, 3, 3406-3418.   | 5.2 | 8         |
| 31 | ATRA Could Correct the Defective S1P-Mediated Cytoskeletal Reorganization in Proplatelet Formation of ITP. <i>Blood</i> , 2019, 134, 218-218.  | 1.4 | 1         |
| 32 | Leukemia-propagating cells demonstrate distinctive gene expression profiles compared with other cell fractions from patients with de novo Philadelphia chromosome-positive ALL. <i>Annals of Hematology</i> , 2018, 97, 799-811.   | 1.8 | 0         |
| 33 | Oral arsenic and all-trans retinoic acid for high-risk acute promyelocytic leukemia. <i>Blood</i> , 2018, 131, 2987-2989.  | 1.4 | 36        |
| 34 | Identification of a novel CPSF6-RARG fusion transcript in acute myeloid leukemia resembling acute promyelocytic leukemia. <i>Leukemia</i> , 2018, 32, 2285-2287.   | 7.2 | 32        |
| 35 | The initial level of MLL-partial tandem duplication affects the clinical outcomes in patients with acute myeloid leukemia. <i>Leukemia and Lymphoma</i> , 2018, 59, 967-972.   | 1.3 | 12        |
| 36 | Outcome and Minimal Residual Disease Monitoring in Patients with t(16;21) Acute Myelogenous Leukemia Undergoing Allogeneic Hematopoietic Stem Cell Transplantation. <i>Biology of Blood and Marrow Transplantation</i> , 2018, 24, 163-168.  | 2.0 | 6         |

| #  | ARTICLE   | IF   | CITATIONS |
|----|---|------|-----------|
| 37 | Allogeneic Stem Cell Transplantation versus Tyrosine Kinase Inhibitors Combined with Chemotherapy in Patients with Philadelphia Chromosome-Positive Acute Lymphoblastic Leukemia. <i>Biology of Blood and Marrow Transplantation</i> , 2018, 24, 741-750.               | 2.0  | 36        |
| 38 | Interferon- $\gamma$ Is Effective for Treatment of Minimal Residual Disease in Patients with t(8;21) Acute Myeloid Leukemia After Allogeneic Hematopoietic Stem Cell Transplantation: Results of a Prospective Registry Study. <i>Oncologist</i> , 2018, 23, 1349-1357. | 3.7  | 17        |
| 39 | A seven-color panel including CD34 and TdT could be applied in >97% patients with T cell lymphoblastic leukemia for minimal residual disease detection independent of the initial phenotype. <i>Leukemia Research</i> , 2018, 72, 12-19.                                | 0.8  | 7         |
| 40 | Prevalence and outcomes of uncommon BCR-ABL1 fusion transcripts in patients with chronic myeloid leukaemia: data from a single centre. <i>British Journal of Haematology</i> , 2018, 182, 693-700.  | 2.5  | 31        |
| 41 | Heterogeneous prognosis among KIT mutation types in adult acute myeloid leukemia patients with t(8;21). <i>Blood Cancer Journal</i> , 2018, 8, 76.  | 6.2  | 21        |
| 42 | High EVI1 Expression Predicts Poor Outcomes in Adult Acute Myeloid Leukemia Patients with Intermediate Cytogenetic Risk Receiving Chemotherapy. <i>Medical Science Monitor</i> , 2018, 24, 758-767.   | 1.1  | 17        |
| 43 | Meis1 is critical to the maintenance of human acute myeloid leukemia cells independent of MLL rearrangements. <i>Annals of Hematology</i> , 2017, 96, 567-574.  | 1.8  | 19        |
| 44 | The dynamics of RUNX1-RUNX1T1 transcript levels after allogeneic hematopoietic stem cell transplantation predict relapse in patients with t(8;21) acute myeloid leukemia. <i>Journal of Hematology and Oncology</i> , 2017, 10, 44.                                     | 17.0 | 51        |
| 45 | Methylation pattern of preferentially expressed antigen of melanoma in acute myeloid leukemia and myelodysplastic syndromes. <i>Oncology Letters</i> , 2017, 13, 2823-2830.   | 1.8  | 3         |
| 46 | Impaired Function of Bone Marrow Mesenchymal Stem Cells from Immune Thrombocytopenia Patients in Inducing Regulatory Dendritic Cell Differentiation Through the Notch-1/Jagged-1 Signaling Pathway. <i>Stem Cells and Development</i> , 2017, 26, 1648-1661.            | 2.1  | 36        |
| 47 | PRAME Gene Copy Number Variation Is Related to Its Expression in Multiple Myeloma. <i>DNA and Cell Biology</i> , 2017, 36, 1099-1107.   | 1.9  | 22        |
| 48 | Haploidentical hematopoietic stem cell transplantation for pediatric Philadelphia chromosome-positive acute lymphoblastic leukemia in the imatinib era. <i>Leukemia Research</i> , 2017, 59, 136-141.   | 0.8  | 8         |
| 49 | Concordant optimal molecular and cytogenetic responses at both 3 and 6 months predict a higher probability of MR4.5 achievement in patients with chronic myeloid leukemia treated with imatinib. <i>Leukemia and Lymphoma</i> , 2017, 58, 1384-1393.                    | 1.3  | 3         |
| 50 | The impact of minimal residual disease prior to unmanipulated haploidentical hematopoietic stem cell transplantation in patients with acute myeloid leukemia in complete remission. <i>Leukemia and Lymphoma</i> , 2017, 58, 1135-1143.                                 | 1.3  | 27        |
| 51 | PRAME overexpression predicted good outcome in pediatric B-cell acute lymphoblastic leukemia patients receiving chemotherapy. <i>Leukemia Research</i> , 2017, 52, 43-49.   | 0.8  | 12        |
| 52 | Ruxolitinib/nilotinib cotreatment inhibits leukemia-propagating cells in Philadelphia chromosome-positive ALL. <i>Journal of Translational Medicine</i> , 2017, 15, 184.  | 4.4  | 11        |
| 53 | Cysteine and glycine-rich protein 2 (CSR2) transcript levels correlate with leukemia relapse and leukemia-free survival in adults with B-cell acute lymphoblastic leukemia and normal cytogenetics. <i>Oncotarget</i> , 2017, 8, 35984-36000.                           | 1.8  | 23        |
| 54 | B-cell acute lymphoblastic leukemia associated with SET-NUP214 rearrangement: A case report and review of the literature. <i>Oncology Letters</i> , 2016, 11, 2644-2650.  | 1.8  | 18        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 55 | Homoharringtonine, aclarubicin and cytarabine (HAA) regimen as the first course of induction therapy is highly effective for acute myeloid leukemia with t(8;21). <i>Leukemia Research</i> , 2016, 44, 40-44.   | 0.8 | 29        |
| 56 | Molecular Detection of BCR-ABL in Chronic Myeloid Leukemia. <i>Methods in Molecular Biology</i> , 2016, 1465, 1-15.   | 0.9 | 7         |
| 57 | Low WT1 transcript levels at diagnosis predicted poor outcomes of acute myeloid leukemia patients with t(8;21) who received chemotherapy or allogeneic hematopoietic stem cell transplantation. <i>Chinese Journal of Cancer</i> , 2016, 35, 46.  | 4.9 | 11        |
| 58 | Minimal residual disease monitoring and preemptive immunotherapy in myelodysplastic syndrome after allogeneic hematopoietic stem cell transplantation. <i>Annals of Hematology</i> , 2016, 95, 1233-1240.   | 1.8 | 16        |
| 59 | Combination of White Blood Cell Count at Presentation With Molecular Response at 3 Months Better Predicts Deep Molecular Responses to Imatinib in Newly Diagnosed Chronic-Phase Chronic Myeloid Leukemia Patients. <i>Medicine (United States)</i> , 2016, 95, e2486.                           | 1.0 | 14        |
| 60 | CD38+ CD58 <sup>+</sup> is an independent adverse prognostic factor in paediatric Philadelphia chromosome negative B cell acute lymphoblastic leukaemia patients. <i>Leukemia Research</i> , 2016, 43, 33-38.   | 0.8 | 16        |
| 61 | Philadelphia Chromosome-Positive Acute Lymphoblastic Leukemia: To Allogeneic Stem Cell Transplantation or Not? a Single Center Experience. <i>Blood</i> , 2016, 128, 2308-2308.   | 1.4 | 1         |
| 62 | Janus Kinase Inhibition By Ruxolitinib Combined with Nilotinib Has Superior Anti-Leukemia Propagating Cells Effect in Philadelphia Chromosome-Positive Acute Lymphoblastic Leukemia. <i>Blood</i> , 2016, 128, 4023-4023.   | 1.4 | 0         |
| 63 | C-KIT- Mutated t(8;21)AML Patients with >3log Reduction of MRD Conferred a Very High Relapse and Need HSCT to Improve Outcome. <i>Blood</i> , 2016, 128, 1620-1620.   | 1.4 | 0         |
| 64 | Haploidentical Hematopoietic Stem Cell Transplantation without In Vitro T Cell Depletion for the Treatment of Philadelphia Chromosome-Positive Acute Lymphoblastic Leukemia. <i>Biology of Blood and Marrow Transplantation</i> , 2015, 21, 1110-1116.  | 2.0 | 44        |
| 65 | Allogeneic stem cell transplant may improve the outcome of adult patients with inv(16) acute myeloid leukemia in first complete remission with poor molecular responses to chemotherapy. <i>Leukemia and Lymphoma</i> , 2015, 56, 3116-3123.  | 1.3 | 31        |
| 66 | Prevalence and prognostic significance of c-KIT mutations in core binding factor acute myeloid leukemia: A comprehensive large-scale study from a single Chinese center. <i>Leukemia Research</i> , 2014, 38, 1435-1440.  | 0.8 | 63        |
| 67 | Monitoring Mixed Lineage Leukemia Expression May Help Identify Patients with Mixed Lineage Leukemia-Rearranged Acute Leukemia Who Are at High Risk of Relapse after Allogeneic Hematopoietic Stem Cell Transplantation. <i>Biology of Blood and Marrow Transplantation</i> , 2014, 20, 929-936. | 2.0 | 28        |
| 68 | Prevalence and Prognostic Significance of c-KIT Mutations in Core Binding Factor Acute Myeloid Leukemia: A Comprehensive Large-Scale Study from a Single Chinese Center. <i>Blood</i> , 2014, 124, 1000-1000.   | 1.4 | 0         |
| 69 | Combined use of WT1 and flow cytometry monitoring can promote sensitivity of predicting relapse after allogeneic HSCT without affecting specificity. <i>Annals of Hematology</i> , 2013, 92, 1111-1119.   | 1.8 | 87        |
| 70 | PRAME and WT1 transcripts constitute a good molecular marker combination for monitoring minimal residual disease in myelodysplastic syndromes. <i>Leukemia and Lymphoma</i> , 2013, 54, 1442-1449.  | 1.3 | 23        |
| 71 | Which method better evaluates the molecular response in newly diagnosed chronic phase chronic myeloid leukemia patients with imatinib treatment, BCR-ABLIS or log reduction from the baseline level?. <i>Leukemia Research</i> , 2013, 37, 1035-1040.   | 0.8 | 28        |
| 72 | MRD-directed risk stratification treatment may improve outcomes of t(8;21) AML in the first complete remission: results from the AML05 multicenter trial. <i>Blood</i> , 2013, 121, 4056-4062.  | 1.4 | 277       |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 73 | Low WT1 Expression At Diagnosis Is a Strong Predictor On Poor Outcome In Patients With t(8;21) Acute Myeloid Leukemia. <i>Blood</i> , 2013, 122, 1346-1346.  | 1.4 | 0         |
| 74 | Imatinib Mesylate Versus Allogeneic HSCT for Patients with Chronic Myelogenous Leukemia In Accelerated Phase: A Single Center Experience In China After a 9-Year Follow-up. <i>Blood</i> , 2010, 116, 2347-2347. | 1.4 | 0         |
| 75 | Nucleophosmin mutations in Chinese adults with acute myelogenous leukemia. <i>Annals of Hematology</i> , 2009, 88, 159-166.  | 1.8 | 51        |
| 76 | Abnormal expression of the programmed cell death 5 gene in acute and chronic myeloid leukemia. <i>Leukemia Research</i> , 2006, 30, 1159-1165.   | 0.8 | 50        |