
## Jesus M Porres

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/314317/publications.pdf Version: 2024-02-01



IFSUS M PODDES

#	Article	IF	CITATIONS
1	Papel del ácido fÃŧico en las legumbres. Journal of Physiology and Biochemistry, 2000, 56, 283-294.	3.0	216
2	Phytase enzymology, applications, and biotechnology. Biotechnology Letters, 2003, 25, 1787-1794.	2.2	183
3	Knockout of cellular glutathione peroxidase gene renders mice susceptible to diquat-induced oxidative stress. Free Radical Biology and Medicine, 1999, 27, 605-611.	2.9	118
4	Different Sensitivity of RecombinantAspergillus nigerPhytase (r-PhyA) andEscherichia colipH 2.5 Acid Phosphatase (r-AppA) to Trypsin and Pepsinin Vitro. Archives of Biochemistry and Biophysics, 1999, 365, 262-267.	3.0	93
5	Shifting the pH Profile of Aspergillus niger PhyA Phytase To Match the Stomach pH Enhances Its Effectiveness as an Animal Feed Additive. Applied and Environmental Microbiology, 2006, 72, 4397-4403.	3.1	77
6	Nutritional assessment of raw and germinated pea (Pisum sativum L.) protein and carbohydrate by in vitro and in vivo techniques. Nutrition, 2005, 21, 230-239.	2.4	63
7	Site-directed mutagenesis of Aspergillus niger NRRL 3135 phytase at residue 300 to enhance catalysis at pH 4.0. Biochemical and Biophysical Research Communications, 2002, 297, 1016-1020.	2.1	59
8	Phytase: Source, Structure and Application. , 2007, , 505-529.		56
9	Improvement of the antioxidant and hypolipidaemic effects of cowpea flours ( <i>Vigna) Tj ETQq1 1 0.784314 the Science of Food and Agriculture, 2015, 95, 1207-1216.</i>	rgBT /Overl 3.5	ock 10 Tf 50 54
10	Dietary Intrinsic Phytate Protects Colon from Lipid Peroxidation in Pigs with a Moderately High Dietary Iron Intake. Proceedings of the Society for Experimental Biology and Medicine, 1999, 221, 80-86.	1.8	53
11	Effects of germination on the composition and nutritive value of proteins in , L. Food Chemistry, 2005, 93, 671-679.	8.2	49
12	Spatial-temporal parameters of gait in women with fibromyalgia. Clinical Rheumatology, 2009, 28, 595-598.	2.2	45
13	Effects of high-whey-protein intake and resistance training on renal, bone and metabolic parameters in rats. British Journal of Nutrition, 2011, 105, 836-845.	2.3	45
14	Nutritional Evaluation of Pea (Pisum sativumL.) Protein Diets after Mild Hydrothermal Treatment and with and without Added Phytase. Journal of Agricultural and Food Chemistry, 2003, 51, 2415-2420.	5.2	37
15	Digestive utilisation of protein and amino acids from raw and heated lentils by growing rats. Journal of the Science of Food and Agriculture, 2002, 82, 1740-1747.	3.5	30
16	Health promoting effects of Lupin (Lupinus albus var. multolupa) protein hydrolyzate and insoluble fiber in a diet-induced animal experimental model of hypercholesterolemia. Food Research International, 2013, 54, 1471-1481.	6.2	30
17	Functional expression of keratinase (kerA) gene from Bacillus licheniformis in Pichia pastoris. Biotechnology Letters, 2002, 24, 631-636.	2.2	29
18	Influence of intracerebroventricular or intraperitoneal administration of cannabinoid receptor agonist (WIN 55,212-2) and inverse agonist (AM 251) on the regulation of food intake and hypothalamic serotonin levels. British Journal of Nutrition, 2009, 101, 1569.	2.3	29

JESUS M PORRES

#	Article	IF	CITATIONS
19	Aerobic interval exercise improves parameters of nonalcoholic fatty liver disease (NAFLD) and other alterations of metabolic syndrome in obese Zucker rats. Applied Physiology, Nutrition and Metabolism, 2015, 40, 1242-1252.	1.9	28
20	Co-inoculation of Halomonas maura and Ensifer meliloti to improve alfalfa yield in saline soils. Applied Soil Ecology, 2015, 87, 81-86.	4.3	28
21	High-protein diets and renal status in rats. Nutricion Hospitalaria, 2013, 28, 232-7.	0.3	28
22	Comparative impacts of glutathione peroxidase-1 gene knockout on oxidative stress induced by reactive oxygen and nitrogen species in mouse hepatocytes. Biochemical Journal, 2001, 359, 687-695.	3.7	26
23	Effect of Natural and Controlled Fermentation on Chemical Composition and Nutrient Dialyzability from Beans (Phaseolus vulgarisL.). Journal of Agricultural and Food Chemistry, 2003, 51, 5144-5149.	5.2	26
24	Nutritional Potential of Raw and Free α-Galactosides Lupin (Lupinus albus Var. multolupa) Seed Flours. Effect of Phytase Treatment on Nitrogen and Mineral Dialyzability. Journal of Agricultural and Food Chemistry, 2005, 53, 3088-3094.	5.2	25
25	Evaluation of zinc and magnesium bioavailability from pea (Pisum sativum, L.) sprouts. Effect of illumination and different germination periods. International Journal of Food Science and Technology, 2006, 41, 618-626.	2.7	24
26	Beneficial effects of legumes on parameters of the metabolic syndrome: a systematic review of trials in animal models. British Journal of Nutrition, 2016, 116, 402-424.	2.3	22
27	Ergogenic effects of quercetin supplementation in trained rats. Journal of the International Society of Sports Nutrition, 2013, 10, 3.	3.9	21
28	Effects of a combined intervention with a lentil protein hydrolysate and a mixed training protocol on the lipid metabolism and hepatic markers of NAFLD in Zucker rats. Food and Function, 2018, 9, 830-850.	4.6	21
29	Nutritional Value. , 2007, , 47-93.		21
30	Ca and P bioavailability of processed lentils as affected by dietary fiber and phytic acid content. Nutrition Research, 1999, 19, 49-64.	2.9	18
31	Novel effects of the cannabinoid inverse agonist AM 251 on parameters related to metabolic syndrome in obese Zucker rats. Metabolism: Clinical and Experimental, 2013, 62, 1641-1650.	3.4	17
32	Effects of interval aerobic training combined with strength exercise on body composition, glycaemic and lipid profile and aerobic capacity of obese rats. Journal of Sports Sciences, 2016, 34, 1452-1460.	2.0	17
33	Germination Improves the Polyphenolic Profile and Functional Value of Mung Bean (Vigna radiata L.). Antioxidants, 2020, 9, 746.	5.1	17
34	Comparative impacts of glutathione peroxidase-1 gene knockout on oxidative stress induced by reactive oxygen and nitrogen species in mouse hepatocytes. Biochemical Journal, 2001, 359, 687.	3.7	16
35	Effect of Heat Treatment and Mineral and Vitamin Supplementation on the Nutritive Use of Protein and Calcium From Lentils (Lens culinaris M.) in Growing Rats. Nutrition, 2003, 19, 451-456.	2.4	16
36	Nutritional evaluation of protein, phosphorus, calcium and magnesium bioavailability from lupin (Lupinus albus var. multolupa)-based diets in growing rats: effect of α-galactoside oligosaccharide extraction and phytase supplementation. British Journal of Nutrition, 2006, 95, 1102-1111.	2.3	16

JESUS M PORRES

#	Article	IF	CITATIONS
37	Medicago sativa L., a functional food to relieve hypertension and metabolic disorders in a spontaneously hypertensive rat model. Journal of Functional Foods, 2016, 26, 470-484.	3.4	16
38	Improvement in food intake and nutritive utilization of protein from Lupinus albus var. multolupa protein isolates supplemented with ascorbic acid. Food Chemistry, 2007, 103, 944-951.	8.2	15
39	Antitumor Effect of the Ethanolic Extract from Seeds of Euphorbia lathyris in Colorectal Cancer. Nutrients, 2021, 13, 566.	4.1	15
40	The Combined Intervention with Germinated Vigna radiata and Aerobic Interval Training Protocol Is an Effective Strategy for the Treatment of Non-Alcoholic Fatty Liver Disease (NAFLD) and Other Alterations Related to the Metabolic Syndrome in Zucker Rats. Nutrients, 2017, 9, 774.	4.1	14
41	Effects of the dietary amount and source of protein, resistance training and anabolic-androgenic steroids on body weight and lipid profile of rats. Nutricion Hospitalaria, 2013, 28, 127-36.	0.3	14
42	Bioavailability of phytic acid–phosphorus and magnesium from lentils (Lens culinaris m.) in growing rats: Influence of thermal treatment and vitamin-mineral supplementation. Nutrition, 2004, 20, 794-799.	2.4	13
43	High-intensity Exercise Modifies the Effects of Stanozolol on Brain Oxidative Stress in Rats. International Journal of Sports Medicine, 2015, 36, 984-991.	1.7	13
44	High-protein diet induces oxidative stress in rat brain: protective action of high-intensity exercise against lipid peroxidation. Nutricion Hospitalaria, 2014, 31, 866-74.	0.3	12
45	Nitrogen Fractions and Mineral Content in Different Lupin Species ( <i>Lupinus albus</i> , <i>Lupinus) Tj ETQq1 I Journal of Agricultural and Food Chemistry, 2007, 55, 7445-7452.</i>	1 0.784314 5.2	rgBT /Overld 11
46	Natural Fermentation of Cowpea (Vigna unguiculata) Flour Improves the Nutritive Utilization of Indispensable Amino Acids and Phosphorus by Growing Rats. Nutrients, 2020, 12, 2186.	4.1	11
47	Antioxidant and antiproliferative potential of ethanolic extracts from Moringa oleifera, Tropaeolum tuberosum and Annona cherimola in colorrectal cancer cells. Biomedicine and Pharmacotherapy, 2021, 143, 112248.	5.6	11
48	Fecal fermentation products of common bean-derived fiber inhibit C/EBPα and PPARγ expression and lipid accumulation but stimulate PPARÎ′ and UCP2 expression in the adipogenesis of 3T3-L1 cells. Journal of Nutritional Biochemistry, 2018, 60, 9-15.	4.2	10
49	Effect of treatment with α-galactosidase, tannase or a cell-wall-degrading enzyme complex on the nutritive utilisation of protein and carbohydrates from pea (Pisum sativum L.) flour. Journal of the Science of Food and Agriculture, 2007, 87, 1356-1363.	3.5	9
50	Aerobic interval exercise improves renal functionality and affects mineral metabolism in obese Zucker rats. American Journal of Physiology - Renal Physiology, 2019, 316, F90-F100.	2.7	9
51	Bioavailability of calcium and magnesium from faba beans(Vicia faba L varmajor), soaked in different pH solutions and cooked, in growing rats. Journal of the Science of Food and Agriculture, 2004, 84, 1514-1520.	3.5	8
52	In Vivo Nutritional Assessment of the Microalga Nannochloropsis gaditana and Evaluation of the Antioxidant and Antiproliferative Capacity of Its Functional Extracts. Marine Drugs, 2022, 20, 318.	4.6	8
53	Improvement of iron availability from phytase-treated Pisum sativum, L. flour. Food Chemistry, 2007, 103, 389-395.	8.2	7
54	A combined healthy strategy for successful weight loss, weight maintenance and improvement of hepatic lipid metabolism. Journal of Nutritional Biochemistry, 2020, 85, 108456.	4.2	7

JESUS M PORRES

#	Article	IF	CITATIONS
55	Synthesis of [77Se]-methylselenocysteine when preparing sauerkraut in the presence of [77Se]-selenite. Metabolic transformation of [77Se]-methylselenocysteine in Wistar rats determined by LC–IDA–ICP–MS. Analytical and Bioanalytical Chemistry, 2014, 406, 7949-7958.	3.7	6
56	The combined treatment with lentil protein hydrolysate and a mixed training protocol is an efficient lifestyle intervention to manage cardiovascular and renal alterations in obese Zucker rats. European Journal of Nutrition, 2020, 59, 3473-3490.	3.9	6
57	High-Intensity Exercise May Compromise Renal Morphology in Rats. International Journal of Sports Medicine, 2014, 35, 639-644.	1.7	5
58	Caloric restriction, physical exercise, and CB1 receptor blockade as an efficient combined strategy for bodyweight control and cardiometabolic status improvement in male rats. Scientific Reports, 2021, 11, 4286.	3.3	5
59	Anemonia sulcata and Its Symbiont Symbiodinium as a Source of Anti-Tumor and Anti-Oxidant Compounds for Colon Cancer Therapy: A Preliminary In Vitro Study. Biology, 2021, 10, 134.	2.8	5
60	In vitro evidence of the antitumor capacity of <i>Solanaceae</i> and <i>Cucurbitaceae</i> in colon cancer: A systematic review. Critical Reviews in Food Science and Nutrition, 2022, 62, 6293-6314.	10.3	5
61	Effect of phytic acid degradation by soaking and exogenous phytase on the bioavailability of magnesium and zinc from Pisum sativum, L European Food Research and Technology, 2007, 226, 105-111.	3.3	4
62	Effects of the amount and source of dietary protein on bone status in rats. Food and Function, 2014, 5, 716.	4.6	4
63	Whey Versus Soy Protein Diets and Renal Status in Rats. Journal of Medicinal Food, 2014, 17, 1011-1016.	1.5	4
64	Exploring Honeybee Abdominal Anatomy through Micro-CT and Novel Multi-Staining Approaches. Insects, 2022, 13, 556.	2.2	4
65	Effects of hydroalcoholic α-galactoside extraction and phytase supplementation on the nutritive utilization of manganese, iron, zinc and potassium from lupin (Lupinus albus var. multolupa)-based diets in growing rats. Food Chemistry, 2008, 109, 554-563.	8.2	3
66	Carbohydrates digestibility and faecal microbiota composition in rats fed diets based on raw or fermented Vigna unguiculata seed meal as the only protein source. , 2022, 1, 100022.		3
67	Bioavailability and biotransformation of linolenic acid from basil seed oil as a novel source of omega-3 fatty acids tested on a rat experimental model. Food and Function, 2022, 13, 7614-7628.	4.6	3
68	Selenium, Selenoproteins, and Age-Related Disorders. , 2013, , 227-239.		2
69	Effects of a moderately high-protein diet and interval aerobic training combined with strength-endurance exercise on markers of bone metabolism, microarchitecture and turnover in obese Zucker rats. Bone, 2016, 92, 116-123.	2.9	2
70	Changes on metabolic parameters induced by acute cannabinoid administration (CBD, THC) in a rat experimental model of nutritional vitamin A deficiency. Nutricion Hospitalaria, 2013, 28, 857-67.	0.3	2
71	Stanozolol Decreases Bone Turnover Markers, Increases Mineralization, and Alters Femoral Geometry in Male Rats. Calcified Tissue International, 2016, 98, 609-618.	3.1	1

Legumes, Genome Maintenance, and Optimal Health. , 2013, , 321-334.

#	Article	IF	CITATIONS
73	Effects of Hypertrophy Exercise in Bone Turnover Markers and Structure in Growing Male Rats. International Journal of Sports Medicine, 2017, 38, 418-425.	1.7	Ο
74	Efectos del ejercicio aeróbico interválico, combinado con entrenamiento de fuerza y de la restricción calórica, sobre la composición corporal de ratas obesas. Revista Andaluza De Medicina Del Deporte, 2017, 10, 3-8.	0.1	0
75	Efectos de un protocolo de entrenamiento de alta intensidad sobre marcadores fisiológicos de estrés en ratas. [Physiological effects of the stress induced by a high-intensity exercise protocol in rats] RICYDE Revista Internacional De Ciencias Del Deporte, 2015, 11, 145-162.	0.2	0