
## Haejun Yim

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3145436/publications.pdf Version: 2024-02-01



Ηλειιινι Υιμ

#	Article	IF	CITATIONS
1	Wound healing ability of acellular fish skin and bovine collagen grafts for split-thickness donor sites in burn patients: Characterization of acellular grafts and clinical application. International Journal of Biological Macromolecules, 2022, 205, 452-461.	7.5	23
2	Time-varying discrimination accuracy of longitudinal biomarkers for the prediction of mortality compared to assessment at fixed time point in severe burns patients. BMC Emergency Medicine, 2021, 21, 1.	1.9	23
3	Trajectories of longitudinal biomarkers for mortality in severely burned patients. Scientific Reports, 2020, 10, 16193.	3.3	8
4	Development of a risk prediction model (Hangang) and comparison with clinical severity scores in burn patients. PLoS ONE, 2019, 14, e0211075.	2.5	11
5	Reliability of resting energy expenditure in major burns: Comparison between measured and predictive equations. Clinical Nutrition, 2019, 38, 2763-2769.	5.0	9
6	Diagnostic performance of plasma and urine neutrophil gelatinase-associated lipocalin, cystatin C, and creatinine for acute kidney injury in burn patients: A prospective cohort study. PLoS ONE, 2018, 13, e0199600.	2.5	8
7	Clinical Outcome of Cryopreserved Acellular Dermal Matrix for Full-Thickness Burns. Macromolecular Research, 2018, 26, 780-787.	2.4	ο
8	Prevalence of Malnutrition in Hospitalized Patients: a Multicenter Cross-sectional Study. Journal of Korean Medical Science, 2018, 33, e10.	2.5	79
9	Analysis of prognostic factors for acute kidney injury with continuous renal replacement therapy in severely burned patients. Burns, 2017, 43, 1418-1426.	1.9	11
10	Effectiveness and Safety of a Thermosensitive Hydrogel Cultured Epidermal Allograft for Burns. Advances in Skin and Wound Care, 2017, 30, 559-564.	1.0	2
11	Assessment of Plasma Neutrophil Gelatinase-Associated Lipocalin for Early Detection of Acute Kidney Injury and Prediction of Mortality in Severely Burned Patients. Journal of Burn Care and Research, 2017, 39, 1.	0.4	5
12	Subgroup analysis of continuous renal replacement therapy in severely burned patients. PLoS ONE, 2017, 12, e0189057.	2.5	11
13	Does inhalation injury predict mortality in burns patients or require redefinition?. PLoS ONE, 2017, 12, e0185195.	2.5	18
14	Effect of extracorporeal shock wave therapy on scar pain in burn patients. Medicine (United States), 2016, 95, e4575.	1.0	27
15	Effectiveness of wound healing using the novel collagen dermal substitute INSUREGRAF®. RSC Advances, 2016, 6, 59692-59701.	3.6	5
16	Development of cell-laden 3D scaffolds for efficient engineered skin substitutes by collagen gelation. RSC Advances, 2016, 6, 21439-21447.	3.6	63
17	The application of cultured epithelial autografts improves survival in burns. Wound Repair and Regeneration, 2015, 23, 340-344.	3.0	25
18	Evaluation of diagnostic biomarkers for acute kidney injury in major burn patients. Annals of Surgical Treatment and Research, 2015, 88, 281.	1.0	19

Haejun Yim

#	Article	IF	CITATIONS
19	Serum cystatin C and microalbuminuria in burn patients with acute kidney injury. European Journal of Clinical Investigation, 2015, 45, 594-600.	3.4	17
20	Epidemiological trends and risk factors in major burns patients in South Korea: A 10-year experience. Burns, 2015, 41, 181-187.	1.9	27
21	Serum Transthyretin Level Is Associated With Clinical Severity Rather Than Nutrition Status in Massively Burned Patients. Journal of Parenteral and Enteral Nutrition, 2014, 38, 966-972.	2.6	16
22	Assessment of biochemical markers in the early post-burn period for predicting acute kidney injury and mortality in patients with major burn injury: comparison of serum creatinine, serum cystatin-C, plasma and urine neutrophil gelatinase-associated lipocalin. Critical Care, 2014, 18, R151.	5.8	52
23	A clinical trial designed to evaluate the safety and effectiveness of a thermosensitive hydrogel-type cultured epidermal allograft for deep second-degree burns. Burns, 2014, 40, 1642-1649.	1.9	9
24	The effect of burn rehabilitation massage therapy on hypertrophic scar after burn: A randomized controlled trial. Burns, 2014, 40, 1513-1520.	1.9	95
25	Improvement of burn pain management through routine pain monitoring and pain management protocol. Burns, 2013, 39, 619-624.	1.9	19
26	Population Pharmacokinetic Analysis of Fluconazole To Predict Therapeutic Outcome in Burn Patients with Candida Infection. Antimicrobial Agents and Chemotherapy, 2013, 57, 1006-1011.	3.2	18
27	Prediction of clinical outcomes for massively-burned patients via serum transthyretin levels in the early postburn period. Journal of Trauma, 2012, 72, 999-1005.	2.3	19
28	Change of serum phosphate level and clinical outcome of hypophosphatemia in massive burn patient. Journal of Trauma and Acute Care Surgery, 2012, 73, 1298-1302.	2.1	16
29	Changes in the Levels of Interleukins 6, 8, and 10, Tumor Necrosis Factor Alpha, and Granulocyte-colony Stimulating Factor in Korean Burn Patients: Relation to Burn Size and Postburn Time. Annals of Laboratory Medicine, 2012, 32, 339-344.	2.5	38
30	Clinical study of cultured epithelial autografts in liquid suspension in severe burn patients. Burns, 2011, 37, 1067-1071.	1.9	38
31	Investigation of relationship between inhalation injury assessment and prognosis in burn patients. [Chapchi] Journal Taehan Oekwa Hakhoe, 2011, 81, 1.	1.1	8
32	Serum Lactate and Base Deficit: Early Predictors of Morbidity and Mortality in Burn Patients with Inhalation Injury. [Chapchi] Journal Taehan Oekwa Hakhoe, 2011, 80, 84.	1.1	3
33	Necrotizing Fasciitis Following a Small Burn. [Chapchi] Journal Taehan Oekwa Hakhoe, 2010, 79, 71.	1.1	7
34	A Clinical Study of Stevens-Johnson Syndrome and Toxic Epidermal Necrolysis: Efficacy of Treatment in Burn Intensive Care Unit. [Chapchi] Journal Taehan Oekwa Hakhoe, 2010, 78, 133.	1,1	4
35	Population pharmacokinetics of meropenem in burn patients. Journal of Antimicrobial Chemotherapy, 2010, 65, 2428-2435.	3.0	47
36	The use of AlloDerm on major burn patients: AlloDerm prevents post-burn joint contracture. Burns, 2010, 36, 322-328.	1.9	80