Daigo Sumi

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/3154694/daigo-sumi-publications-by-year.pdf

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

| 80 | 2,774 | 31 | 51 |
|-------------|----------------|---------|---------|
| papers | citations | h-index | g-index |
| 83 | 2,995 | 5.1 | 4.61 |
| ext. papers | ext. citations | avg, IF | L-index |

| # | Paper | IF | Citations |
|----|--|-----|-----------|
| 80 | Bismuth 2022 , 121-139 | | O |
| 79 | Elevated serum periostin levels among arsenic-exposed individuals and their associations with the features of asthma <i>Chemosphere</i> , 2022 , 134277 | 8.4 | 0 |
| 78 | Effects of individual amino acid mutations of zinc transporter ZIP8 on manganese- and cadmium-transporting activity. <i>Biochemical and Biophysical Research Communications</i> , 2022 , 616, 26-32 | 3.4 | 2 |
| 77 | T helper 2-driven immune dysfunction in chronic arsenic-exposed individuals and its link to the features of allergic asthma. <i>Toxicology and Applied Pharmacology</i> , 2021 , 420, 115532 | 4.6 | 3 |
| 76 | Arsenite suppresses IL-2-dependent tumoricidal activities of natural killer cells. <i>Toxicology and Applied Pharmacology</i> , 2021 , 412, 115353 | 4.6 | 3 |
| 75 | Comparisons of segment-specific toxicity of platinum-based agents and cadmium using S1, S2, and S3 cells derived from mouse kidney proximal tubules. <i>Toxicology in Vitro</i> , 2021 , 75, 105179 | 3.6 | 1 |
| 74 | Arsenic Secondary Methylation Capacity Is Inversely Associated with Arsenic Exposure-Related Muscle Mass Reduction. <i>International Journal of Environmental Research and Public Health</i> , 2021 , 18, | 4.6 | 2 |
| 73 | Arsenite suppresses the transcriptional activity of EVI1 through the binding to CCHC-type Zn finger domain. <i>Biochemical and Biophysical Research Communications</i> , 2020 , 529, 910-915 | 3.4 | 1 |
| 72 | Arsenite suppresses NO production evoked by lipopolysaccharide and poly(I:C) via the suppression of interferon-lexpression in RAW264.7 cells. <i>Journal of Toxicological Sciences</i> , 2019 , 44, 83-92 | 1.9 | O |
| 71 | Chronic exposure to submicromolar arsenite promotes the migration of human esophageal Het1A cells induced by heparin-binding EGF-like growth factor. <i>Archives of Toxicology</i> , 2019 , 93, 3523-3534 | 5.8 | 2 |
| 70 | Toxicometallomics of Cadmium, Manganese and Arsenic with Special Reference to the Roles of Metal Transporters. <i>Toxicological Research</i> , 2019 , 35, 311-317 | 3.7 | 9 |
| 69 | Synergistic augmentation of ATP-induced interleukin-6 production by arsenite in HaCaT cells. <i>Archives of Toxicology</i> , 2016 , 90, 1307-13 | 5.8 | 3 |
| 68 | Arsenic trioxide augments all-trans retinoic acid-induced differentiation of HL-60 cells. <i>Life Sciences</i> , 2016 , 149, 42-50 | 6.8 | 7 |
| 67 | Hydrogen peroxide triggers a novel alternative splicing of arsenic (+3lbxidation state) methyltransferase gene. <i>Biochemical and Biophysical Research Communications</i> , 2016 , 480, 18-22 | 3.4 | |
| 66 | High accumulation of arsenic in the esophagus of mice after exposure to arsenite. <i>Archives of Toxicology</i> , 2015 , 89, 1751-8 | 5.8 | 4 |
| 65 | Arsenite retards the cardiac differentiation of rat cardiac myoblast H9c2 cells. <i>Biochemical and Biophysical Research Communications</i> , 2013 , 436, 175-9 | 3.4 | 15 |
| 64 | Involvement of Nrf2 activation in the upregulation of S100A9 by exposure to inorganic arsenite. <i>International Journal of Molecular Medicine</i> , 2013 , 31, 259-64 | 4.4 | 4 |

(2008-2012)

| 63 | Role of arsenic (+3 oxidation state) methyltransferase in arsenic metabolism and toxicity. <i>Biological and Pharmaceutical Bulletin</i> , 2012 , 35, 1870-5 | 2.3 | 27 |
|----|---|-----|----|
| 62 | Alternative splicing variants of human arsenic (+3 oxidation state) methyltransferase. <i>Biochemical and Biophysical Research Communications</i> , 2011 , 415, 48-53 | 3.4 | 9 |
| 61 | Peroxiredoxin 6 is a molecular target for 1,2-naphthoquinone, an atmospheric electrophile, in human pulmonary epithelial A549 cells. <i>Journal of Toxicological Sciences</i> , 2011 , 36, 817-21 | 1.9 | 12 |
| 60 | Chronic exposure to arsenite induces S100A8 and S100A9 expression in rat RBL-2H3 mast cells. Journal of Toxicological Sciences, 2011 , 36, 135-9 | 1.9 | 3 |
| 59 | Inhibition of DNA binding activity of cAMP response element-binding protein by 1,2-naphthoquinone through chemical modification of Cys-286. <i>Chemico-Biological Interactions</i> , 2011 , 192, 272-7 | 5 | 13 |
| 58 | Rat H9c2 cardiac myocytes are sensitive to arsenite due to a modest activation of transcription factor Nrf2. <i>Archives of Toxicology</i> , 2011 , 85, 1509-16 | 5.8 | 19 |
| 57 | Initial response and cellular protection through the Keap1/Nrf2 system during the exposure of primary mouse hepatocytes to 1,2-naphthoquinone. <i>Chemical Research in Toxicology</i> , 2011 , 24, 559-67 | 4 | 47 |
| 56 | Reduction of arsenic-induced cytotoxicity through Nrf2/HO-1 signaling in HepG2 cells. <i>Journal of Toxicological Sciences</i> , 2010 , 35, 419-23 | 1.9 | 34 |
| 55 | 1,2-Naphthoquinone suppresses lipopolysaccharide-dependent activation of IKK/INF-B/NO signaling: an alternative mechanism for the disturbance of inducible NO synthase-catalyzed NO formation. <i>Journal of Toxicological Sciences</i> , 2010 , 35, 891-8 | 1.9 | 15 |
| 54 | Diesel exhaust particles induce a Th2 phenotype in mouse nalle mononuclear cells in vitro. <i>Experimental and Therapeutic Medicine</i> , 2010 , 1, 761-767 | 2.1 | 4 |
| 53 | Signal transduction pathways and transcription factors triggered by arsenic trioxide in leukemia cells. <i>Toxicology and Applied Pharmacology</i> , 2010 , 244, 385-92 | 4.6 | 44 |
| 52 | Carbon monoxide derived from heme oxygenase-2 mediates reduction of methylmercury toxicity in SH-SY5Y cells. <i>Toxicology and Applied Pharmacology</i> , 2010 , 249, 86-90 | 4.6 | 5 |
| 51 | Role of aquaporin 9 in cellular accumulation of arsenic and its cytotoxicity in primary mouse hepatocytes. <i>Toxicology and Applied Pharmacology</i> , 2009 , 237, 232-6 | 4.6 | 43 |
| 50 | Catechol estrogens mediated activation of Nrf2 through covalent modification of its quinone metabolite to Keap1. <i>Journal of Toxicological Sciences</i> , 2009 , 34, 627-35 | 1.9 | 35 |
| 49 | L-Xylulose reductase is involved in 9,10-phenanthrenequinone-induced apoptosis in human T lymphoma cells. <i>Free Radical Biology and Medicine</i> , 2008 , 44, 1191-202 | 7.8 | 52 |
| 48 | Redox cycling of 9,10-phenanthraquinone to cause oxidative stress is terminated through its monoglucuronide conjugation in human pulmonary epithelial A549 cells. <i>Free Radical Biology and Medicine</i> , 2008 , 44, 1645-55 | 7.8 | 48 |
| 47 | Biological Effects of and Responses to Exposure to Electrophilic Environmental Chemicals. <i>Journal of Health Science</i> , 2008 , 54, 267-272 | | 14 |
| 46 | Downregulation of arginase II and renal apoptosis by inorganic mercury: overexpression of arginase II reduces its apoptosis. <i>Archives of Toxicology</i> , 2008 , 82, 67-73 | 5.8 | 12 |

| 45 | Reduction of arginase I activity and manganese levels in the liver during exposure of rats to methylmercury: a possible mechanism. <i>Archives of Toxicology</i> , 2008 , 82, 803-8 | 5.8 | 15 |
|----|---|------|-----|
| 44 | Activation of the Nrf2 pathway, but decreased gamma-glutamylcysteine synthetase heavy subunit chain levels and caspase-3-dependent apoptosis during exposure of primary mouse hepatocytes to diphenylarsinic acid. <i>Toxicology and Applied Pharmacology</i> , 2007 , 223, 218-24 | 4.6 | 11 |
| 43 | Effects of naphthoquinone on airway responsiveness in the presence or absence of antigen in mice. <i>Archives of Toxicology</i> , 2007 , 81, 575-81 | 5.8 | 20 |
| 42 | Chemical knockdown of protein-tyrosine phosphatase 1B by 1,2-naphthoquinone through covalent modification causes persistent transactivation of epidermal growth factor receptor. <i>Journal of Biological Chemistry</i> , 2007 , 282, 33396-33404 | 5.4 | 87 |
| 41 | Effects of a pomegranate fruit extract rich in punicalagin on oxidation-sensitive genes and eNOS activity at sites of perturbed shear stress and atherogenesis. <i>Cardiovascular Research</i> , 2007 , 73, 414-23 | 9.9 | 69 |
| 40 | Arsenic and Other Metal Contamination of Groundwater in the Mekong River Delta, Vietnam. <i>Journal of Health Science</i> , 2007 , 53, 344-346 | | 22 |
| 39 | Arsenic: signal transduction, transcription factor, and biotransformation involved in cellular response and toxicity. <i>Annual Review of Pharmacology and Toxicology</i> , 2007 , 47, 243-62 | 17.9 | 206 |
| 38 | 1,2-Naphthoquinone disrupts the function of cAMP response element-binding protein through covalent modification. <i>Biochemical and Biophysical Research Communications</i> , 2007 , 361, 243-8 | 3.4 | 25 |
| 37 | Cytoprotective role of Nrf2/Keap1 system in methylmercury toxicity. <i>Biochemical and Biophysical Research Communications</i> , 2007 , 363, 645-50 | 3.4 | 109 |
| 36 | Inhibition of endothelial nitric oxide synthase activity and suppression of endothelium-dependent vasorelaxation by 1,2-naphthoquinone, a component of diesel exhaust particles. <i>Archives of Toxicology</i> , 2006 , 80, 280-5 | 5.8 | 33 |
| 35 | Serine 1179 phosphorylation of endothelial nitric oxide synthase caused by 2,4,6-trinitrotoluene through PI3K/Akt signaling in endothelial cells. <i>Toxicology and Applied Pharmacology</i> , 2006 , 214, 55-60 | 4.6 | 6 |
| 34 | Modulating role of estradiol on arginase II expression in hyperlipidemic rabbits as an atheroprotective mechanism. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006 , 103, 10485-10490 | 11.5 | 35 |
| 33 | Selective iNOS inhibitor, ONO1714 successfully retards the development of high-cholesterol diet induced atherosclerosis by novel mechanism. <i>Atherosclerosis</i> , 2006 , 187, 316-24 | 3.1 | 32 |
| 32 | Sulforaphane, an activator of Nrf2, suppresses cellular accumulation of arsenic and its cytotoxicity in primary mouse hepatocytes. <i>FEBS Letters</i> , 2006 , 580, 1771-4 | 3.8 | 79 |
| 31 | Pomegranate juice protects nitric oxide against oxidative destruction and enhances the biological actions of nitric oxide. <i>Nitric Oxide - Biology and Chemistry</i> , 2006 , 15, 93-102 | 5 | 116 |
| 30 | Sp1 transcription factor expression is regulated by estrogen-related receptor alpha1. <i>Biochemical and Biophysical Research Communications</i> , 2005 , 328, 165-72 | 3.4 | 18 |
| 29 | Monomethylarsonous Acid Inhibits Endothelial Nitric Oxide Synthase Activity. <i>Journal of Health Science</i> , 2005 , 51, 728-730 | | 6 |
| 28 | 2,4,6-Trinitrotoluene inhibits endothelial nitric oxide synthase activity and elevates blood pressure in rats. <i>Archives of Toxicology</i> , 2005 , 79, 705-10 | 5.8 | 10 |

(2001-2005)

| 27 | in ischemia-induced angiogenesis in the mouse hindlimb. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005 , 102, 17202-6 | 11.5 | 62 |
|----|---|------|-----|
| 26 | Regulation of inducible nitric oxide synthase expression in advanced glycation end product-stimulated raw 264.7 cells: the role of heme oxygenase-1 and endogenous nitric oxide. <i>Diabetes</i> , 2004 , 53, 1841-50 | 0.9 | 42 |
| 25 | Nitric oxide: inhibitory effects on endothelial cell calcium signaling, prostaglandin I2 production and nitric oxide synthase expression. <i>Cardiovascular Research</i> , 2004 , 62, 194-201 | 9.9 | 28 |
| 24 | Gene transfer of endothelial NO synthase, but not eNOS plus inducible NOS, regressed atherosclerosis in rabbits. <i>Cardiovascular Research</i> , 2004 , 61, 339-51 | 9.9 | 47 |
| 23 | Increased expression of elastolytic cysteine proteases, cathepsins S and K, in the neointima of balloon-injured rat carotid arteries. <i>American Journal of Pathology</i> , 2004 , 164, 243-51 | 5.8 | 80 |
| 22 | A new HMG-CoA reductase inhibitor, pitavastatin remarkably retards the progression of high cholesterol induced atherosclerosis in rabbits. <i>Atherosclerosis</i> , 2004 , 176, 255-63 | 3.1 | 30 |
| 21 | Estrogen-related receptor alpha 1 up-regulates endothelial nitric oxide synthase expression. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 14451-6 | 11.5 | 90 |
| 20 | 17beta-estradiol inhibits NADPH oxidase activity through the regulation of p47phox mRNA and protein expression in THP-1 cells. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2003 , 1640, 113-8 | 4.9 | 38 |
| 19 | Sarpogrelate HCl, a selective 5-HT2A antagonist, retards the progression of atherosclerosis through a novel mechanism. <i>Atherosclerosis</i> , 2003 , 168, 23-31 | 3.1 | 54 |
| 18 | The long-term effect of estriol on endothelial function and bone mineral density in octogenarian women. <i>Journal of the American Geriatrics Society</i> , 2002 , 50, 777-8 | 5.6 | 10 |
| 17 | Anti-atherosclerotic effect of beta-blocker with nitric oxide-releasing action on the severe atherosclerosis. <i>Journal of Cardiovascular Pharmacology</i> , 2002 , 39, 298-309 | 3.1 | 16 |
| 16 | Estriol retards and stabilizes atherosclerosis through an NO-mediated system. <i>Life Sciences</i> , 2002 , 71, 31-42 | 6.8 | 11 |
| 15 | Cerivastatin, a hydroxymethylglutaryl coenzyme a reductase inhibitor, improves endothelial function in elderly diabetic patients within 3 days. <i>Circulation</i> , 2001 , 104, 376-9 | 16.7 | 229 |
| 14 | A HMG-CoA reductase inhibitor possesses a potent anti-atherosclerotic effect other than serum lipid lowering effectsthe relevance of endothelial nitric oxide synthase and superoxide anion scavenging action. <i>Atherosclerosis</i> , 2001 , 155, 347-57 | 3.1 | 90 |
| 13 | Up-regulation of endothelial nitric oxide synthase through beta(2)-adrenergic receptorthe role of a beta-blocker with NO-releasing action. <i>Biochemical and Biophysical Research Communications</i> , 2001 , 280, 589-94 | 3.4 | 26 |
| 12 | Estrogen prevents destabilization of endothelial nitric oxide synthase mRNA induced by tumor necrosis factor alpha through estrogen receptor mediated system. <i>Life Sciences</i> , 2001 , 69, 1651-60 | 6.8 | 27 |
| 11 | Temporal effects of 17beta-estradiol on caveolin-1 mRNA and protein in bovine aortic endothelial cells. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2001 , 281, H1327-33 | 5.2 | 30 |
| 10 | HMG-CoA reductase inhibitor stabilizes rabbit atheroma by increasing basal NO and decreasing superoxide. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2001 , 281, H75-83 | 5.2 | 25 |

| 9 | Inhibition of inducible nitric oxide synthase gene expression by indomethacin or ibuprofen in beta-amyloid protein-stimulated J774 cells. <i>European Journal of Pharmacology</i> , 2000 , 408, 137-41 | 5.3 | 22 | |
|---|---|-----|----|--|
| 8 | Physiological concentration of 17beta-estradiol retards the progression of severe atherosclerosis induced by a high-cholesterol diet plus balloon catheter injury: role of NO. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2000 , 20, 1613-21 | 9.4 | 37 | |
| 7 | Dehydroepiandrosterone retards atherosclerosis formation through its conversion to estrogen: the possible role of nitric oxide. <i>Arteriosclerosis, Thrombosis, and Vascular Biology,</i> 2000 , 20, 782-92 | 9.4 | 90 | |
| 6 | Expression of inducible nitric oxide synthase and Fas/Fas ligand correlates with the incidence of apoptotic cell death in atheromatous plaques of human coronary arteries. <i>Nitric Oxide - Biology and Chemistry</i> , 2000 , 4, 561-71 | 5 | 35 | |
| 5 | Endothelium-dependent relaxation of rabbit atherosclerotic aorta was not restored by control of hyperlipidemia: the possible role of peroxynitrite (ONOO(-)). <i>Atherosclerosis</i> , 1999 , 147, 349-63 | 3.1 | 27 | |
| 4 | A HMG-CoA reductase inhibitor improved regression of atherosclerosis in the rabbit aorta without affecting serum lipid levels: possible relevance of up-regulation of endothelial NO synthase mRNA. <i>Biochemical and Biophysical Research Communications</i> , 1999 , 259, 414-9 | 3.4 | 68 | |
| 3 | Overexpression of phospholipid hydroperoxide glutathione peroxidase suppressed cell death due to oxidative damage in rat basophile leukemia cells (RBL-2H3). <i>Biochemical and Biophysical Research Communications</i> , 1996 , 222, 432-8 | 3.4 | 65 | |
| 2 | Import into mitochondria of phospholipid hydroperoxide glutathione peroxidase requires a leader sequence. <i>Biochemical and Biophysical Research Communications</i> , 1996 , 227, 433-9 | 3.4 | 95 | |
| 1 | Gender Differences in the Risk of Metabolic Syndrome Among Chronic Arsenic-Exposed Individuals in Bangladesh. <i>Exposure and Health</i> ,1 | 8.8 | 1 | |