

Wen-Bin Guo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3173990/publications.pdf>

Version: 2024-02-01

163
papers

6,364
citations

71102

41
h-index

88630

70
g-index

164
all docs

164
docs citations

164
times ranked

5513
citing authors

#	ARTICLE	IF	CITATIONS
1	Impaired robust interhemispheric function integration of depressive brain from REST-meta-MDD database in China. <i>Bipolar Disorders</i> , 2022, 24, 400-411.	1.9	8
2	Abnormal global-brain functional connectivity and its relationship with cognitive deficits in drug-naive first-episode adolescent-onset schizophrenia. <i>Brain Imaging and Behavior</i> , 2022, , 1.	2.1	7
3	Reduced erythrocyte membrane polyunsaturated fatty acid levels indicate diminished treatment response in patients with multi- versus first-episode schizophrenia. <i>NPJ Schizophrenia</i> , 2022, 8, 7.	3.6	17
4	Disrupted Cerebellar-Default Mode Network Functional Connectivity in Major Depressive Disorder With Gastrointestinal Symptoms. <i>Frontiers in Cellular Neuroscience</i> , 2022, 16, 833592.	3.7	5
5	Epidemiological, Radiographical, and Laboratorial Characteristics of Chinese Asymptomatic Cases With COVID-19: A Systematic Review and Meta-Analysis. <i>Frontiers in Public Health</i> , 2022, 10, 808471.	2.7	2
6	Abnormal Default Mode Network Homogeneity in Major Depressive Disorder With Gastrointestinal Symptoms at Rest. <i>Frontiers in Aging Neuroscience</i> , 2022, 14, 804621.	3.4	5
7	Editorial: Brain and Somatization Symptoms in Psychiatric Disorders, Volume II. <i>Frontiers in Psychiatry</i> , 2022, 13, 881245.	2.6	0
8	Abnormal spontaneous neural activity as a potential predictor of early treatment response in patients with obsessive-compulsive disorder. <i>Journal of Affective Disorders</i> , 2022, 309, 27-36.	4.1	18
9	Reduced nucleus accumbens functional connectivity in reward network and default mode network in patients with recurrent major depressive disorder. <i>Translational Psychiatry</i> , 2022, 12, .	4.8	20
10	Altered Regional Activity and Network Homogeneity within the Fronto-Limbic Network at Rest in Medicine-Free Obsessive-Compulsive Disorder. <i>Brain Sciences</i> , 2022, 12, 857.	2.3	0
11	Less reduced gray matter volume in the subregions of superior temporal gyrus predicts better treatment efficacy in drug-naive, first-episode schizophrenia. <i>Brain Imaging and Behavior</i> , 2021, 15, 1997-2004.	2.1	6
12	Increased regional homogeneity modulated by metacognitive training predicts therapeutic efficacy in patients with schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021, 271, 783-798.	3.2	11
13	Frequency-specific alterations of the frontal-cerebellar circuit in first-episode, drug-naive somatization disorder. <i>Journal of Affective Disorders</i> , 2021, 280, 319-325.	4.1	1
14	Voxel-Wise Brain-Wide Functional Connectivity Abnormalities in Patients with Primary Blepharospasm at Rest. <i>Neural Plasticity</i> , 2021, 2021, 1-9.	2.2	10
15	Disrupted Regional Homogeneity in Melancholic and Non-melancholic Major Depressive Disorder at Rest. <i>Frontiers in Psychiatry</i> , 2021, 12, 618805.	2.6	19
16	Editorial: Emotional Disturbance and Brain Imaging in Neuropsychiatric Disorders. <i>Frontiers in Psychiatry</i> , 2021, 12, 632244.	2.6	0
17	Adverse Psychological Reactions and Psychological Aids for Medical Staff During the COVID-19 Outbreak in China. <i>Frontiers in Psychiatry</i> , 2021, 12, 580067.	2.6	9
18	Disrupted hemispheric connectivity specialization in patients with major depressive disorder: Evidence from the REST-meta-MDD Project. <i>Journal of Affective Disorders</i> , 2021, 284, 217-228.	4.1	23

#	ARTICLE	IF	CITATIONS
19	Abnormal Default-Mode Network Homogeneity in Melancholic and Nonmelancholic Major Depressive Disorder at Rest. <i>Neural Plasticity</i> , 2021, 2021, 1-12.	2.2	14
20	Shared and distinct homotopic connectivity changes in melancholic and non-melancholic depression. <i>Journal of Affective Disorders</i> , 2021, 287, 268-275.	4.1	37
21	Associations Between Internet Addiction and Gender, Anxiety, Coping Styles and Acceptance in University Freshmen in South China. <i>Frontiers in Psychiatry</i> , 2021, 12, 558080.	2.6	19
22	Disrupted Regional Homogeneity in Major Depressive Disorder With Gastrointestinal Symptoms at Rest. <i>Frontiers in Psychiatry</i> , 2021, 12, 636820.	2.6	13
23	Decreased Nucleus Accumbens Connectivity at Rest in Medication-Free Patients with Obsessive-Compulsive Disorder. <i>Neural Plasticity</i> , 2021, 2021, 1-7.	2.2	5
24	Disrupted Asymmetry of Inter- and Intra-Hemispheric Functional Connectivity at Rest in Medication-Free Obsessive-Compulsive Disorder. <i>Frontiers in Neuroscience</i> , 2021, 15, 634557.	2.8	4
25	Increased Global-Brain Functional Connectivity Is Associated with Dyslipidemia and Cognitive Impairment in First-Episode, Drug-Naive Patients with Bipolar Disorder. <i>Neural Plasticity</i> , 2021, 2021, 1-10.	2.2	4
26	Increased subcortical region volume induced by electroconvulsive therapy in patients with schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021, 271, 1285-1295.	3.2	3
27	Dissociation Pattern in Default-Mode Network Homogeneity in Drug-Naive Bipolar Disorder. <i>Frontiers in Psychiatry</i> , 2021, 12, 699292.	2.6	0
28	Disrupted intrinsic functional brain topology in patients with major depressive disorder. <i>Molecular Psychiatry</i> , 2021, 26, 7363-7371.	7.9	82
29	Increased Homotopic Connectivity in the Prefrontal Cortex Modulated by Olanzapine Predicts Therapeutic Efficacy in Patients with Schizophrenia. <i>Neural Plasticity</i> , 2021, 2021, 1-11.	2.2	8
30	Clubhouse Model of Psychiatric Rehabilitation in China to Promote Recovery of People With Schizophrenia: A Systematic Review and Meta-Analysis. <i>Frontiers in Psychiatry</i> , 2021, 12, 730552.	2.6	4
31	Altered Functional Connectivity Strength at Rest in Medication-Free Obsessive-Compulsive Disorder. <i>Neural Plasticity</i> , 2021, 2021, 1-9.	2.2	6
32	Efficacy of Treatments Targeting Hypothalamic-Pituitary-Adrenal Systems for Major Depressive Disorder: A Meta-Analysis. <i>Frontiers in Pharmacology</i> , 2021, 12, 732157.	3.5	11
33	Brain structural alterations in MDD patients with gastrointestinal symptoms: Evidence from the REST-meta-MDD project. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021, 111, 110386.	4.8	18
34	Suicidality in patients with premenstrual dysphoric disorder—A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2021, 295, 339-346.	4.1	15
35	A Selective Review of the Excitatory-Inhibitory Imbalance in Schizophrenia: Underlying Biology, Genetics, Microcircuits, and Symptoms. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 664535.	3.7	30
36	Altered Fractional Amplitude of Low-Frequency Fluctuation in Major Depressive Disorder and Bipolar Disorder. <i>Frontiers in Psychiatry</i> , 2021, 12, 739210.	2.6	8

#	ARTICLE	IF	CITATIONS
37	Abnormal interhemispheric homotopic functional connectivity is correlated with gastrointestinal symptoms in patients with major depressive disorder. <i>Journal of Psychiatric Research</i> , 2021, 144, 234-240.	3.1	8
38	Comparing Brain Functional Activities in Patients With Blepharospasm and Dry Eye Disease Measured With Resting-State fMRI. <i>Frontiers in Neurology</i> , 2021, 12, 607476.	2.4	1
39	Abnormal Network Homogeneity in the Right Superior Medial Frontal Gyrus in Cervical Dystonia. <i>Frontiers in Neurology</i> , 2021, 12, 729068.	2.4	3
40	Shared and Distinct Fractional Amplitude of Low-Frequency Fluctuation Patterns in Major Depressive Disorders With and Without Gastrointestinal Symptoms. <i>Frontiers in Psychiatry</i> , 2021, 12, 744898.	2.6	6
41	Differentiating Melancholic and Non-melancholic Major Depressive Disorder Using Fractional Amplitude of Low-Frequency Fluctuations. <i>Frontiers in Psychiatry</i> , 2021, 12, 763770.	2.6	4
42	Global Functional Connectivity Analysis Indicating Dysconnectivity of the Hate Circuit in Major Depressive Disorder. <i>Frontiers in Aging Neuroscience</i> , 2021, 13, 803080.	3.4	4
43	Altered Brain Functional Asymmetry in Patients With Major Depressive Disorder Related to Gastrointestinal Symptoms. <i>Frontiers in Neuroscience</i> , 2021, 15, 797598.	2.8	3
44	Editorial: Neurobiological Biomarkers for Developing Novel Treatments of Substance and Non-substance Addiction. <i>Frontiers in Psychiatry</i> , 2021, 12, 811032.	2.6	1
45	Resting-state functional hypoconnectivity of amygdala in clinical high risk state and first-episode schizophrenia. <i>Brain Imaging and Behavior</i> , 2020, 14, 1840-1849.	2.1	9
46	Reduced connectivity in anterior cingulate cortex as an early predictor for treatment response in drug-naive, first-episode schizophrenia: A global-brain functional connectivity analysis. <i>Schizophrenia Research</i> , 2020, 215, 337-343.	2.0	24
47	Increased cerebellar "default-mode network connectivity at rest in obsessive-compulsive disorder. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2020, 270, 1015-1024.	3.2	9
48	Region-specific insular volumetric decreases in drug-naive, first-episode schizophrenia and their unaffected siblings. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2020, 183, 106-112.	1.7	7
49	Altered resting-state dynamic functional brain networks in major depressive disorder: Findings from the REST-meta-MDD consortium. <i>NeuroImage: Clinical</i> , 2020, 26, 102163.	2.7	76
50	Abnormal default-mode network homogeneity and its correlations with neurocognitive deficits in drug-naive first-episode adolescent-onset schizophrenia. <i>Schizophrenia Research</i> , 2020, 215, 140-147.	2.0	17
51	Enhanced Connectivity of Thalamo-Cortical Networks in First-Episode, Treatment-Naive Somatization Disorder. <i>Frontiers in Psychiatry</i> , 2020, 11, 555836.	2.6	4
52	Reduced regional homogeneity and neurocognitive impairment in patients with moderate-to-severe obstructive sleep apnea. <i>Sleep Medicine</i> , 2020, 75, 418-427.	1.6	14
53	Abnormal Spontaneous Brain Activities of Limbic-Cortical Circuits in Patients With Dry Eye Disease. <i>Frontiers in Human Neuroscience</i> , 2020, 14, 574758.	2.0	9
54	Biotypes of major depressive disorder: Neuroimaging evidence from resting-state default mode network patterns. <i>NeuroImage: Clinical</i> , 2020, 28, 102514.	2.7	51

#	ARTICLE	IF	CITATIONS
55	Mental Health of Pregnant and Postpartum Women During the Coronavirus Disease 2019 Pandemic: A Systematic Review and Meta-Analysis. <i>Frontiers in Psychology</i> , 2020, 11, 617001.	2.1	164
56	Abnormal Functional Asymmetry in the Salience and Auditory Networks in First-episode, Drug-naive Somatization Disorder. <i>Neuroscience</i> , 2020, 444, 1-8.	2.3	6
57	Reduced Global-Brain Functional Connectivity of the Cerebello-Thalamo-Cortical Network in Patients With Dry Eye Disease. <i>Frontiers in Human Neuroscience</i> , 2020, 14, 572693.	2.0	8
58	Decreased Resting-State Interhemispheric Functional Connectivity in Medication-Free Obsessive-Compulsive Disorder. <i>Frontiers in Psychiatry</i> , 2020, 11, 559729.	2.6	17
59	Disrupted Regional Homogeneity in Drug-Naive Patients With Bipolar Disorder. <i>Frontiers in Psychiatry</i> , 2020, 11, 825.	2.6	13
60	Chinese Taoist Cognitive Therapy for Symptoms of Depression and Anxiety in Adults in China: A Systematic Review and Meta-Analysis. <i>Frontiers in Psychology</i> , 2020, 11, 769.	2.1	5
61	Altered Global Brain Functional Connectivity in Drug-Naive Patients With Obsessive-Compulsive Disorder. <i>Frontiers in Psychiatry</i> , 2020, 11, 98.	2.6	28
62	Clinical Characteristics of Children With COVID-19: A Meta-Analysis. <i>Frontiers in Pediatrics</i> , 2020, 8, 431.	1.9	108
63	Editorial: Dynamic Functional Connectivity in Neuropsychiatric Disorders: Methods and Applications. <i>Frontiers in Neuroscience</i> , 2020, 14, 332.	2.8	4
64	Metacognitive Training Modulates Default-Mode Network Homogeneity During 8-Week Olanzapine Treatment in Patients With Schizophrenia. <i>Frontiers in Psychiatry</i> , 2020, 11, 234.	2.6	9
65	Enhanced baseline activity in the left ventromedial putamen predicts individual treatment response in drug-naive, first-episode schizophrenia: Results from two independent study samples. <i>EBioMedicine</i> , 2019, 46, 248-255.	6.1	24
66	Reduced Brain Activity in the Right Putamen as an Early Predictor for Treatment Response in Drug-Naive, First-Episode Schizophrenia. <i>Frontiers in Psychiatry</i> , 2019, 10, 741.	2.6	20
67	Increased Nucleus Accumbens Connectivity in Resting-State Patients With Drug-Naive, First-Episode Somatization Disorder. <i>Frontiers in Psychiatry</i> , 2019, 10, 585.	2.6	9
68	Preliminary Clinical Investigation of Combinatorial Pharmacogenomic Testing for the Optimized Treatment of Depression: A Randomized Single-Blind Study. <i>Frontiers in Neuroscience</i> , 2019, 13, 960.	2.8	15
69	Cognitive deficits in subjects at risk for psychosis, first-episode and chronic schizophrenia patients. <i>Psychiatry Research</i> , 2019, 274, 235-242.	3.3	20
70	Voxel-based global-brain functional connectivity alterations in first-episode drug-naive patients with somatization disorder. <i>Journal of Affective Disorders</i> , 2019, 254, 82-89.	4.1	20
71	Abnormal spontaneous neural activity of brain regions in patients with primary blepharospasm at rest. <i>Journal of the Neurological Sciences</i> , 2019, 403, 44-49.	0.6	14
72	Dysfunction in Serotonergic and Noradrenergic Systems and Somatic Symptoms in Psychiatric Disorders. <i>Frontiers in Psychiatry</i> , 2019, 10, 286.	2.6	43

#	ARTICLE	IF	CITATIONS
73	Increased frontal gray matter volume in individuals with prodromal psychosis. <i>CNS Neuroscience and Therapeutics</i> , 2019, 25, 987-994.	3.9	16
74	Brain structural abnormalities as potential markers for detecting individuals with ultra-high risk for psychosis: A systematic review and meta-analysis. <i>Schizophrenia Research</i> , 2019, 209, 22-31.	2.0	49
75	Alterations of Interhemispheric Functional Connectivity and Degree Centrality in Cervical Dystonia: A Resting-State fMRI Study. <i>Neural Plasticity</i> , 2019, 2019, 1-11.	2.2	29
76	Enhanced Global-Brain Functional Connectivity in the Left Superior Frontal Gyrus as a Possible Endophenotype for Schizophrenia. <i>Frontiers in Neuroscience</i> , 2019, 13, 145.	2.8	29
77	Editorial: Brain and Somatization Symptoms in Psychiatric Disorders. <i>Frontiers in Psychiatry</i> , 2019, 10, 146.	2.6	10
78	Altered network homogeneity of the default-mode network in drug-naive obsessive-compulsive disorder. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2019, 93, 77-83.	4.8	28
79	Reduced default mode network functional connectivity in patients with recurrent major depressive disorder. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 9078-9083.	7.1	441
80	Functional asymmetry of thalamocortical networks in subjects at ultra-high risk for psychosis and first-episode schizophrenia. <i>European Neuropsychopharmacology</i> , 2019, 29, 519-528.	0.7	24
81	Cerebellar structural and functional abnormalities in first-episode and drug-naïve patients with schizophrenia: A meta-analysis. <i>Psychiatry Research - Neuroimaging</i> , 2019, 283, 24-33.	1.8	46
82	Reduced Global-Brain Functional Connectivity and Its Relationship With Symptomatic Severity in Cervical Dystonia. <i>Frontiers in Neurology</i> , 2019, 10, 1358.	2.4	18
83	Cerebellar abnormalities in first-episode, drug-naïve schizophrenia at rest. <i>Psychiatry Research - Neuroimaging</i> , 2018, 276, 73-79.	1.8	29
84	Increased coherence-based regional homogeneity in resting-state patients with first-episode, drug-naïve somatization disorder. <i>Journal of Affective Disorders</i> , 2018, 235, 150-154.	4.1	12
85	Voxel-wise brain-wide functional connectivity abnormalities in first-episode, drug-naïve patients with major depressive disorder. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2018, 177, 447-453.	1.7	25
86	Altered functional connectivity strength and its correlations with cognitive function in subjects with ultra-high risk for psychosis at rest. <i>CNS Neuroscience and Therapeutics</i> , 2018, 24, 1140-1148.	3.9	21
87	Decreased Resting-State Interhemispheric Functional Connectivity Correlated with Neurocognitive Deficits in Drug-Naïve First-Episode Adolescent-Onset Schizophrenia. <i>International Journal of Neuropsychopharmacology</i> , 2018, 21, 33-41.	2.1	43
88	Decreased white matter FA values in the left inferior frontal gyrus is a possible intermediate phenotype of schizophrenia: evidences from a novel group strategy. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2018, 268, 89-98.	3.2	10
89	Abnormal neural activity as a potential biomarker for drug-naïve first-episode adolescent-onset schizophrenia with coherence regional homogeneity and support vector machine analyses. <i>Schizophrenia Research</i> , 2018, 192, 408-415.	2.0	52
90	Abnormal regional homogeneity as a potential imaging biomarker for adolescent-onset schizophrenia: A resting-state fMRI study and support vector machine analysis. <i>Schizophrenia Research</i> , 2018, 192, 179-184.	2.0	80

#	ARTICLE	IF	CITATIONS
91	Treatment effects of olanzapine on homotopic connectivity in drug-free schizophrenia at rest. <i>World Journal of Biological Psychiatry</i> , 2018, 19, S106-S114.	2.6	31
92	Increased anterior default-mode network homogeneity in first-episode, drug-naive major depressive disorder: A replication study. <i>Journal of Affective Disorders</i> , 2018, 225, 767-772.	4.1	19
93	Decreased interhemispheric coordination in the posterior default-mode network and visual regions as trait alterations in first-episode, drug-naive major depressive disorder. <i>Brain Imaging and Behavior</i> , 2018, 12, 1251-1258.	2.1	23
94	Abnormal long- and short-range functional connectivity in adolescent-onset schizophrenia patients: A resting-state fMRI study. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2018, 81, 445-451.	4.8	55
95	Emotional Roles of Mono-Aminergic Neurotransmitters in Major Depressive Disorder and Anxiety Disorders. <i>Frontiers in Psychology</i> , 2018, 9, 2201.	2.1	126
96	Disrupted asymmetry of inter- and intra-hemispheric functional connectivity in patients with drug-naive, first-episode schizophrenia and their unaffected siblings. <i>EBioMedicine</i> , 2018, 36, 429-435.	6.1	32
97	Voxel-Mirrored Homotopic Connectivity of Resting-State Functional Magnetic Resonance Imaging in Blepharospasm. <i>Frontiers in Psychology</i> , 2018, 9, 1620.	2.1	21
98	Regional white matter volume abnormalities in first-episode somatization disorder. <i>International Journal of Psychophysiology</i> , 2018, 133, 12-16.	1.0	6
99	Bidirectional Causal Connectivity in the Cortico-Limbic-Cerebellar Circuit Related to Structural Alterations in First-Episode, Drug-Naive Somatization Disorder. <i>Frontiers in Psychiatry</i> , 2018, 9, 162.	2.6	16
100	Altered Serum Tumor Necrosis Factor and Interleukin-1 β in First-Episode Drug-Naive and Chronic Schizophrenia. <i>Frontiers in Neuroscience</i> , 2018, 12, 296.	2.8	28
101	Abnormal functional connectivity strength in patients with adolescent-onset schizophrenia: a resting-state fMRI study. <i>European Child and Adolescent Psychiatry</i> , 2017, 26, 839-845.	4.7	25
102	Anatomical distance affects cortical-subcortical connectivity in first-episode, drug-naive somatization disorder. <i>Journal of Affective Disorders</i> , 2017, 217, 153-158.	4.1	11
103	Using short-range and long-range functional connectivity to identify schizophrenia with a family-based case-control design. <i>Psychiatry Research - Neuroimaging</i> , 2017, 264, 60-67.	1.8	18
104	Olanzapine modulates the default-mode network homogeneity in recurrent drug-free schizophrenia at rest. <i>Australian and New Zealand Journal of Psychiatry</i> , 2017, 51, 1000-1009.	2.3	21
105	Hyperactivity of the default-mode network in first-episode, drug-naive schizophrenia at rest revealed by family-based case-control and traditional case-control designs. <i>Medicine (United States)</i> , 2017, 96, e6223.	1.0	40
106	Family-based case-control study of homotopic connectivity in first-episode, drug-naive schizophrenia at rest. <i>Scientific Reports</i> , 2017, 7, 43312.	3.3	22
107	Aberrant default mode network homogeneity in patients with first-episode treatment-naive melancholic depression. <i>International Journal of Psychophysiology</i> , 2017, 112, 46-51.	1.0	26
108	Olanzapine modulation of long- and short-range functional connectivity in the resting brain in a sample of patients with schizophrenia. <i>European Neuropsychopharmacology</i> , 2017, 27, 48-58.	0.7	37

#	ARTICLE	IF	CITATIONS
109	Relationship between long-term use of a typical antipsychotic medication by Chinese schizophrenia patients and the bone turnover markers serum osteocalcin and $\hat{1}^2$ -CrossLaps. <i>Schizophrenia Research</i> , 2016, 176, 259-263.	2.0	21
110	Abnormal regional homogeneity as potential imaging biomarker for psychosis risk syndrome: a resting-state fMRI study and support vector machine analysis. <i>Scientific Reports</i> , 2016, 6, 27619.	3.3	25
111	Patients with first-episode, drug-naïve schizophrenia and subjects at ultra-high risk of psychosis shared increased cerebellar-default mode network connectivity at rest. <i>Scientific Reports</i> , 2016, 6, 26124.	3.3	46
112	Clinical significance of increased cerebellar default-mode network connectivity in resting-state patients with drug-naïve somatization disorder. <i>Medicine (United States)</i> , 2016, 95, e4043.	1.0	22
113	Differences among first-episode schizophrenia patients, healthy siblings, and controls at the individual level. <i>International Journal of Psychophysiology</i> , 2016, 104, 24-32.	1.0	3
114	Decreased interhemispheric functional connectivity in insula and angular gyrus/supramarginal gyrus: Significant findings in first-episode, drug-naïve somatization disorder. <i>Psychiatry Research - Neuroimaging</i> , 2016, 248, 48-54.	1.8	37
115	Abnormal default-mode network homogeneity and its correlations with personality in drug-naïve somatization disorder at rest. <i>Journal of Affective Disorders</i> , 2016, 193, 81-88.	4.1	31
116	Decreased long- and short-range functional connectivity at rest in drug-naïve major depressive disorder. <i>Australian and New Zealand Journal of Psychiatry</i> , 2016, 50, 763-769.	2.3	21
117	Resting-state cerebellar-cerebral networks are differently affected in first-episode, drug-naïve schizophrenia patients and unaffected siblings. <i>Scientific Reports</i> , 2015, 5, 17275.	3.3	45
118	Increased Cerebellar Functional Connectivity With the Default-Mode Network in Unaffected Siblings of Schizophrenia Patients at Rest. <i>Schizophrenia Bulletin</i> , 2015, 41, 1317-1325.	4.3	48
119	Dissociation of anatomical and functional alterations of the default-mode network in first-episode, drug-naïve schizophrenia. <i>Clinical Neurophysiology</i> , 2015, 126, 2276-2281.	1.5	10
120	Increased Causal Connectivity Related to Anatomical Alterations as Potential Endophenotypes for Schizophrenia. <i>Medicine (United States)</i> , 2015, 94, e1493.	1.0	11
121	Unidirectionally affected causal connectivity of cortico-limbic-cerebellar circuit by structural deficits in drug-naïve major depressive disorder. <i>Journal of Affective Disorders</i> , 2015, 172, 410-416.	4.1	31
122	Disrupted cortical hubs in functional brain networks in social anxiety disorder. <i>Clinical Neurophysiology</i> , 2015, 126, 1711-1716.	1.5	126
123	Abnormal regional homogeneity and its correlations with personality in first-episode, treatment-naïve somatization disorder. <i>International Journal of Psychophysiology</i> , 2015, 97, 108-112.	1.0	28
124	Dissociation of functional and anatomical brain abnormalities in unaffected siblings of schizophrenia patients. <i>Clinical Neurophysiology</i> , 2015, 126, 927-932.	1.5	28
125	Decreased insular connectivity in drug-naïve major depressive disorder at rest. <i>Journal of Affective Disorders</i> , 2015, 179, 31-37.	4.1	65
126	Regional white matter abnormalities in drug-naïve, first-episode schizophrenia patients and their healthy unaffected siblings. <i>Australian and New Zealand Journal of Psychiatry</i> , 2015, 49, 246-254.	2.3	25

#	ARTICLE	IF	CITATIONS
127	Decreased regional activity and network homogeneity of the fronto-limbic network at rest in drug-naive major depressive disorder. Australian and New Zealand Journal of Psychiatry, 2015, 49, 550-556.	2.3	34
128	Increased Cerebellar-Default-Mode-Network Connectivity in Drug-Naive Major Depressive Disorder at Rest. Medicine (United States), 2015, 94, e560.	1.0	54
129	Dysfunctional resting-state connectivities of brain regions with structural deficits in drug-naive first-episode schizophrenia adolescents. Schizophrenia Research, 2015, 168, 353-359.	2.0	31
130	Increased short-range and long-range functional connectivity in first-episode, medication-naive schizophrenia at rest. Schizophrenia Research, 2015, 166, 144-150.	2.0	48
131	Alterations in white matter integrity in first-episode, treatment-naive patients with somatization disorder. Neuroscience Letters, 2015, 599, 102-108.	2.1	17
132	Increased functional connectivity strength of right inferior temporal gyrus in first-episode, drug-naive somatization disorder. Australian and New Zealand Journal of Psychiatry, 2015, 49, 74-81.	2.3	59
133	Abnormal Causal Connectivity by Structural Deficits in First-Episode, Drug-Naive Schizophrenia at Rest. Schizophrenia Bulletin, 2015, 41, 57-65.	4.3	71
134	Multivariate classification of social anxiety disorder using whole brain functional connectivity. Brain Structure and Function, 2015, 220, 101-115.	2.3	321
135	Abnormal Default-Mode Network Homogeneity in First-Episode, Drug-Naive Major Depressive Disorder. PLoS ONE, 2014, 9, e91102.	2.5	72
136	Dissociation of Regional Activity in Default Mode Network in Medication-Naive, First-Episode Somatization Disorder. PLoS ONE, 2014, 9, e99273.	2.5	45
137	Decreased default-mode network homogeneity in unaffected siblings of schizophrenia patients at rest. Psychiatry Research - Neuroimaging, 2014, 224, 218-224.	1.8	23
138	Disrupted white matter integrity in first-episode, drug-naive, late-onset depression. Journal of Affective Disorders, 2014, 163, 70-75.	4.1	26
139	Abnormal default-mode network homogeneity in first-episode, drug-naive schizophrenia at rest. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2014, 49, 16-20.	4.8	94
140	Decreased gray matter volume in the left middle temporal gyrus as a candidate biomarker for schizophrenia: A study of drug naive, first-episode schizophrenia patients and unaffected siblings. Schizophrenia Research, 2014, 159, 43-50.	2.0	43
141	Decreased resting-state interhemispheric functional connectivity in unaffected siblings of schizophrenia patients. Schizophrenia Research, 2014, 152, 170-175.	2.0	50
142	Decreased resting-state interhemispheric coordination in first-episode, drug-naive paranoid schizophrenia. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2014, 48, 14-19.	4.8	76
143	Functional and anatomical brain deficits in drug-naive major depressive disorder. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2014, 54, 1-6.	4.8	61
144	Decreased regional activity of default-mode network in unaffected siblings of schizophrenia patients at rest. European Neuropsychopharmacology, 2014, 24, 545-552.	0.7	31

#	ARTICLE	IF	CITATIONS
145	Dissociation of regional activity in the default mode network in first-episode, drug-naive major depressive disorder at rest. <i>Journal of Affective Disorders</i> , 2013, 151, 1097-1101.	4.1	49
146	Is there a cerebellar compensatory effort in first-episode, treatment-naive major depressive disorder at rest?. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2013, 46, 13-18.	4.8	82
147	Abnormal amplitude low-frequency oscillations in medication-naive, first-episode patients with major depressive disorder: A resting-state fMRI study. <i>Journal of Affective Disorders</i> , 2013, 146, 401-406.	4.1	231
148	Abnormal resting-state cerebellar-cerebral functional connectivity in treatment-resistant depression and treatment sensitive depression. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2013, 44, 51-57.	4.8	160
149	Decreased interhemispheric resting-state functional connectivity in first-episode, drug-naive major depressive disorder. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2013, 41, 24-29.	4.8	121
150	Reversal alterations of amplitude of low-frequency fluctuations in early and late onset, first-episode, drug-naive depression. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2013, 40, 153-159.	4.8	58
151	Decreased Interhemispheric Coordination in Treatment-Resistant Depression: A Resting-State fMRI Study. <i>PLoS ONE</i> , 2013, 8, e71368.	2.5	68
152	Abnormal neural activity of brain regions in treatment-resistant and treatment-sensitive major depressive disorder: A resting-state fMRI study. <i>Journal of Psychiatric Research</i> , 2012, 46, 1366-1373.	3.1	96
153	Right lateralized white matter abnormalities in first-episode, drug-naive paranoid schizophrenia. <i>Neuroscience Letters</i> , 2012, 531, 5-9.	2.1	85
154	Alterations of the amplitude of low-frequency fluctuations in treatment-resistant and treatment-response depression: A resting-state fMRI study. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2012, 37, 153-160.	4.8	148
155	Altered white matter integrity of forebrain in treatment-resistant depression: A diffusion tensor imaging study with tract-based spatial statistics. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2012, 38, 201-206.	4.8	71
156	Altered white matter integrity in young adults with first-episode, treatment-naive, and treatment-responsive depression. <i>Neuroscience Letters</i> , 2012, 522, 139-144.	2.1	78
157	Early and late onset, first-episode, treatment-naive depression: same clinical symptoms, different regional neural activities. <i>Journal of Affective Disorders</i> , 2012, 143, 56-63.	4.1	80
158	Abnormal regional spontaneous neural activity in first-episode, treatment-naive patients with late-life depression: A resting-state fMRI study. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2012, 39, 326-331.	4.8	155
159	Classification of Different Therapeutic Responses of Major Depressive Disorder with Multivariate Pattern Analysis Method Based on Structural MR Scans. <i>PLoS ONE</i> , 2012, 7, e40968.	2.5	125
160	Disrupted regional homogeneity in treatment-resistant depression: A resting-state fMRI study. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2011, 35, 1297-1302.	4.8	110
161	Abnormal neural activities in first-episode, treatment-naive, short-illness-duration, and treatment-response patients with major depressive disorder: A resting-state fMRI study. <i>Journal of Affective Disorders</i> , 2011, 135, 326-331.	4.1	127
162	Difference ERPs Effects of the Difference Introduction on the Recognition of Chinese Emotional Content Words in Healthy Subjects. , 0, , .		1

#	ARTICLE	IF	CITATIONS
163	Multiple-Network Alterations in Major Depressive Disorder With Gastrointestinal Symptoms at Rest Revealed by Global Functional Connectivity Analysis. <i>Frontiers in Neuroscience</i> , 0, 16, .	2.8	2