

Mitsi A Blount

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3179328/publications.pdf>

Version: 2024-02-01

37
papers

1,070
citations

430874

18
h-index

414414

32
g-index

38
all docs

38
docs citations

38
times ranked

746
citing authors

#	ARTICLE	IF	CITATIONS
1	Purinergic signaling is enhanced in the absence of UT-A1 and UT-A3. <i>Physiological Reports</i> , 2021, 9, e14636.	1.7	1
2	Lack of urea transporters, UT-A1 and UT-A3, increases nitric oxide accumulation to dampen medullary sodium reabsorption through ENaC. <i>American Journal of Physiology - Renal Physiology</i> , 2019, 316, F539-F549.	2.7	2
3	Chronic lithium treatment induces novel patterns of pendrin localization and expression. <i>American Journal of Physiology - Renal Physiology</i> , 2018, 315, F313-F322.	2.7	5
4	UT (Urea Transporter). , 2018, , 5862-5872.		0
5	Metformin, an AMPK activator, stimulates the phosphorylation of aquaporin 2 and urea transporter A1 in inner medullary collecting ducts. <i>American Journal of Physiology - Renal Physiology</i> , 2016, 310, F1008-F1012.	2.7	46
6	Transgenic Restoration of Urea Transporter A1 Confers Maximal Urinary Concentration in the Absence of Urea Transporter A3. <i>Journal of the American Society of Nephrology: JASN</i> , 2016, 27, 1448-1455.	6.1	19
7	UT (Urea Transporter). , 2016, , 1-10.		0
8	Activation of protein kinase C β increases phosphorylation of the UT-A1 urea transporter at serine 494 in the inner medullary collecting duct. <i>American Journal of Physiology - Cell Physiology</i> , 2015, 309, C608-C615.	4.6	11
9	Vasopressin regulation of multisite phosphorylation of UT-A1 in the inner medullary collecting duct. <i>American Journal of Physiology - Renal Physiology</i> , 2015, 308, F49-F55.	2.7	12
10	Absence of PKC-Alpha Attenuates Lithium-Induced Nephrogenic Diabetes Insipidus. <i>PLoS ONE</i> , 2014, 9, e101753.	2.5	17
11	Novel activators of aquaporin 2 membrane expression for the treatment of nephrogenic diabetes insipidus: less is more. Focus on "High-throughput chemical screening identifies AG-490 as a stimulator of aquaporin 2 membrane expression and urine concentration". <i>American Journal of Physiology - Cell Physiology</i> , 2014, 307, C595-C596.	4.6	2
12	Genes and Proteins of Urea Transporters. <i>Sub-Cellular Biochemistry</i> , 2014, 73, 45-63.	2.4	18
13	Urine concentration in the diabetic mouse requires both urea and water transporters. <i>American Journal of Physiology - Renal Physiology</i> , 2013, 304, F103-F111.	2.7	13
14	Acute calcineurin inhibition with tacrolimus increases phosphorylated UT-A1. <i>American Journal of Physiology - Renal Physiology</i> , 2012, 302, F998-F1004.	2.7	11
15	Lack of protein kinase C β leads to impaired urine concentrating ability and decreased aquaporin-2 in angiotensin II-induced hypertension. <i>American Journal of Physiology - Renal Physiology</i> , 2012, 303, F37-F44.	2.7	19
16	The Role of Nitric Oxide in the Dysregulation of the Urine Concentration Mechanism in Diabetes Mellitus. <i>Frontiers in Physiology</i> , 2012, 3, 176.	2.8	14
17	Molecular mechanisms of urea transport in health and disease. <i>Pflügers Archiv European Journal of Physiology</i> , 2012, 464, 561-572.	2.8	40
18	The urea transporter UT-A1 is phosphorylated at serines 486 and 499 downstream of cyclic AMP production. <i>FASEB Journal</i> , 2012, 26, 885.11.	0.5	0

#	ARTICLE	IF	CITATIONS
19	Urea Transport in the Kidney. , 2011, 1, 699-729.		69
20	Metal ion stimulators of PDE5 cause similar conformational changes in the enzyme as does cGMP or sildenafil. Cellular Signalling, 2011, 23, 778-784.	3.6	4
21	Regulation of renal urea transport by vasopressin. Transactions of the American Clinical and Climatological Association, 2011, 122, 82-92.	0.5	26
22	A timely characterization of vasopressin-sensitive adenylyl cyclase isoforms in the mouse inner medullary collecting duct. American Journal of Physiology - Renal Physiology, 2010, 298, F857-F858.	2.7	2
23	Expression of transporters involved in urine concentration recovers differently after cessation of lithium treatment. American Journal of Physiology - Renal Physiology, 2010, 298, F601-F608.	2.7	28
24	Phosphorylation of UT-A1 on serine 486 correlates with membrane accumulation and urea transport activity in both rat IMCDs and cultured cells. American Journal of Physiology - Renal Physiology, 2010, 298, F935-F940.	2.7	33
25	Epac Regulates UT-A1 to Increase Urea Transport in Inner Medullary Collecting Ducts. Journal of the American Society of Nephrology: JASN, 2009, 20, 2018-2024.	6.1	48
26	Phosphorylation Increases Affinity of the Phosphodiesterase-5 Catalytic Site for Tadalafil. Journal of Pharmacology and Experimental Therapeutics, 2008, 325, 62-68.	2.5	18
27	Phosphorylation of UT-A1 urea transporter at serines 486 and 499 is important for vasopressin-regulated activity and membrane accumulation. American Journal of Physiology - Renal Physiology, 2008, 295, F295-F299.	2.7	83
28	Urea transporters UT-A1 and UT-A3 accumulate in the plasma membrane in response to increased hypertonicity. American Journal of Physiology - Renal Physiology, 2008, 295, F1336-F1341.	2.7	34
29	Candesartan augments compensatory changes in medullary transport proteins in the diabetic rat kidney. American Journal of Physiology - Renal Physiology, 2008, 294, F1448-F1452.	2.7	19
30	AVP causes transient formation of cAMP and activation of phosphodiesterase activity in MDCK cells. FASEB Journal, 2008, 22, 1216.13.	0.5	0
31	Conversion of Phosphodiesterase-5 (PDE5) Catalytic Site to Higher Affinity by PDE5 Inhibitors. Journal of Pharmacology and Experimental Therapeutics, 2007, 323, 730-737.	2.5	23
32	Forskolin stimulates phosphorylation and membrane accumulation of UT-A3. American Journal of Physiology - Renal Physiology, 2007, 293, F1308-F1313.	2.7	76
33	Phosphorylation of phosphodiesterase-5 is promoted by a conformational change induced by sildenafil, vardenafil, or tadalafil. Frontiers in Bioscience - Landmark, 2007, 12, 1899.	3.0	19
34	Vasopressin Increases Plasma Membrane Accumulation of Urea Transporter UT-A1 in Rat Inner Medullary Collecting Ducts. Journal of the American Society of Nephrology: JASN, 2006, 17, 2680-2686.	6.1	81
35	A 46-Amino Acid Segment in Phosphodiesterase-5 GAF-B Domain Provides for High Vardenafil Potency over Sildenafil and Tadalafil and Is Involved in Phosphodiesterase-5 Dimerization. Molecular Pharmacology, 2006, 70, 1822-1831.	2.3	34
36	Binding of Tritiated Sildenafil, Tadalafil, or Vardenafil to the Phosphodiesterase-5 Catalytic Site Displays Potency, Specificity, Heterogeneity, and cGMP Stimulation. Molecular Pharmacology, 2004, 66, 144-152.	2.3	168

#	ARTICLE	IF	CITATIONS
37	[3H]Sildenafil Binding to Phosphodiesterase-5 Is Specific, Kinetically Heterogeneous, and Stimulated by cGMP. <i>Molecular Pharmacology</i> , 2003, 63, 1364-1372.	2.3	75