

Eisuke Inoue

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3186524/publications.pdf>

Version: 2024-02-01

15
papers

337
citations

1163117

8
h-index

996975

15
g-index

15
all docs

15
docs citations

15
times ranked

403
citing authors

#	ARTICLE	IF	CITATIONS
1	A large observational cohort study of rheumatoid arthritis, IORRA: Providing context for today's treatment options. <i>Modern Rheumatology</i> , 2020, 30, 1-6.	1.8	83
2	Associated factors for falls and fear of falling in Japanese patients with rheumatoid arthritis. <i>Clinical Rheumatology</i> , 2009, 28, 1325-1330.	2.2	61
3	Proposal of Classification Criteria for HTLV-1-Associated Myelopathy/Tropical Spastic Paraparesis Disease Activity. <i>Frontiers in Microbiology</i> , 2018, 9, 1651.	3.5	48
4	Effectiveness of Daily Prednisolone to Slow Progression of Human T-Lymphotropic Virus Type 1-Associated Myelopathy/Tropical Spastic Paraparesis: A Multicenter Retrospective Cohort Study. <i>Neurotherapeutics</i> , 2017, 14, 1084-1094.	4.4	29
5	Risk factors for established vertebral fractures in Japanese patients with rheumatoid arthritis: Results from a large prospective observational cohort study. <i>Modern Rheumatology</i> , 2015, 25, 373-378.	1.8	27
6	Incidence of falls and fear of falling in Japanese patients with rheumatoid arthritis. <i>Modern Rheumatology</i> , 2011, 21, 51-56.	1.8	21
7	Profiles of EQ-5D utility scores in the daily practice of Japanese patients with rheumatoid arthritis; Analysis of the IORRA database. <i>Modern Rheumatology</i> , 2016, 26, 40-45.	1.8	19
8	Use of cerebrospinal fluid CXCL10 and neopterin as biomarkers in HTLV-1-associated myelopathy/tropical spastic paraparesis treated with steroids. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2020, 91, 321-323.	1.9	11
9	Vitamin D deficiency is a risk factor for new fractures in Japanese postmenopausal women with rheumatoid arthritis: results from the IORRA cohort study. <i>Archives of Osteoporosis</i> , 2021, 16, 119.	2.4	8
10	Factors associated with frailty in Japanese patients with rheumatoid arthritis: results from the Institute of Rheumatology Rheumatoid Arthritis cohort study. <i>Clinical Rheumatology</i> , 2022, 41, 405-410.	2.2	8
11	Factors associated with decreasing serum 25(OH)D among Japanese patients with rheumatoid arthritis: Results from the IORRA cohort study. <i>Modern Rheumatology</i> , 2019, 29, 430-435.	1.8	7
12	Factors associated with osteoporosis medication use in Japanese patients with rheumatoid arthritis: Results from the Institute of Rheumatology Rheumatoid Arthritis cohort study. <i>Osteoporosis and Sarcopenia</i> , 2020, 6, 82-87.	1.9	6
13	Polypharmacy, declined walking speed, bent back, and disability associated with a history of falls in Japanese patients with rheumatoid arthritis: Results from the IORRA cohort study. <i>Modern Rheumatology</i> , 2021, 31, 764-767.	1.8	5
14	A simple screening test to assess risk of falls in Japanese patients with rheumatoid arthritis: Results from the IORRA cohort study. <i>Modern Rheumatology</i> , 2021, 31, 506-509.	1.8	2
15	Adherence to denosumab for the treatment of osteoporosis in Japanese patients with rheumatoid arthritis: Results from the IORRA cohort study. <i>Modern Rheumatology</i> , 2021, 31, 510-513.	1.8	2