

Ignace Mc Janssen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/318901/publications.pdf>

Version: 2024-02-01

35
papers

1,475
citations

279487

23
h-index

360668

35
g-index

35
all docs

35
docs citations

35
times ranked

2002
citing authors

#	ARTICLE	IF	CITATIONS
1	Wait-and-see policy or laparoscopic cholecystectomy after endoscopic sphincterotomy for bile-duct stones: a randomised trial. <i>Lancet, The</i> , 2002, 360, 761-765.	6.3	309
2	Secondary surgery after sleeve gastrectomy: Roux-en-Y gastric bypass or biliopancreatic diversion with duodenal switch. <i>Surgery for Obesity and Related Diseases</i> , 2015, 11, 771-777.	1.0	138
3	Alterations of Hormonally Active Fibroblast Growth Factors after Roux-en-Y Gastric Bypass Surgery. <i>Digestive Diseases</i> , 2011, 29, 48-51.	0.8	118
4	Aspiration Therapy As a Tool to Treat Obesity: 1- to 4-Year Results in a 201-Patient Multi-Center Post-Market European Registry Study. <i>Obesity Surgery</i> , 2018, 28, 1860-1868.	1.1	63
5	Vitamin D absorption: consequences of gastric bypass surgery. <i>European Journal of Endocrinology</i> , 2011, 164, 827-832.	1.9	53
6	Revisional surgery after failed gastric banding: results of one-stage conversion to RYGB in 195 patients. <i>Surgery for Obesity and Related Diseases</i> , 2014, 10, 1077-1083.	1.0	50
7	Striatal dopamine receptor binding in morbidly obese women before and after gastric bypass surgery and its relationship with insulin sensitivity. <i>Diabetologia</i> , 2014, 57, 1078-1080.	2.9	50
8	A Longer Biliopancreatic Limb in Roux-en-Y Gastric Bypass Improves Weight Loss in the First Years After Surgery: Results of a Randomized Controlled Trial. <i>Obesity Surgery</i> , 2018, 28, 3744-3755.	1.1	50
9	Altered neural responsivity to food cues in relation to food preferences, but not appetite-related hormone concentrations after RYGB-surgery. <i>Behavioural Brain Research</i> , 2018, 353, 194-202.	1.2	42
10	Striatal dopamine D2/3 receptor availability increases after long-term bariatric surgery-induced weight loss. <i>European Neuropsychopharmacology</i> , 2016, 26, 1190-1200.	0.3	39
11	The magnitude and progress of lean body mass, fat-free mass, and skeletal muscle mass loss following bariatric surgery: A systematic review and meta-analysis. <i>Obesity Reviews</i> , 2022, 23, e13370.	3.1	39
12	What happens after gastric band removal without additional bariatric surgery?. <i>Surgery for Obesity and Related Diseases</i> , 2014, 10, 1092-1096.	1.0	36
13	Weight loss induced by very low calorie diet is associated with a more beneficial systemic inflammatory profile than by Roux-en-Y gastric bypass. <i>Metabolism: Clinical and Experimental</i> , 2016, 65, 1614-1620.	1.5	36
14	Body image dissatisfaction and depression in postbariatric patients is associated with less weight loss and a desire for body contouring surgery. <i>Surgery for Obesity and Related Diseases</i> , 2018, 14, 1507-1515.	1.0	36
15	Subclinical hypothyroidism and its relation to obesity in patients before and after Roux-en-Y gastric bypass. <i>Surgery for Obesity and Related Diseases</i> , 2015, 11, 1257-1263.	1.0	34
16	Effect of Carbohydrate Restriction in Patients with Hyperinsulinemic Hypoglycemia after Roux-en-Y Gastric Bypass. <i>Obesity Surgery</i> , 2014, 24, 1850-1855.	1.1	33
17	Safety experience with the duodenal-jejunal bypass liner: an endoscopic treatment for diabetes and obesity. <i>Gastrointestinal Endoscopy</i> , 2015, 82, 845-852.	0.5	28
18	A long biliopancreatic and short alimentary limb results in more weight loss in revisional RYGB surgery. Outcomes of the randomized controlled ELEGANCE REDO trial. <i>Surgery for Obesity and Related Diseases</i> , 2019, 15, 60-69.	1.0	28

#	ARTICLE	IF	CITATIONS
19	Decreased Peritoneal Tissue Plasminogen Activator During Prolonged Laparoscopic Surgery. <i>Journal of Surgical Research</i> , 2009, 151, 89-93.	0.8	27
20	Duodenal-jejunal bypass liner implantation provokes rapid weight loss and improved glycemic control, accompanied by elevated fasting ghrelin levels. <i>Endoscopy International Open</i> , 2014, 2, E21-E27.	0.9	27
21	An optimized multivitamin supplement lowers the number of vitamin and mineral deficiencies three years after Roux-en-Y gastric bypass: a cohort study. <i>Surgery for Obesity and Related Diseases</i> , 2016, 12, 659-667.	1.0	26
22	Gonadal status and outcome of bariatric surgery in obese men. <i>Clinical Endocrinology</i> , 2014, 81, 378-386.	1.2	23
23	Regulation of angiopoietin-like 4 and lipoprotein lipase in human adipose tissue. <i>Journal of Clinical Lipidology</i> , 2018, 12, 773-783.	0.6	23
24	Intra-gastric band erosion: Experiences with gastrointestinal endoscopic removal. <i>World Journal of Gastroenterology</i> , 2015, 21, 1567.	1.4	20
25	Laparoscopic Adjustable Gastric Banding: A Prospective Comparison of Two Commonly Used Bands. <i>Obesity Surgery</i> , 2002, 12, 579-582.	1.1	19
26	Preperitoneal Bupivacaine Infiltration Reduces Postoperative Opioid Consumption, Acute Pain, and Chronic Postsurgical Pain After Bariatric Surgery: a Randomized Controlled Trial. <i>Obesity Surgery</i> , 2018, 28, 3102-3110.	1.1	18
27	The feasibility of delivering a duodenal-jejunal bypass liner (endobarrier) endoscopically with patients under conscious sedation. <i>Surgical Endoscopy and Other Interventional Techniques</i> , 2014, 28, 325-330.	1.3	17
28	Changes in glycemic control and body weight after explantation of the duodenal-jejunal bypass liner. <i>Gastrointestinal Endoscopy</i> , 2017, 85, 409-415.	0.5	17
29	Evaluation of carbohydrate restriction as primary treatment for post-gastric bypass hypoglycemia. <i>Surgery for Obesity and Related Diseases</i> , 2017, 13, 404-410.	1.0	15
30	Management of vitamin K deficiency after biliopancreatic diversion with or without duodenal switch. <i>Surgery for Obesity and Related Diseases</i> , 2016, 12, 338-344.	1.0	14
31	Radiographic appearance of endoscopic duodenal-jejunal bypass liner for treatment of obesity and type 2 diabetes. <i>Surgery for Obesity and Related Diseases</i> , 2009, 5, 371-374.	1.0	12
32	Gastric pouch emptying of solid food in patients with successful and unsuccessful weight loss after Roux-en-Y gastric bypass surgery. <i>Surgery for Obesity and Related Diseases</i> , 2017, 13, 1840-1846.	1.0	12
33	Patients' Preoperative Estimate of Target Weight and Actual Outcome after Bariatric Surgery. <i>Obesity Surgery</i> , 2017, 27, 1729-1734.	1.1	10
34	A randomized controlled trial comparing oral and intravenous iron supplementation after Roux-en-Y gastric bypass surgery. <i>Clinical Nutrition</i> , 2020, 39, 3779-3785.	2.3	7
35	Leakage of Adjustable Gastric Bands. <i>Obesity Surgery</i> , 1999, 9, 258-260.	1.1	6