Tun-Jun Tsai

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3193378/publications.pdf

Version: 2024-02-01

91872 76322 5,564 130 40 69 citations h-index g-index papers 130 130 130 6019 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Effect of Pentoxifylline in Addition to Losartan on Proteinuria and GFR in CKD: A 12-Month Randomized Trial. American Journal of Kidney Diseases, 2008, 52, 464-474.	1.9	325
2	Renoprotective effect of combining pentoxifylline with angiotensin-converting enzyme inhibitor or angiotensin II receptor blocker in advanced chronic kidney disease. Journal of the Formosan Medical Association, 2014, 113, 219-226.	1.7	283
3	Platelet-derived growth factor receptor signaling activates pericyte–myofibroblast transition in obstructive and post-ischemic kidney fibrosis. Kidney International, 2011, 80, 1170-1181.	5.2	273
4	Targeting Endothelium-Pericyte Cross Talk by Inhibiting VEGF Receptor Signaling Attenuates Kidney Microvascular Rarefaction and Fibrosis. American Journal of Pathology, 2011, 178, 911-923.	3.8	224
5	Transforming Growth Factor \hat{l}^2 -1 Stimulates Profibrotic Epithelial Signaling to Activate Pericyte-Myofibroblast Transition in Obstructive Kidney Fibrosis. American Journal of Pathology, 2013, 182, 118-131.	3.8	206
6	Acute-on-chronic kidney injury at hospital discharge is associated with long-term dialysis and mortality. Kidney International, 2011, 80, 1222-1230.	5. 2	163
7	Pentoxifylline Attenuates Tubulointerstitial Fibrosis by Blocking Smad3/4-Activated Transcription and Profibrogenic Effects of Connective Tissue Growth Factor. Journal of the American Society of Nephrology: JASN, 2005, 16, 2702-2713.	6.1	142
8	Rate of decline of residual renal function is associated with all-cause mortality and technique failure in patients on long-term peritoneal dialysis. Nephrology Dialysis Transplantation, 2009, 24, 2909-2914.	0.7	122
9	Lineage Tracing Reveals Distinctive Fates for Mesothelial Cells and Submesothelial Fibroblasts during Peritoneal Injury. Journal of the American Society of Nephrology: JASN, 2014, 25, 2847-2858.	6.1	117
10	Cognitive-behavioral therapy for sleep disturbance decreases inflammatory cytokines and oxidative stress in hemodialysis patients. Kidney International, 2011, 80, 415-422.	5.2	108
11	Pentoxifylline Attenuated the Renal Disease Progression in Rats with Remnant Kidney. Journal of the American Society of Nephrology: JASN, 2002, 13, 2916-2929.	6.1	106
12	Comparison of residual renal function in patients undergoing twiceâ€weekly versus threeâ€timesâ€weekly haemodialysis. Nephrology, 2009, 14, 59-64.	1.6	105
13	Adiponectin in peritoneal dialysis patients: a comparison with hemodialysis patients and subjects with normal renal function. American Journal of Kidney Diseases, 2004, 43, 1047-1055.	1.9	95
14	Cognitive-Behavioral Therapy for Sleep Disturbance in Patients Undergoing Peritoneal Dialysis: A Pilot Randomized Controlled Trial. American Journal of Kidney Diseases, 2008, 52, 314-323.	1.9	89
15	Multidisciplinary Care Program for Advanced Chronic Kidney Disease: Reduces Renal Replacement and Medical Costs. American Journal of Medicine, 2015, 128, 68-76.	1.5	88
16	Impact of timing of renal replacement therapy initiation on outcome of septic acute kidney injury. Critical Care, 2011, 15, R134.	5 . 8	87
17	Higher systemic inflammation is associated with poorer sleep quality in stable haemodialysis patients. Nephrology Dialysis Transplantation, 2008, 24, 247-251.	0.7	82
18	Primary aldosteronism. Journal of Hypertension, 2011, 29, 1778-1786.	0.5	81

#	Article	IF	CITATIONS
19	The 90-day mortality and the subsequent renal recovery in critically ill surgical patients requiring acute renal replacement therapy. American Journal of Surgery, 2009, 198, 325-332.	1.8	78
20	Pentoxifylline attenuates experimental mesangial proliferative glomerulonephritis. Kidney International, 1999, 56, 932-943.	5. 2	74
21	Sexual dysfunction in female hemodialysis patients: A multicenter study. Kidney International, 2005, 68, 760-765.	5.2	74
22	Pentoxifylline ameliorates proteinuria through suppression of renal monocyte chemoattractant protein-1 in patients with proteinuric primary glomerular diseases. Kidney International, 2006, 69, 1410-1415.	5.2	66
23	Dual Regulation of Tumor Necrosis Factor-α-Induced CCL2/Monocyte Chemoattractant Protein-1 Expression in Vascular Smooth Muscle Cells by Nuclear Factor-ÎB and Activator Protein-1: Modulation by Type III Phosphodiesterase Inhibition. Journal of Pharmacology and Experimental Therapeutics, 2004, 309. 978-986.	2.5	62
24	Predictors of Faster Decline of Residual Renal Function in Taiwanese Peritoneal Dialysis Patients. Peritoneal Dialysis International, 2008, 28, 191-195.	2.3	62
25	Clinical Outcomes and Predictors for ESRD and Mortality in Primary GN. Clinical Journal of the American Society of Nephrology: CJASN, 2012, 7, 1401-1408.	4.5	61
26	Economic, Social, and Psychological Factors Associated With Healthâ€Related Quality of Life of Chronic Hemodialysis Patients in Northern Taiwan: A Multicenter Study. Artificial Organs, 2009, 33, 61-68.	1.9	60
27	The association of higher depressive symptoms and sexual dysfunction in male haemodialysis patients. Nephrology Dialysis Transplantation, 2007, 22, 857-861.	0.7	59
28	Outcomes of Stage 3–5 Chronic Kidney Disease before End-Stage Renal Disease at a Single Center in Taiwan. Nephron Clinical Practice, 2008, 109, c109-c118.	2.3	58
29	Health-Related Quality of Life of Hemodialysis Patients in Taiwan: A Multicenter Study. Blood Purification, 2004, 22, 490-498.	1.8	55
30	Vasodilator Agents Modulate Rat Glomerular Mesangial Cell Growth and Collagen Synthesis. Nephron, 1995, 70, 91-99.	1.8	52
31	Pentoxifylline Inhibits PDGF-induced Proliferation of and TGF-β-stimulated Collagen Synthesis by Vascular Smooth Muscle Cells. Journal of Molecular and Cellular Cardiology, 1999, 31, 773-783.	1.9	52
32	Pentoxifylline suppresses renal tumour necrosis factor-Â and ameliorates experimental crescentic glomerulonephritis in rats. Nephrology Dialysis Transplantation, 2004, 19, 1106-1115.	0.7	51
33	Dipyridamole inhibits TGF-β–induced collagen gene expression in human peritoneal mesothelial cells. Kidney International, 2001, 60, 1249-1257.	5.2	49
34	Nasal Carriage of Methicillin-resistant Staphylococcus aureus Is Associated with Higher All-Cause Mortality in Hemodialysis Patients. Clinical Journal of the American Society of Nephrology: CJASN, 2011, 6, 167-174.	4.5	49
35	Inhibition by pentoxifylline of TNFâ€Î±â€stimulated fractalkine production in vascular smooth muscle cells: evidence for mediation by NFâ€ <i>ΰ</i> B downâ€regulation. British Journal of Pharmacology, 2003, 138, 950-958.	5.4	45
36	Pentoxifylline inhibits human peritoneal mesothelial cell growth and collagen synthesis: Effects on TGF- \hat{l}^2 . Kidney International, 2000, 57, 2626-2633.	5. 2	44

#	Article	IF	CITATIONS
37	Chronic Fatigue in Long-Term Peritoneal Dialysis Patients. American Journal of Nephrology, 2001, 21, 479-485.	3.1	44
38	Pentoxifylline modulates intracellular signalling of TGF-Â in cultured human peritoneal mesothelial cells: implications for prevention of encapsulating peritoneal sclerosis. Nephrology Dialysis Transplantation, 2003, 18, 670-676.	0.7	44
39	Early activation of bradykinin B2 receptor aggravates reactive oxygen species generation and renal damage in ischemia/reperfusion injury. Free Radical Biology and Medicine, 2006, 41, 1304-1314.	2.9	43
40	Association of serum fetuin A with truncal obesity and dyslipidemia in non-diabetic hemodialysis patients. European Journal of Endocrinology, 2009, 160, 777-783.	3.7	42
41	Safety Issues of Long-Term Glucose Load in Patients on Peritoneal Dialysisâ€"A 7-Year Cohort Study. PLoS ONE, 2012, 7, e30337.	2.5	42
42	Interleukin-18 is a strong predictor of hospitalization in haemodialysis patients. Nephrology Dialysis Transplantation, 2004, 19, 2810-2815.	0.7	40
43	Mini-Laparotomy Implantation of Peritoneal Dialysis Catheters: Outcome and Rescue. Peritoneal Dialysis International, 2010, 30, 513-518.	2.3	40
44	Pentoxifylline Inhibits Transforming Growth Factor-Beta Signaling and Renal Fibrosis in Experimental Crescentic Glomerulonephritis in Rats. American Journal of Nephrology, 2009, 29, 43-53.	3.1	37
45	Impact of Peritoneal Membrane Transport on Technique Failure and Patient Survival in a Population on Automated Peritoneal Dialysis. ASAIO Journal, 1999, 45, 568-573.	1.6	36
46	Systemic Lupus Erythematosus and Peritoneal Dialysis: Outcomes and Infectious Complications. Peritoneal Dialysis International, 2001, 21, 143-148.	2.3	35
47	Surgical Management of Refractory Exit-Site/Tunnel Infection of Tenckhoff Catheter: Technical Innovations of Partial Replantation. Peritoneal Dialysis International, 1999, 19, 451-454.	2.3	34
48	Pentoxifylline Inhibits Platelet-Derived Growth Factor-Stimulated Cyclin D1 Expression in Mesangial Cells by Blocking Akt Membrane Translocation. Molecular Pharmacology, 2003, 64, 811-822.	2.3	34
49	Lifetime Costs for Peritoneal Dialysis and Hemodialysis in Patients in Taiwan. Peritoneal Dialysis International, 2013, 33, 671-678.	2.3	34
50	Blockade of cysteine-rich protein 61 attenuates renal inflammation and fibrosis after ischemic kidney injury. American Journal of Physiology - Renal Physiology, 2014, 307, F581-F592.	2.7	34
51	Poor Renal Outcome of Antineutrophil Cytoplasmic Antibody Negative Pauci-immune Glomerulonephritis in Taiwanese. Journal of the Formosan Medical Association, 2006, 105, 804-812.	1.7	33
52	Associations of metabolic syndrome and its components with cardiovascular outcomes among non-diabetic patients undergoing maintenance peritoneal dialysis. Nephrology Dialysis Transplantation, 2011, 26, 4047-4054.	0.7	33
53	Pentoxifylline: A potential therapy for chronic kidney disease. Nephrology, 2004, 9, 198-204.	1.6	32
54	Low-Density Lipoprotein Cholesterol: Association with Mortality and Hospitalization in Hemodialysis Patients. Blood Purification, 2005, 23, 134-140.	1.8	32

#	Article	IF	CITATIONS
55	Higher plasma interleukin-18 levels associated with poor quality of sleep in peritoneal dialysis patients. Nephrology Dialysis Transplantation, 2007, 22, 3606-3609.	0.7	32
56	Relationship between Dialysis Adequacy and Quality of Life in Long-Term Peritoneal Dialysis Patients. Peritoneal Dialysis International, 2000, 20, 534-540.	2.3	31
57	Initial Glucose Load Predicts Technique Survival in Patients on Chronic Peritoneal Dialysis. American Journal of Nephrology, 2008, 28, 765-771.	3.1	31
58	Clinical Findings and Outcomes of Intra-Hemodialysis Cardiopulmonary Resuscitation. American Journal of Nephrology, 1999, 19, 468-473.	3.1	30
59	Treating baclofen overdose by hemodialysis. American Journal of Emergency Medicine, 2012, 30, 1654.e5-1654.e7.	1.6	30
60	Tumor necrosis factor- \hat{l}_{\pm} stimulates fractalkine production by mesangial cells and regulates monocyte transmigration: Down-regulation by cAMP. Kidney International, 2003, 63, 474-486.	5.2	29
61	Plasma Interleukin-18 Levels in Chronic Renal Failure and Continuous Ambulatory Peritoneal Dialysis. Blood Purification, 2005, 23, 144-148.	1.8	28
62	Impact of Near-Death Experiences on Dialysis Patients: A Multicenter Collaborative Study. American Journal of Kidney Diseases, 2007, 50, 124-132.e2.	1.9	28
63	Cysteine-Rich Protein 61 Plays a Proinflammatory Role in Obstructive Kidney Fibrosis. PLoS ONE, 2013, 8, e56481.	2.5	27
64	Sexual Dysfunction in Peritoneal Dialysis Patients. American Journal of Nephrology, 2007, 27, 615-621.	3.1	26
65	Effect of Diuretic Use on 30-Day Postdialysis Mortality in Critically III Patients Receiving Acute Dialysis. PLoS ONE, 2012, 7, e30836.	2.5	25
66	Dipyridamole inhibits PDGF-stimulated human peritoneal mesothelial cell proliferation. Kidney International, 2001, 60, 872-881.	5.2	24
67	Expression of CX3CL1/fractalkine by mesangial cells in vitro and in acute anti-Thy1 glomerulonephritis in rats. Nephrology Dialysis Transplantation, 2003, 18, 2505-2514.	0.7	24
68	Fibrin-Induced Epithelial-to-Mesenchymal Transition of Peritoneal Mesothelial Cells as a Mechanism of Peritoneal Fibrosis: Effects of Pentoxifylline. PLoS ONE, 2012, 7, e44765.	2.5	24
69	Natural Changes in Peritoneal Equilibration Test Results in Continuous Ambulatory Peritoneal Dialysis Patients: A Retrospective, Seven Year Cohortâ€∫Survey. Artificial Organs, 2000, 24, 261-264.	1.9	23
70	The aggressiveness of urinary tract urothelial carcinoma increases with the severity of chronic kidney disease. BJU International, 2009, 104, 1471-1474.	2.5	23
71	Tamoxifen Downregulates Connective Tissue Growth Factor to Ameliorate Peritoneal Fibrosis. Blood Purification, 2011, 31, 252-258.	1.8	23
72	Dipyridamole inhibits human peritoneal mesothelial cell proliferation in vitro and attenuates rat peritoneal fibrosis in vivo. Kidney International, 2001, 59, 2316-2324.	5.2	22

#	Article	IF	CITATIONS
73	Preservation of peritoneal morphology and function by pentoxifylline in a rat model of peritoneal dialysis: molecular studies. Nephrology Dialysis Transplantation, 2008, 23, 3831-3840.	0.7	22
74	Therapeutic efficacy of pentoxifylline on proteinuria and renal progression: an update. Journal of Biomedical Science, 2017, 24, 84.	7.0	22
75	Effect of Intraperitoneally Administered Agents on Human Peritoneal Mesothelial Cell Growth. Nephron, 1995, 71, 23-28.	1.8	21
76	Effects of Pentoxifylline on Peritoneal Fibroblasts and Silica-Induced Peritoneal Fibrosis. Peritoneal Dialysis International, 2003, 23, 228-236.	2.3	21
77	Life expectancy, expected years of life lost and survival of hemodialysis and peritoneal dialysis patients. Journal of Nephrology, 2010, 23, 677-82.	2.0	21
78	The Renoprotective Potential of Pentoxifylline in Chronic Kidney Disease. Journal of the Chinese Medical Association, 2005, 68, 99-105.	1.4	19
79	Diltiazem suppresses collagen synthesis and IL- $1\hat{l}^2$ -induced TGF- \hat{l}^21 production on human peritoneal mesothelial cells. Nephrology Dialysis Transplantation, 2006, 21, 1340-1347.	0.7	19
80	Hyperuricemia Associated With Rapid Renal Function Decline in Elderly Taiwanese Subjects. Journal of the Formosan Medical Association, 2009, 108, 921-928.	1.7	19
81	Association of Low Serum Fetuin A Levels With Poor Arteriovenous Access Patency in Patients Undergoing Maintenance Hemodialysis. American Journal of Kidney Diseases, 2010, 56, 720-727.	1.9	19
82	Risk Factors for High Dialysate Glucose use in PD Patients—A Retrospective 5-Year Cohort Study. Peritoneal Dialysis International, 2010, 30, 448-455.	2.3	19
83	Pleiotropic Effects of Sevelamer Beyond Phosphate Binding in End-Stage Renal Disease Patients. Clinical Drug Investigation, 2011, 31, 257-267.	2.2	19
84	Seven-Year Follow-Up of Peritoneal Dialysis Patients in Taiwan. Peritoneal Dialysis International, 2009, 29, 450-457.	2.3	18
85	Comparison of self-reported health-related quality of life between Taiwan hemodialysis and peritoneal dialysis patients: a multi-center collaborative study. Quality of Life Research, 2011, 20, 399-405.	3.1	18
86	Fibroadhesive Form of Tuberculous Peritonitis: Chyloperitoneum in a Patient Undergoing Automated Peritoneal Dialysis. Nephron, 1996, 72, 708-711.	1.8	17
87	Viral Hepatitis in Continuous Ambulatory Peritoneal Dialysis Patients in an Endemic Area for Hepatitis B and C Infection: The Taiwan Experience. Blood Purification, 1997, 15, 195-199.	1.8	17
88	YC-1-inhibited proliferation of rat mesangial cells through suppression of cyclin D1â€"Independent of cGMP pathway and partially reversed by p38 MAPK inhibitor. European Journal of Pharmacology, 2005, 517, 1-10.	3.5	17
89	Factors associated with metabolic acidosis in patients receiving parenteral nutrition. Nephrology, 2007, 12, 3-7.	1.6	17
90	Lercanidipine-Induced Chyloperitoneum in Patients on Peritoneal Dialysis. Peritoneal Dialysis International, 2008, 28, 632-636.	2.3	17

#	Article	IF	CITATIONS
91	Are Both Psychological and Physical Dimensions in Health-Related Quality of Life Associated with Mortality in Hemodialysis Patients: A 7-Year Taiwan Cohort Study. Blood Purification, 2010, 30, 98-105.	1.8	17
92	Viral Hepatitis Infection Should Be Considered for Evaluating Uremic Pruritus in Continuous Ambulatory Peritoneal Dialysis Patients. Blood Purification, 1998, 16, 147-153.	1.8	16
93	Association between serum aspartate transaminase and homocysteine levels in hemodialysis patients. American Journal of Kidney Diseases, 2002, 40, 1195-1201.	1.9	16
94	Correlations Between Spiritual Beliefs and Healthâ€Related Quality of Life of Chronic Hemodialysis Patients in Taiwan. Artificial Organs, 2009, 33, 576-579.	1.9	16
95	loxitalamate Induces Renal Tubular Apoptosis via Activation of Renal Efferent Nerve–Mediated Adrenergic Signaling, Renin Activity, and Reactive Oxygen Species Production in Rats. Toxicological Sciences, 2010, 114, 149-158.	3.1	16
96	Autonomic dysfunction in chronic kidney disease: An old problem in a new era. Journal of the Formosan Medical Association, 2016, 115, 687-688.	1.7	16
97	Antibiotics induce apoptosis of human peritoneal mesothelial cells. Nephrology, 2003, 8, 142-149.	1.6	15
98	Women on hemodialysis have lower self-reported health-related quality of life scores but better survival than men. Journal of Nephrology, 2013, 26, 366-374.	2.0	14
99	Effects of Intraperitoneal Antibiotics on Human Peritoneal Mesothelial Cell Growth. Nephron, 1996, 74, 694-700.	0.6	13
100	Extracellular Matrix Proteins Modulate Human Peritoneal Mesothelial Cell Behavior. Nephron, 1997, 75, 188-195.	0.6	13
101	Plasma Interleukin-18 Levels in Hemodialysis Patients: Increased by Dialysis Process and Association with Interleukin-6 and Tumor Necrotic Factor-α. Blood Purification, 2006, 24, 174-179.	1.8	12
102	Impact of Spiritual and Religious Activity on Quality of Sleep in Hemodialysis Patients. Blood Purification, 2008, 26, 221-225.	1.8	12
103	N-Acetylcysteine-Mediated Antioxidation Prevents Hyperglycemia-Induced Apoptosis and Collagen Synthesis in Rat Mesangial Cells. American Journal of Nephrology, 2009, 29, 192-202.	3.1	12
104	Disintegrin Modulates Rat Glomerular Mesangial Cell Behavior. Nephron, 1995, 70, 83-90.	1.8	11
105	Bradykinin enhances reactive oxygen species generation, mitochondrial injury, and cell death induced by ATP depletion—A role of the phospholipase CCa2+ pathway. Free Radical Biology and Medicine, 2007, 43, 702-710.	2.9	11
106	Benefits of Sevelamer on Markers of Bone Turnover in Taiwanese Hemodialysis Patients. Journal of the Formosan Medical Association, 2010, 109, 663-672.	1.7	11
107	Hemorrhagic Stroke in Chronic Dialysis Patients. Renal Failure, 2004, 26, 165-170.	2.1	10
108	Effects of pentoxifylline on peritoneal fibroblasts and silica-induced peritoneal fibrosis. Peritoneal Dialysis International, 2003, 23, 228-36.	2.3	10

#	Article	IF	Citations
109	Peritoneal thickening is not inevitable in long-term peritoneal dialysis and is associated with peritoneal transport characteristics: a two-centre sonographic study. Nephrology Dialysis Transplantation, 2007, 23, 1005-1010.	0.7	9
110	Seven-year follow-up of peritoneal dialysis patients in Taiwan. Peritoneal Dialysis International, 2009, 29, 450-7.	2.3	9
111	RENAL KALLIKREIN IN CHRONIC HYPOXIC RATS. Clinical and Experimental Pharmacology and Physiology, 1996, 23, 819-824.	1.9	8
112	Intraperitoneal Vascular Endothelial Growth Factor C Level Is Related to Peritoneal Dialysis Ultrafiltration. Blood Purification, 2009, 28, 69-74.	1.8	8
113	Peritoneal fibrosis and its prevention. Nephrology, 2002, 7, 227-232.	1.6	7
114	Peritoneal Fibrosing Syndrome: Pathogenetic Mechanism and Current Therapeutic Strategies. Journal of the Chinese Medical Association, 2005, 68, 401-405.	1.4	7
115	Influence of Relative Hypoparathyroidism on the Responsiveness to Recombinant Human Erythropoietin in Hemodialysis Patients. Blood Purification, 2003, 21, 220-224.	1.8	6
116	<i>In Vitro</i> Study of Peritoneal Fibrosis. Peritoneal Dialysis International, 2007, 27, 72-75.	2.3	5
117	Outcomes following Dialysis for Acute Kidney Injury among Different Stages of Chronic Kidney Disease. American Journal of Nephrology, 2011, 34, 95-103.	3.1	5
118	Antineutrophil cytoplasmic antibody-associated glomerulonephritis in Taiwanese. Nephrology, 2004, 9, 297-303.	1.6	4
119	Urinary kallikrein excretion is related to renal function change and inflammatory status in chronic kidney disease patients receiving angiotensin II receptor blocker treatment. Nephrology, 2008, 13, 198-203.	1.6	4
120	Maintenance haemodialysis and delayed administration of appropriate antibiotics increase 30-day mortality among patients with non-hospital-acquired meticillin-resistant Staphylococcus aureus bacteraemia. International Journal of Antimicrobial Agents, 2010, 35, 511-512.	2.5	4
121	Combining body mass index and serum potassium to urine potassium clearance ratio is an alternative method to predict primary aldosteronism. Clinica Chimica Acta, 2011, 412, 1637-1642.	1.1	4
122	Renovascular disease in Taiwan: A long-term nationwide population study. International Journal of Cardiology, 2013, 168, 541-542.	1.7	2
123	Cardiopulmonary resuscitation in dialysis patients. Journal of Interventional Cardiac Electrophysiology, 2002, 6, 160-162.	1.0	1
124	Peritoneal Permeability in Patients with Encapsulating Peritoneal Sclerosis. American Journal of Kidney Diseases, 2006, 48, 875.	1.9	1
125	Associations between urinary cysteine-rich protein 61 excretion and kidney function decline in outpatients with chronic kidney disease: a prospective cohort study in Taiwan. BMJ Open, 2021, 11, e051165.	1.9	1
126	Primary biliary cirrhosis associated with minimal change disease. Nephrology Dialysis Transplantation, 2007, 22, 966-967.	0.7	0

Tun-Jun Tsai

#	Article	IF	CITATIONS
127	Gender effect on quality of life in hemodialysis patients: response to Einollahi and Motalebi. Journal of Nephrology, 2014, 27, 593-593.	2.0	0
128	A tribute to Professor Wan-Yu Chen. Journal of the Formosan Medical Association, 2015, 114, 791-792.	1.7	0
129	Web-based pulse analysis system for detection of acute kidney injury. , 2015, , .		O
130	Lysophosphatidic Acid and Renal Fibrosis. Recent Patents on Endocrine, Metabolic & Immune Drug Discovery, 2008, 2, 204-210.	0.6	0