

Fredrik H Nystrom

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3199724/fredrik-h-nystrom-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

86
papers

2,448
citations

28
h-index

47
g-index

91
ext. papers

2,834
ext. citations

4.8
avg, IF

4.62
L-index

#	Paper	IF	Citations
86	A cross-over study of postprandial effects from moist snuff and red wine on metabolic rate, appetite-related hormones and glucose.. <i>Drug and Alcohol Dependence</i> , 2022 , 236, 109479	4.9	
85	Plasma mid-regional pro-atrial natriuretic peptide predicts cardiovascular events in patients with type 2 diabetes independently of subclinical organ damage. <i>Diabetes Research and Clinical Practice</i> , 2021 , 182, 109095	7.4	0
84	Circulating TNFR1I levels predict incidence of ischemic heart disease and total mortality, independently of intima media thickness and pulse wave velocity in male with type 2 diabetes. <i>Heart and Vessels</i> , 2021 , 36, 1591-1596	2.1	
83	Relationships between cardiovascular risk factors and white-coat hypertension diagnosed by home blood pressure recordings in a middle-aged population. <i>Journal of Hypertension</i> , 2021 , 39, 2009-2014	1.9	3
82	Plasma Protein Profile of Carotid Artery Atherosclerosis and Atherosclerotic Outcomes: Meta-Analyses and Mendelian Randomization Analyses. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2021 , 41, 1777-1788	9.4	4
81	Prevalence of Subclinical Coronary Artery Atherosclerosis in the General Population. <i>Circulation</i> , 2021 , 144, 916-929	16.7	17
80	Education of the primary health care staff based on acceptance and commitment therapy is associated with reduced sick leave in a prospective controlled trial. <i>BMC Family Practice</i> , 2021 , 22, 179	2.6	
79	A Pilot Study of Perioperative External Circumferential Cryoablation of Human Renal Arteries for Sympathetic Denervation. <i>Vascular Specialist International</i> , 2020 , 36, 151-157	1	2
78	Is there an association between serum 25(OH)D and mental well-being in patients with type 2 diabetes? Results from a cohort study in primary care. <i>Hormones</i> , 2020 , 19, 361-367	3.1	
77	Weight gain and blood pressure. <i>Journal of Hypertension</i> , 2020 , 38, 387-394	1.9	2
76	Growth differentiation factor 15 (GDF-15) is a potential biomarker of both diabetic kidney disease and future cardiovascular events in cohorts of individuals with type 2 diabetes: a proteomics approach. <i>Uppsala Journal of Medical Sciences</i> , 2020 , 125, 37-43	2.8	15
75	A pilot study of hypertension management using a telemedicine treatment approach. <i>Blood Pressure Monitoring</i> , 2020 , 25, 18-21	1.3	1
74	The association between plasma proteomics and incident cardiovascular disease identifies MMP-12 as a promising cardiovascular risk marker in patients with chronic kidney disease. <i>Atherosclerosis</i> , 2020 , 307, 11-15	3.1	3
73	The association between alcohol consumption, cardiac biomarkers, left atrial size and re-ablation in patients with atrial fibrillation referred for catheter ablation. <i>PLoS ONE</i> , 2019 , 14, e0215121	3.7	4
72	Expression of GHR and Downstream Signaling Genes in Human Adipose Tissue-Relation to Obesity and Weight Change. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 1459-1470	5.6	8
71	Subjects with high fasting insulin also have higher postprandial GLP-1 and glucagon levels than controls with lower insulin. <i>Nutrition Research</i> , 2019 , 72, 111-120	4	
70	Predictors of improvement in arrhythmia-specific symptoms and health-related quality of life after catheter ablation of atrial fibrillation. <i>Clinical Cardiology</i> , 2019 , 42, 247-255	3.3	11

69	The association between circulating endostatin and a disturbed circadian blood pressure pattern in patients with type 2 diabetes. <i>Blood Pressure</i> , 2018 , 27, 215-221	1.7	1
68	A randomised study in young subjects of the effects of eating extra fruit or nuts on periodontal inflammation. <i>BDJ Open</i> , 2018 , 4, 17022	3.3	4
67	Overweight and obesity impair left ventricular systolic function as measured by left ventricular ejection fraction and global longitudinal strain. <i>Cardiovascular Diabetology</i> , 2018 , 17, 113	8.7	30
66	Multiplex proteomics for prediction of major cardiovascular events in type 2 diabetes. <i>Diabetologia</i> , 2018 , 61, 1748-1757	10.3	28
65	Skin microvascular endothelial dysfunction is associated with type 2 diabetes independently of microalbuminuria and arterial stiffness. <i>Diabetes and Vascular Disease Research</i> , 2017 , 14, 363-371	3.3	20
64	A randomized cross-over study of the effects of macronutrient composition and meal frequency on GLP-1, ghrelin and energy expenditure in humans. <i>Peptides</i> , 2017 , 93, 20-26	3.8	6
63	A randomized cross-over study of the acute effects of running 5 km on glucose, insulin, metabolic rate, cortisol and Troponin T. <i>PLoS ONE</i> , 2017 , 12, e0179401	3.7	6
62	Association of soluble tumor necrosis factor receptors 1 and 2 with nephropathy, cardiovascular events, and total mortality in type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2016 , 15, 40	8.7	50
61	Diastolic orthostatic hypertension and cardiovascular prognosis in type 2 diabetes: a prospective cohort study. <i>Cardiovascular Diabetology</i> , 2016 , 15, 83	8.7	17
60	A Randomized Study of the Effects of Additional Fruit and Nuts Consumption on Hepatic Fat Content, Cardiovascular Risk Factors and Basal Metabolic Rate. <i>PLoS ONE</i> , 2016 , 11, e0147149	3.7	25
59	Microdialysis and proteomics of subcutaneous interstitial fluid reveals increased galectin-1 in type 2 diabetes patients. <i>Metabolism: Clinical and Experimental</i> , 2016 , 65, 998-1006	12.7	13
58	A randomized trial of cold-exposure on energy expenditure and supraclavicular brown adipose tissue volume in humans. <i>Metabolism: Clinical and Experimental</i> , 2016 , 65, 926-34	12.7	20
57	Aortic pulse wave velocity predicts incident cardiovascular events in patients with type 2 diabetes treated in primary care. <i>Journal of Diabetes and Its Complications</i> , 2016 , 30, 1223-8	3.2	18
56	Phosphatidylethanol Compared with Other Blood Tests as a Biomarker of Moderate Alcohol Consumption in Healthy Volunteers: A Prospective Randomized Study. <i>Alcohol and Alcoholism</i> , 2015 , 50, 399-406	3.5	59
55	Left ventricular diastolic function, assessed by echocardiography and tissue Doppler imaging, is a strong predictor of cardiovascular events, superior to global left ventricular longitudinal strain, in patients with type 2 diabetes. <i>European Heart Journal Cardiovascular Imaging</i> , 2015 , 16, 1000-7	4.1	34
54	Comparisons of automated blood pressures in a primary health care setting with self-measurements at the office and at home using the Omron i-C10 device. <i>Blood Pressure Monitoring</i> , 2015 , 20, 98-103	1.3	26
53	A prospective observational study of all-cause mortality in relation to serum 25-OH vitamin D3 and parathyroid hormone levels in patients with type 2 diabetes. <i>Diabetology and Metabolic Syndrome</i> , 2015 , 7, 53	5.6	9
52	Serum leptin levels are independently related to the incidence of ischemic heart disease in a prospective study of patients with type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2015 , 14, 62	8.7	18

51	Microcirculation assessment using an individualized model for diffuse reflectance spectroscopy and conventional laser Doppler flowmetry. <i>Journal of Biomedical Optics</i> , 2014 , 19, 057002	3.5	12
50	Associations of genetic polymorphisms in the renin-angiotensin system with central aortic and ambulatory blood pressure in type 2 diabetic patients. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , 2014 , 15, 61-8	3	2
49	Advice to follow a low-carbohydrate diet has a favourable impact on low-grade inflammation in type 2 diabetes compared with advice to follow a low-fat diet. <i>Annals of Medicine</i> , 2014 , 46, 182-7	1.5	49
48	Sagittal abdominal diameter is a more independent measure compared with waist circumference to predict arterial stiffness in subjects with type 2 diabetes--a prospective observational cohort study. <i>Cardiovascular Diabetology</i> , 2013 , 12, 55	8.7	33
47	Toe brachial index in middle aged patients with diabetes mellitus type 2: not just a peripheral issue. <i>Diabetes Research and Clinical Practice</i> , 2013 , 100, 195-202	7.4	20
46	Effects of exercise with or without blueberries in the diet on cardio-metabolic risk factors: an exploratory pilot study in healthy subjects. <i>Upsala Journal of Medical Sciences</i> , 2013 , 118, 247-55	2.8	17
45	A randomized cross-over trial of the postprandial effects of three different diets in patients with type 2 diabetes. <i>PLoS ONE</i> , 2013 , 8, e79324	3.7	12
44	A randomized trial of protein supplementation compared with extra fast food on the effects of resistance training to increase metabolism. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2012 , 72, 471-8	2	5
43	Is there an underestimation of intima-media thickness based on M-mode ultrasound technique in the abdominal aorta?. <i>Clinical Physiology and Functional Imaging</i> , 2012 , 32, 1-4	2.4	2
42	Ethyl glucuronide in human hair after daily consumption of 16 or 32 g of ethanol for 3 months. <i>Forensic Science International</i> , 2012 , 215, 51-5	2.6	75
41	Ambulatory systolic blood pressure predicts left ventricular mass in type 2 diabetes, independent of central systolic blood pressure. <i>Blood Pressure Monitoring</i> , 2012 , 17, 139-44	1.3	14
40	Cardiovascular risk factors related to the PPARIPro12Ala polymorphism in patients with type 2 diabetes are gender dependent. <i>Blood Pressure</i> , 2012 , 21, 122-7	1.7	5
39	Circadian blood pressure variation in patients with type 2 diabetes--relationship to macro- and microvascular subclinical organ damage. <i>Primary Care Diabetes</i> , 2011 , 5, 167-73	2.4	18
38	Transient increase in HDL-cholesterol during weight gain by hyperalimantation in healthy subjects. <i>Obesity</i> , 2011 , 19, 812-7	8	5
37	Effects of moderate red wine consumption on liver fat and blood lipids: a prospective randomized study. <i>Annals of Medicine</i> , 2011 , 43, 545-54	1.5	38
36	Identification of adipocyte genes regulated by caloric intake. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011 , 96, E413-8	5.6	57
35	A hierarchical whole-body modeling approach elucidates the link between in Vitro insulin signaling and in Vivo glucose homeostasis. <i>Journal of Biological Chemistry</i> , 2011 , 286, 26028-41	5.4	59
34	Young healthy individuals develop lack of energy when adopting an obesity provoking behaviour for 4 weeks: a phenomenological analysis. <i>Scandinavian Journal of Caring Sciences</i> , 2010 , 24, 565-71	2.3	4

33	Weight gain by hyperalimantation elevates C-reactive protein levels but does not affect circulating levels of adiponectin or resistin in healthy subjects. <i>European Journal of Endocrinology</i> , 2010 , 163, 879-85	6.5	14
32	Comparison of the subjective sense of high or low metabolism and objectively measured resting metabolic rate. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2010 , 70, 334-7	2	3
31	Reduced arteriovenous shunting capacity after local heating and redistribution of baseline skin blood flow in type 2 diabetes assessed with velocity-resolved quantitative laser Doppler flowmetry. <i>Diabetes</i> , 2010 , 59, 1578-84	0.9	35
30	Doubling of water intake increases daytime blood pressure and reduces vertigo in healthy subjects. <i>Clinical and Experimental Hypertension</i> , 2010 , 32, 439-43	2.2	6
29	An obesity provoking behaviour negatively influences young normal weight subjectsThealth related quality of life and causes depressive symptoms. <i>Eating Behaviors</i> , 2010 , 11, 247-52	3	7
28	Association between admission supine systolic blood pressure and 1-year mortality in patients admitted to the intensive care unit for acute chest pain. <i>JAMA - Journal of the American Medical Association</i> , 2010 , 303, 1167-72	27.4	18
27	Long-term increase of fat mass after a four week intervention with fast food based hyper-alimentation and limitation of physical activity. <i>Nutrition and Metabolism</i> , 2010 , 7, 68	4.6	10
26	Short-term overeating induces insulin resistance in fat cells in lean human subjects. <i>Molecular Medicine</i> , 2009 , 15, 228-34	6.2	25
25	Regulation of the fibrosis and angiogenesis promoter SPARC/osteonectin in human adipose tissue by weight change, leptin, insulin, and glucose. <i>Diabetes</i> , 2009 , 58, 1780-8	0.9	86
24	Men develop more intraabdominal obesity and signs of the metabolic syndrome after hyperalimantation than women. <i>Metabolism: Clinical and Experimental</i> , 2009 , 58, 995-1001	12.7	27
23	Two weeks of overfeeding with candy, but not peanuts, increases insulin levels and body weight. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2009 , 69, 598-605	2	46
22	Retinol-binding protein-4 attenuates insulin-induced phosphorylation of IRS1 and ERK1/2 in primary human adipocytes. <i>FASEB Journal</i> , 2007 , 21, 3696-704	0.9	106
21	Human, but not rat, IRS1 targets to the plasma membrane in both human and rat adipocytes. <i>Biochemical and Biophysical Research Communications</i> , 2007 , 363, 840-5	3.4	10
20	PPAR-gamma response element activity in intact primary human adipocytes: effects of fatty acids. <i>Nutrition</i> , 2006 , 22, 60-8	4.8	52
19	Phosphorylation of IRS1 at serine 307 and serine 312 in response to insulin in human adipocytes. <i>Biochemical and Biophysical Research Communications</i> , 2006 , 342, 1183-7	3.4	16
18	Angiotensin type 1 receptor antagonists induce human in-vitro adipogenesis through peroxisome proliferator-activated receptor-gamma activation. <i>Journal of Hypertension</i> , 2006 , 24, 1809-16	1.9	71
17	Separation and characterization of caveolae subclasses in the plasma membrane of primary adipocytes; segregation of specific proteins and functions. <i>FEBS Journal</i> , 2006 , 273, 3381-92	5.7	45
16	The effect of percutaneous dilatation of renal arterial stenosis on captopril renography in hypertension. <i>Blood Pressure</i> , 2005 , 14, 359-65	1.7	2

15	Glucose transport is equally sensitive to insulin stimulation, but basal and insulin-stimulated transport is higher, in human omental compared with subcutaneous adipocytes. <i>Metabolism: Clinical and Experimental</i> , 2005 , 54, 781-5	12.7	14
14	Insulin resistance in human adipocytes occurs downstream of IRS1 after surgical cell isolation but at the level of phosphorylation of IRS1 in type 2 diabetes. <i>FEBS Journal</i> , 2005 , 272, 141-51	5.7	61
13	Attenuation of insulin-stimulated insulin receptor substrate-1 serine 307 phosphorylation in insulin resistance of type 2 diabetes. <i>Journal of Biological Chemistry</i> , 2005 , 280, 34389-92	5.4	63
12	Triacylglycerol is synthesized in a specific subclass of caveolae in primary adipocytes. <i>Journal of Biological Chemistry</i> , 2005 , 280, 5-8	5.4	110
11	Subcutaneous adipocytes from obese hyperinsulinemic women with polycystic ovary syndrome exhibit normal insulin sensitivity but reduced maximal insulin responsiveness. <i>European Journal of Endocrinology</i> , 2005 , 153, 831-5	6.5	13
10	Lipids and glycosphingolipids in caveolae and surrounding plasma membrane of primary rat adipocytes. <i>FEBS Journal</i> , 2004 , 271, 2028-36		112
9	Colocalization of insulin receptor and insulin receptor substrate-1 to caveolae in primary human adipocytes. Cholesterol depletion blocks insulin signalling for metabolic and mitogenic control. <i>FEBS Journal</i> , 2004 , 271, 2471-9		72
8	Expression of a mutant IRS inhibits metabolic and mitogenic signalling of insulin in human adipocytes. <i>Molecular and Cellular Endocrinology</i> , 2004 , 221, 1-8	4.4	21
7	Cell surface orifices of caveolae and localization of caveolin to the necks of caveolae in adipocytes. <i>Molecular Biology of the Cell</i> , 2003 , 14, 3967-76	3.5	111
6	Individualized growth hormone substitution with normalized IGF-I levels does not stimulate the renin-angiotensin-aldosterone system. <i>Clinical Endocrinology</i> , 2002 , 57, 473-9	3.4	5
5	Insulin stimulates increased catalytic activity of phosphoinositide-dependent kinase-1 by a phosphorylation-dependent mechanism. <i>Biochemistry</i> , 2001 , 40, 11851-9	3.2	31
4	Less difference between office and ambulatory blood pressure in women than in men both before and during antihypertensive treatment. <i>Blood Pressure</i> , 2000 , 9, 340-5	1.7	2
3	Caveolin-1 interacts with the insulin receptor and can differentially modulate insulin signaling in transfected Cos-7 cells and rat adipose cells. <i>Molecular Endocrinology</i> , 1999 , 13, 2013-24		160
2	Insulin signalling: metabolic pathways and mechanisms for specificity. <i>Cellular Signalling</i> , 1999 , 11, 563-74	4.9	162
1	Population-based reference values for IGF-I and IGF-binding protein-1: relations with metabolic and anthropometric variables. <i>European Journal of Endocrinology</i> , 1997 , 136, 165-72	6.5	47