## Ivan Sadowski

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3201489/publications.pdf Version: 2024-02-01



| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | TORC1 signaling modulates Cdk8-dependent <i>GAL</i> gene expression in <i>Saccharomyces cerevisiae</i> . Genetics, 2021, 219, .  | 1.2 | 1         |
| 2  | Diversity of small molecule HIVâ€1 latency reversing agents identified in low―and highâ€ŧhroughput small<br>molecule screens. Medicinal Research Reviews, 2020, 40, 881-908.   | 5.0 | 23        |
| 3  | Management of inadvertent template contamination in production of oligonucleotide qPCR reagents.<br>BioTechniques, 2020, 69, 401-403.  | 0.8 | 1         |
| 4  | Regulation of Skn7-dependent, oxidative stress-induced genes by the RNA polymerase II-CTD<br>phosphatase, Fcp1, and Mediator kinase subunit, Cdk8, in yeast. Journal of Biological Chemistry, 2019,<br>294, 16080-16094.               | 1.6 | 9         |
| 5  | Strategies to eradicate HIV from infected patients: elimination of latent provirus reservoirs. Cellular and Molecular Life Sciences, 2019, 76, 3583-3600.  | 2.4 | 77        |
| 6  | Compounds producing an effective combinatorial regimen for disruption of <scp>HIV</scp> â€1 latency.<br>EMBO Molecular Medicine, 2018, 10, 160-174.  | 3.3 | 25        |
| 7  | HIV Provirus Stably Reproduces Parental Latent and Induced Transcription Phenotypes Regardless of the Chromosomal Integration Site. Journal of Virology, 2016, 90, 5302-5314.  | 1.5 | 18        |
| 8  | Direct non-productive HIV-1 infection in a T-cell line is driven by cellular activation state and NFήB.<br>Retrovirology, 2014, 11, 17.  | 0.9 | 37        |
| 9  | A Doubly Fluorescent HIV-1 Reporter Shows that the Majority of Integrated HIV-1 Is Latent Shortly after Infection. Journal of Virology, 2013, 87, 4716-4727.   | 1.5 | 88        |
| 10 | The PhosphoGRID Saccharomyces cerevisiae protein phosphorylation site database: version 2.0 update.<br>Database: the Journal of Biological Databases and Curation, 2013, 2013, bat026-bat026.  | 1.4 | 96        |
| 11 | An Upstream YY1 Binding Site on the HIV-1 LTR Contributes to Latent Infection. PLoS ONE, 2013, 8, e77052.  | 1.1 | 33        |
| 12 | Cdk8 Regulates Stability of the Transcription Factor Phd1 To Control Pseudohyphal Differentiation of Saccharomyces cerevisiae. Molecular and Cellular Biology, 2012, 32, 664-674.  | 1.1 | 40        |
| 13 | Identification and functional analysis of a second RBF-2 binding site within the HIV-1 promoter.<br>Virology, 2011, 418, 57-66.  | 1.1 | 26        |
| 14 | The Suv39H1 methyltransferase inhibitor chaetocin causes induction of integrated HIV-1 without producing a T cell response. FEBS Letters, 2011, 585, 3549-3554.  | 1.3 | 76        |
| 15 | PhosphoGRID: a database of experimentally verified in vivo protein phosphorylation sites from the<br>budding yeast Saccharomyces cerevisiae. Database: the Journal of Biological Databases and Curation,<br>2010, 2010, bap026-bap026. | 1.4 | 90        |
| 16 | Dominant marker vectors for selecting yeast mating products. Yeast, 2008, 25, 595-599.   | 0.8 | 12        |
| 17 | Specific interaction of TFIIâ€I with an upstream element on the HIVâ€I LTR regulates induction of latent provirus. FEBS Letters, 2008, 582, 3903-3908.   | 1.3 | 25        |
| 18 | Factors Controlling Chromatin Organization and Nucleosome Positioning for Establishment and Maintenance of HIV Latency. Current HIV Research, 2008, 6, 286-295.  | 0.2 | 41        |

Ivan Sadowski

| #  | Article  | IF   | CITATIONS |
|----|--|------|-----------|
| 19 | Disintegrator vectors for single-copy yeast chromosomal integration. Yeast, 2007, 24, 447-455.   | 0.8  | 47        |
| 20 | Induction of chromosomally integrated HIV-1 LTR requires RBF-2 (USF/TFII-I) and RAS/MAPK signaling.<br>Virus Genes, 2007, 35, 215-223.   | 0.7  | 51        |
| 21 | TFII-I Regulates Induction of Chromosomally Integrated Human Immunodeficiency Virus Type 1 Long<br>Terminal Repeat in Cooperation with USF. Journal of Virology, 2005, 79, 4396-4406.                              | 1.5  | 48        |
| 22 | TFII-I and USF (RBF-2) regulate Ras/MAPK-responsive HIV-1 transcription in T cells. European Journal of Cancer, 2005, 41, 2528-2536.   | 1.3  | 31        |
| 23 | Srb10/Cdk8 regulates yeast filamentous growth by phosphorylating the transcription factor Ste12.<br>Nature, 2003, 421, 187-190.  | 13.7 | 144       |
| 24 | Multiple Signals Regulate GAL Transcription in Yeast. Molecular and Cellular Biology, 2000, 20, 3880-3886.   | 1.1  | 69        |
| 25 | Purification of RBF-2, a transcription factor with specificity for the most conservedcis-element of naturally occurring HIV-1 LTRs. Journal of Biomedical Science, 1999, 6, 320-332.                               | 2.6  | 15        |
| 26 | GAL4 Is Regulated by the RNA Polymerase II Holoenzyme–Associated Cyclin-Dependent Protein Kinase<br>SRB10/CDK8. Molecular Cell, 1999, 3, 673-678.  | 4.5  | 128       |
| 27 | Naturally Occurring Human Immunodeficiency Virus Type 1 Long Terminal Repeats Have a Frequently<br>Observed Duplication That Binds RBF-2 and Represses Transcription. Journal of Virology, 1998, 72,<br>6465-6474. | 1.5  | 38        |
| 28 | Characterization of the Basal and Pheromone-Stimulated Phosphorylation States of Ste12p. FEBS<br>Journal, 1997, 245, 241-251.  | 0.2  | 50        |
| 29 | GAL4-VP16 is an unusually potent transcriptional activator. Nature, 1988, 335, 563-564.  | 13.7 | 1,433     |