

Abdulsamie Hanano

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3208943/abdulsamie-hanano-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

26

papers

384

citations

12

h-index

19

g-index

27

ext. papers

517

ext. citations

5.4

avg, IF

3.36

L-index

#	Paper	IF	Citations
26	Characterization of lipid droplets from a <i>Taxus media</i> cell suspension and their potential involvement in trafficking and secretion of paclitaxel.. <i>Plant Cell Reports</i> , 2022 , 1	5.1	0
25	Functional involvement of caleosin/ peroxygenase PdPXG4 in the accumulation of date palm leaf lipid droplets after exposure to dioxins. <i>Environmental Pollution</i> , 2021 , 281, 116966	9.3	1
24	Involvement of hepatic lipid droplets and their associated proteins in the detoxification of aflatoxin B in aflatoxin-resistance BALB/C mouse. <i>Toxicology Reports</i> , 2020 , 7, 795-804	4.8	2
23	Dioxin impacts on lipid metabolism of soil microbes: towards effective detection and bioassessment strategies. <i>Bioresources and Bioprocessing</i> , 2020 , 7,	5.2	1
22	Exposure of NRRL 3357 to the Environmental Toxin, 2,3,7,8-Tetrachlorinated Dibenzo–Dioxin, Results in a Hyper Aflatoxicogenic Phenotype: A Possible Role for Caleosin/Peroxygenase (AfPXG). <i>Frontiers in Microbiology</i> , 2019 , 10, 2338	5.7	6
21	The cytochrome P450 of <i>Bacillus megaterium</i> A14K is induced by 2,3,7,8-Tetrachlorinated dibenzo-p-dioxin: Biophysical, molecular and biochemical determinants. <i>Chemosphere</i> , 2019 , 216, 258-270	8.4	7
20	Arabidopsis plants exposed to dioxin result in a WRINKLED seed phenotype due to 20S proteasomal degradation of WR1. <i>Journal of Experimental Botany</i> , 2018 , 69, 1781-1794	7	8
19	The Peroxygenase Activity of the Caleosin, AfPXG, Modulates the Biosynthesis of Aflatoxins and Their Trafficking and Extracellular Secretion via Lipid Droplets. <i>Frontiers in Microbiology</i> , 2018 , 9, 158	5.7	10
18	Evolutionary, structural and functional analysis of the caleosin/ peroxygenase gene family in the Fungi. <i>BMC Genomics</i> , 2018 , 19, 976	4.5	8
17	Evolutionary and genomic analysis of the caleosin/ peroxygenase (CLO/PXG) gene/protein families in the Viridiplantae. <i>PLoS ONE</i> , 2018 , 13, e0196669	3.7	12
16	Identification of a dioxin-responsive oxylipin signature in roots of date palm: involvement of a 9-hydroperoxide fatty acid reductase, caleosin/ peroxygenase PdPXG2. <i>Scientific Reports</i> , 2018 , 8, 13181	4.9	11
15	Biochemical, Molecular, and Transcriptional Highlights of the Biosynthesis of an Effective Biosurfactant Produced by PHA3, a Petroleum-Dwelling Bacteria. <i>Frontiers in Microbiology</i> , 2017 , 8, 77	5.7	7
14	Specific Caleosin/Peroxygenase and Lipoxygenase Activities Are Tissue-Differentially Expressed in Date Palm (<i>L.</i>) Seedlings and Are Further Induced Following Exposure to the Toxin 2,3,7,8-tetrachlorodibenzo-p-dioxin. <i>Frontiers in Plant Science</i> , 2016 , 7, 2025	6.2	12
13	Biochemical, Transcriptional, and Bioinformatic Analysis of Lipid Droplets from Seeds of Date Palm (<i>Phoenix dactylifera</i> L.) and Their Use as Potent Sequestration Agents against the Toxic Pollutant, 2,3,7,8-Tetrachlorinated Dibenzo-p-Dioxin. <i>Frontiers in Plant Science</i> , 2016 , 7, 836	6.2	13
12	Immuno-detection of dioxins using a recombinant protein of aryl hydrocarbon receptor (AhR) fused with sfGFP. <i>BMC Biotechnology</i> , 2016 , 16, 51	3.5	3
11	<i>Saccharomyces cerevisiae</i> SHSY detoxifies petroleum n-alkanes by an induced CYP52A58 and an enhanced order in cell surface hydrophobicity. <i>Chemosphere</i> , 2015 , 135, 418-26	8.4	13
10	A Caleosin-Like Protein with Peroxygenase Activity Mediates <i>Aspergillus flavus</i> Development, Aflatoxin Accumulation, and Seed Infection. <i>Applied and Environmental Microbiology</i> , 2015 , 81, 6129-44	4.8	17

9	Involvement of the caleosin/peroxygenase RD20 in the control of cell death during Arabidopsis responses to pathogens. <i>Plant Signaling and Behavior</i> , 2015 , 10, e991574	2.5	16
8	Differential tissue accumulation of 2,3,7,8-Tetrachlorinated dibenzo-p-dioxin in Arabidopsis thaliana affects plant chronology, lipid metabolism and seed yield. <i>BMC Plant Biology</i> , 2015 , 15, 193	5.3	19
7	Removal of petroleum-crude oil from aqueous solution by Saccharomyces cerevisiae SHSY strain necessitates at least an inducible CYP450ALK homolog gene. <i>Journal of Basic Microbiology</i> , 2014 , 54, 358-68	2.7	6
6	The reductase activity of the Arabidopsis caleosin RESPONSIVE TO DESSICATION20 mediates gibberellin-dependent flowering time, abscisic acid sensitivity, and tolerance to oxidative stress. <i>Plant Physiology</i> , 2014 , 166, 109-24	6.6	40
5	Traceability of polychlorinated dibenzo-dioxins/furans pollutants in soil and their ecotoxicological effects on genetics, functions and composition of bacterial community. <i>Chemosphere</i> , 2014 , 108, 326-33	8.4	17
4	Phytotoxicity effects and biological responses of Arabidopsis thaliana to 2,3,7,8-tetrachlorinated dibenzo-p-dioxin exposure. <i>Chemosphere</i> , 2014 , 104, 76-84	8.4	18
3	Silencing of Erwinia amylovora sy69 AHL-quorum sensing by a Bacillus simplex AHL-inducible aiiA gene encoding a zinc-dependent N-acyl-homoserine lactonase. <i>Plant Pathology</i> , 2014 , 63, 773-783	2.8	7
2	Plant seed peroxygenase is an original heme-oxygenase with an EF-hand calcium binding motif. <i>Journal of Biological Chemistry</i> , 2006 , 281, 33140-51	5.4	107
1	Stereochemical features of the hydrolysis of 9,10-epoxystearic acid catalysed by plant and mammalian epoxide hydrolases. <i>Biochemical Journal</i> , 2002 , 366, 471-80	3.8	23