

# Bingcheng Xu

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/3210538/bingcheng-xu-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

61  
papers

1,201  
citations

21  
h-index

32  
g-index

65  
ext. papers

1,535  
ext. citations

4.2  
avg, IF

4.3  
L-index

#	Paper	IF	Citations
61	Nitrogen supply improved plant growth and Cd translocation in maize at the silking and physiological maturity under moderate Cd stress.. <i>Ecotoxicology and Environmental Safety</i> , <b>2021</b> , 230, 113137	7	2
60	Characterization of Root System Architecture Traits in Diverse Soybean Genotypes Using a Semi-Hydroponic System.. <i>Plants</i> , <b>2021</b> , 10,	4.5	2
59	Soil Water Availability Changes in Amount and Timing Favor the Growth and Competitiveness of Grass Than a Co-dominant Legume in Their Mixtures. <i>Frontiers in Plant Science</i> , <b>2021</b> , 12, 723839	6.2	0
58	Arbuscular mycorrhizal symbioses alleviating salt stress in maize is associated with a decline in root-to-leaf gradient of Na/K ratio. <i>BMC Plant Biology</i> , <b>2021</b> , 21, 457	5.3	4
57	Responses of soil respiration to rainfall depth and frequency in semiarid grassland communities. <i>Ecohydrology</i> , <b>2021</b> , 14, e2326	2.5	0
56	Aboveground biomass production and dominant species type determined canopy storage capacity of abandoned grassland communities on semiarid Loess Plateau. <i>Ecohydrology</i> , <b>2021</b> , 14, e2265	2.5	3
55	Effects of water and nitrogen on growth and relative competitive ability of introduced versus native C4 grass species in the semi-arid Loess Plateau of China. <i>Journal of Arid Land</i> , <b>2021</b> , 13, 730-743	2.2	
54	N and P addition increase soil respiration but decrease contribution of heterotrophic respiration in semiarid grassland. <i>Agriculture, Ecosystems and Environment</i> , <b>2021</b> , 318, 107493	5.7	2
53	Root morphology and rhizosheath acid phosphatase activity in legume and graminoid species respond differently to low phosphorus supply. <i>Rhizosphere</i> , <b>2021</b> , 19, 100391	3.5	4
52	Effects of plant diversity on semiarid grassland stability depends on functional group composition and dynamics under N and P addition. <i>Science of the Total Environment</i> , <b>2021</b> , 799, 149482	10.2	1
51	Grassland productivity and diversity changes in responses to N and P addition depend primarily on tall clonal and annual species in semiarid Loess Plateau. <i>Ecological Engineering</i> , <b>2020</b> , 145, 105727	3.9	10
50	Physiological responses of yellow-horn seedlings to high temperatures under drought condition. <i>Plant Biotechnology Reports</i> , <b>2020</b> , 14, 111-120	2.5	4
49	Maize genotypes with deep root systems tolerate salt stress better than those with shallow root systems during early growth. <i>Journal of Agronomy and Crop Science</i> , <b>2020</b> , 206, 711-721	3.9	11
48	Responses of soil respiration to rainfall pulses in a natural grassland community on the semi-arid Loess Plateau of China. <i>Catena</i> , <b>2019</b> , 178, 199-208	5.8	14
47	Surface water storage characteristics of main herbaceous species in semiarid Loess Plateau of China. <i>Ecohydrology</i> , <b>2019</b> , 12, e2145	2.5	3
46	Optimal Wheat Seeding Rate is Influenced by Cultivar-Specific Topsoil and Subsoil Root Traits. <i>Agronomy Journal</i> , <b>2019</b> , 111, 3150-3160	2.2	3
45	Dissecting root trait variability in maize genotypes using the semi-hydroponic phenotyping platform. <i>Plant and Soil</i> , <b>2019</b> , 439, 75-90	4.2	18

44	Abscisic acid and brassinolide combined application synergistically enhances drought tolerance and photosynthesis of tall fescue under water stress. <i>Scientia Horticulturae</i> , <b>2018</b> , 228, 1-9	4.1	37
43	Accumulation of N and P in the Legume in Controlled Mixtures with the Grass under Varying Water and Fertilization Conditions. <i>Frontiers in Plant Science</i> , <b>2018</b> , 9, 165	6.2	6
42	Overexpression of alfalfa Orange gene in tobacco enhances carotenoid accumulation and tolerance to multiple abiotic stresses. <i>Plant Physiology and Biochemistry</i> , <b>2018</b> , 130, 613-622	5.4	13
41	Variability in leaf wettability and surface water retention of main species in semiarid Loess Plateau of China. <i>Ecohydrology</i> , <b>2018</b> , 11, e2021	2.5	6
40	Soil Moisture Availability at Early Growth Stages Strongly Affected Root Growth of When Mixed With. <i>Frontiers in Plant Science</i> , <b>2018</b> , 9, 1050	6.2	7
39	Stress-induced expression of the sweetpotato gene IbLEA14 in poplar confers enhanced tolerance to multiple abiotic stresses. <i>Environmental and Experimental Botany</i> , <b>2018</b> , 156, 261-270	5.9	4
38	Small rainfall pulses affected leaf photosynthesis rather than biomass production of dominant species in semiarid grassland community on Loess Plateau of China. <i>Functional Plant Biology</i> , <b>2017</b> , 44, 1229-1242	2.7	11
37	Down-regulation of GIGANTEA-like genes increases plant growth and salt stress tolerance in poplar. <i>Plant Biotechnology Journal</i> , <b>2017</b> , 15, 331-343	11.6	33
36	Moderate Drought Stress Affected Root Growth and Grain Yield in Old, Modern and Newly Released Cultivars of Winter Wheat. <i>Frontiers in Plant Science</i> , <b>2017</b> , 8, 672	6.2	104
35	Transgenic poplar expressing codA exhibits enhanced growth and abiotic stress tolerance. <i>Plant Physiology and Biochemistry</i> , <b>2016</b> , 100, 75-84	5.4	28
34	N:P ratio of the grass <i>Bothriochloa ischaemum</i> mixed with the legume <i>Lespedeza davurica</i> under varying water and fertilizer supplies. <i>Plant and Soil</i> , <b>2016</b> , 400, 67-79	4.2	13
33	Eco-Physiological Responses of Dominant Species to Watering in a Natural Grassland Community on the Semi-Arid Loess Plateau of China. <i>Frontiers in Plant Science</i> , <b>2016</b> , 7, 663	6.2	15
32	Diurnal and seasonal variations of soil respiration rate under different row-spacing in a <i>Panicum virgatum</i> L. field on semi-arid Loess Plateau of China. <i>Journal of Arid Land</i> , <b>2016</b> , 8, 341-349	2.2	9
31	Overexpressing Arabidopsis ABF3 increases tolerance to multiple abiotic stresses and reduces leaf size in alfalfa. <i>Plant Physiology and Biochemistry</i> , <b>2016</b> , 109, 199-208	5.4	30
30	Transgenic poplar expressing Arabidopsis YUCCA6 exhibits auxin-overproduction phenotypes and increased tolerance to abiotic stress. <i>Plant Physiology and Biochemistry</i> , <b>2015</b> , 94, 19-27	5.4	85
29	Morphological changes in roots of <i>Bothriochloa ischaemum</i> intercropped with <i>Lespedeza davurica</i> following phosphorus application and water stress. <i>Plant Biosystems</i> , <b>2015</b> , 149, 298-306	1.6	6
28	Transgenic alfalfa plants expressing the sweetpotato Orange gene exhibit enhanced abiotic stress tolerance. <i>PLoS ONE</i> , <b>2015</b> , 10, e0126050	3.7	36
27	Overexpression of codA gene confers enhanced tolerance to abiotic stresses in alfalfa. <i>Plant Physiology and Biochemistry</i> , <b>2014</b> , 85, 31-40	5.4	42

26	Transgenic alfalfa plants expressing AtNDPK2 exhibit increased growth and tolerance to abiotic stresses. <i>Plant Physiology and Biochemistry</i> , <b>2014</b> , 84, 67-77	5.4	33
25	Does a mixture of old and modern winter wheat cultivars increase yield and water use efficiency in water-limited environments?. <i>Field Crops Research</i> , <b>2014</b> , 156, 12-21	5.5	32
24	Photosynthetic activity and efficiency of <i>Bothriochloa ischaemum</i> and <i>Lespedeza davurica</i> in mixtures across growth periods under water stress. <i>Acta Physiologiae Plantarum</i> , <b>2014</b> , 36, 1033-1044	2.6	21
23	Biomass production, relative competitive ability and water use efficiency of two dominant species in semiarid Loess Plateau under different water supply and fertilization treatments. <i>Ecological Research</i> , <b>2013</b> , 28, 781-792	1.9	9
22	Effects of Root Pruning on Non-Hydraulic Root-Sourced Signal, Drought Tolerance and Water Use Efficiency of Winter Wheat. <i>Journal of Integrative Agriculture</i> , <b>2013</b> , 12, 989-998	3.2	5
21	Two perennial legumes ( <i>Astragalus adsurgens</i> Pall. and <i>Lespedeza davurica</i> S.) adapted to semiarid environments are not as productive as lucerne ( <i>Medicago sativa</i> L.), but use less water. <i>Grass and Forage Science</i> , <b>2013</b> , 68, 469-478	2.3	18
20	Calcium regulates the cell-to-cell water flow pathway in maize roots during variable water conditions. <i>Plant Physiology and Biochemistry</i> , <b>2012</b> , 58, 212-9	5.4	14
19	Biomass allocation, relative competitive ability and water use efficiency of two dominant species in semiarid Loess Plateau under water stress. <i>Plant Science</i> , <b>2011</b> , 181, 644-51	5.3	21
18	Biomass production and relative competitiveness of a C3 legume and a C4 grass co-dominant in the semiarid Loess Plateau of China. <i>Plant and Soil</i> , <b>2011</b> , 347, 25-39	4.2	19
17	The relationship between competitive ability and yield stability in an old and a modern winter wheat cultivar. <i>Plant and Soil</i> , <b>2011</b> , 347, 7-23	4.2	40
16	Velocity and pattern of ice propagation and deep supercooling in woody stems of <i>Castanea sativa</i> , <i>Morus nigra</i> and <i>Quercus robur</i> measured by IDTA. <i>Tree Physiology</i> , <b>2010</b> , 30, 1037-45	4.2	36
15	Effect of lowering the root/shoot ratio by pruning roots on water use efficiency and grain yield of winter wheat. <i>Field Crops Research</i> , <b>2010</b> , 115, 158-164	5.5	52
14	Seasonal Root Biomass and Distribution of Switchgrass and Milk Vetch Intercropping under 2:1 Row Replacement in a Semiarid Region in Northwest China. <i>Communications in Soil Science and Plant Analysis</i> , <b>2010</b> , 41, 1959-1973	1.5	14
13	Does root pruning increase yield and water-use efficiency of winter wheat?. <i>Crop and Pasture Science</i> , <b>2010</b> , 61, 899	2.2	24
12	Seedling biomass partition and water use efficiency of switchgrass and milkvetch in monocultures and mixtures in response to various water availabilities. <i>Environmental Management</i> , <b>2010</b> , 46, 599-609	3.1	5
11	Grain yield, dry matter accumulation and remobilization, and root respiration in winter wheat as affected by seeding rate and root pruning. <i>European Journal of Agronomy</i> , <b>2010</b> , 33, 257-266	5	52
10	Effect of Root Redundancy on Grain Yield and Water Use Efficiency of Winter Wheat*. <i>Ying Yong Yu Huan Jing Sheng Wu Xue Bao = Chinese Journal of Applied and Environmental Biology</i> , <b>2010</b> , 16, 305-308		1
9	Effect of Root Pruning on Root Efficiency, Water Use and Yield of Winter Wheat*. <i>Ying Yong Yu Huan Jing Sheng Wu Xue Bao = Chinese Journal of Applied and Environmental Biology</i> , <b>2010</b> , 2009, 606-609		

8	Effects of root pruning on the growth and water use efficiency of winter wheat. <i>Plant Growth Regulation</i> , <b>2009</b> , 57, 233-241	3.2	12
7	Switchgrass and milkvetch intercropping under 2:1 row-replacement in semiarid region, northwest China: Aboveground biomass and water use efficiency. <i>European Journal of Agronomy</i> , <b>2008</b> , 28, 485-492 <sup>5</sup>		5 <sup>2</sup>
6	Effects of root pruning on competitive ability and water use efficiency in winter wheat. <i>Field Crops Research</i> , <b>2008</b> , 105, 56-63	5.5	46
5	Soil-Water Threshold Range of Chemical Signals and Drought Tolerance Was Mediated by ROS Homeostasis in Winter Wheat During Progressive Soil Drying. <i>Journal of Plant Growth Regulation</i> , <b>2008</b> , 27, 309-319	4.7	36
4	Aboveground biomass production and soil water dynamics of four leguminous forages in semiarid region, northwest China. <i>South African Journal of Botany</i> , <b>2006</b> , 72, 507-516	2.9	23
3	Gas exchange, biomass partition, and water relationships of three grass seedlings under water stress. <i>Weed Biology and Management</i> , <b>2006</b> , 6, 79-88	1.4	42
2	Hydraulic and Non-hydraulic Root-sourced Signals in Old and Modern Spring Wheat Cultivars in a Semiarid Area. <i>Journal of Plant Growth Regulation</i> , <b>2006</b> , 25, 120-136	4.7	27
1	Effect of silicon on morpho-physiological attributes, yield and cadmium accumulation in two maize genotypes with contrasting root system size and health risk assessment. <i>Plant and Soil</i> , 1	4.2	0