

# Hiroshi Okada

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/3213135/hiroshi-okada-publications-by-year.pdf>

**Version:** 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

64

papers

712

citations

14

h-index

25

g-index

79

ext. papers

951

ext. citations

4.3

avg, IF

3.63

L-index

#	Paper	IF	Citations
64	Relationship between eosinophils counts and muscle mass decline in older people with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort.. <i>Experimental Gerontology</i> , <b>2022</b> , 159, 1116-1121	4.5	71
63	Characterization of Peripheral Blood TCR in Patients with Type 1 Diabetes Mellitus by BD Rhapsody™ VDJ CDR3 Assay. <i>Cells</i> , <b>2022</b> , 11, 1623	7.9	0
62	Asymptomatic postprandial hypotension in patients with diabetes: The KAMOGAWA-HBP study. <i>Journal of Diabetes Investigation</i> , <b>2021</b> , 12, 837-844	3.9	0
61	Sarcopenia Is Associated With a Risk of Mortality in People With Type 2 Diabetes Mellitus. <i>Frontiers in Endocrinology</i> , <b>2021</b> , 12, 783363	5.7	0
60	Association between Geriatric Nutrition Risk Index and The Presence of Sarcopenia in People with Type 2 Diabetes Mellitus: A Cross-Sectional Study. <i>Nutrients</i> , <b>2021</b> , 13,	6.7	1
59	Impact of extracellular-to-intracellular fluid volume ratio on albuminuria in patients with type 2 diabetes: A cross-sectional and longitudinal cohort study. <i>Journal of Diabetes Investigation</i> , <b>2021</b> , 12, 1202-1211	3.9	1
58	Association of mean corpuscular volume with sarcopenia and visceral obesity in individuals without anemia. <i>Journal of Diabetes Investigation</i> , <b>2021</b> , 12, 1287-1292	3.9	1
57	Short energy intake is associated with muscle mass loss in older patients with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort. <i>Clinical Nutrition</i> , <b>2021</b> , 40, 1613-1620	5.9	7
56	Habitual Miso (Fermented Soybean Paste) Consumption Is Associated with Glycemic Variability in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , <b>2021</b> , 13,	6.7	1
55	Association between variability in body mass index and development of type 2 diabetes: Panasonic cohort study. <i>BMJ Open Diabetes Research and Care</i> , <b>2021</b> , 9,	4.5	2
54	Habitual Dietary Intake Affects the Altered Pattern of Gut Microbiome by Acarbose in Patients with Type 2 Diabetes. <i>Nutrients</i> , <b>2021</b> , 13,	6.7	3
53	Distinct associations of intraperitoneal and retroperitoneal visceral adipose tissues with metabolic syndrome and its components. <i>Clinical Nutrition</i> , <b>2021</b> , 40, 3479-3484	5.9	1
52	Effect of coronavirus disease 2019 pandemic on the lifestyle and glycemic control in patients with type 2 diabetes: a cross-section and retrospective cohort study. <i>Endocrine Journal</i> , <b>2021</b> , 68, 201-210	2.9	18
51	Vitamin Intake and Loss of Muscle Mass in Older People with Type 2 Diabetes: A Prospective Study of the KAMOGAWA-DM Cohort. <i>Nutrients</i> , <b>2021</b> , 13,	6.7	2
50	Decreased peripheral perfusion measured by perfusion index is a novel indicator for cardiovascular death in patients with type 2 diabetes and established cardiovascular disease. <i>Scientific Reports</i> , <b>2021</b> , 11, 2135	4.9	1
49	Let-7e-5p Regulates IGF2BP2, and Induces Muscle Atrophy.. <i>Frontiers in Endocrinology</i> , <b>2021</b> , 12, 791363	5.7	
48	Intraperitoneal, but not retroperitoneal, visceral adipose tissue is associated with diabetes mellitus: a cross-sectional, retrospective pilot analysis. <i>Diabetology and Metabolic Syndrome</i> , <b>2020</b> , 12, 103	5.6	2

47	Usefulness of Exercise for Home Blood Pressure Control in People with Diabetes: A Study Protocol for a Crossover Randomized Controlled Trial. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , <b>2020</b> , 13, 4747-4753	3.4	1
46	Low-attenuation muscle is a predictor of diabetes mellitus: A population-based cohort study. <i>Nutrition</i> , <b>2020</b> , 74, 110752	4.8	12
45	Decreased microcirculatory function measured by perfusion index is predictive of cardiovascular death. <i>Heart and Vessels</i> , <b>2020</b> , 35, 930-935	2.1	1
44	Peripheral perfusion, measured by perfusion index, is a novel indicator for renal events in patients with type 2 diabetes mellitus. <i>Scientific Reports</i> , <b>2020</b> , 10, 6054	4.9	
43	Comparison of the Efficacy of Repaglinide Versus the Combination of Mitiglinide and Voglibose on Glycemic Variability in Japanese Patients with Type 2 Diabetes. <i>Current Pharmaceutical Design</i> , <b>2020</b> , 25, 4600-4605	3.3	1
42	Habitual Miso (Fermented Soybean Paste) Consumption Is Associated with a Low Prevalence of Sarcopenia in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , <b>2020</b> , 13,	6.7	6
41	Effect of probiotics, G9-1, on gastrointestinal symptoms in patients with type 2 diabetes mellitus: study protocol for open-label, single-arm, exploratory research trial (Big STAR study). <i>Journal of Clinical Biochemistry and Nutrition</i> , <b>2020</b> , 67, 223-227	3.1	0
40	Trunk muscle quality and quantity predict the development of metabolic syndrome and the increase in the number of its components in individuals without metabolic syndrome. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , <b>2020</b> , 30, 1161-1168	4.5	4
39	Relationship between nonalcoholic fatty liver disease and muscle quality as well as quantity evaluated by computed tomography. <i>Liver International</i> , <b>2020</b> , 40, 120-130	7.9	14
38	Decreased microcirculatory function measured by perfusion index is a novel indicator of diabetic kidney disease in patients with type 2 diabetes. <i>Journal of Diabetes Investigation</i> , <b>2020</b> , 11, 681-687	3.9	2
37	Eating Fast Is Associated with Nonalcoholic Fatty Liver Disease in Men But Not in Women with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , <b>2020</b> , 12,	6.7	5
36	Combined effect of hemoglobin and mean corpuscular volume levels on incident metabolic syndrome: A population-based cohort study. <i>Clinical Nutrition ESPEN</i> , <b>2020</b> , 40, 314-319	1.3	0
35	The Effects of Metformin on the Gut Microbiota of Patients with Type 2 Diabetes: A Two-Center, Quasi-Experimental Study. <i>Life</i> , <b>2020</b> , 10,	3	9
34	Relationship between metabolic syndrome and trunk muscle quality as well as quantity evaluated by computed tomography. <i>Clinical Nutrition</i> , <b>2020</b> , 39, 1818-1825	5.9	12
33	Association between Body Weight Gain from 20 Years of Age and Diabetic Nephropathy in Japanese Patients with Type 2 Diabetes. <i>Kidney and Blood Pressure Research</i> , <b>2019</b> , 44, 583-589	3.1	
32	Frequent Usage of Convenience Stores is Associated with Low Diet Quality. <i>Nutrients</i> , <b>2019</b> , 11,	6.7	8
31	Potential impact of the joint association of total bilirubin and gamma-glutamyltransferase with metabolic syndrome. <i>Diabetology and Metabolic Syndrome</i> , <b>2019</b> , 11, 12	5.6	7
30	2457-PUB: Microcirculatory Alterations Are Associated with Increased Risk of Total Death in Patients with Type 2 Diabetes. <i>Diabetes</i> , <b>2019</b> , 68, 2457-PUB	0.9	

29	The perfusion index is a useful screening tool for peripheral artery disease. <i>Heart and Vessels</i> , <b>2019</b> , 34, 583-589	2.1	5
28	The Association Between Taste Impairment and Serum Zinc Concentration in Adult Patients With Type 2 Diabetes. <i>Canadian Journal of Diabetes</i> , <b>2018</b> , 42, 520-524	2.1	3
27	Sarcopenia is associated with blood pressure variability in older patients with type 2 diabetes: A cross-sectional study of the KAMOGAWA-DM cohort study. <i>Geriatrics and Gerontology International</i> , <b>2018</b> , 18, 1345-1349	2.9	23
26	Valacyclovir-induced Neurotoxicity in a Patient with a Preserved Renal Function. <i>Internal Medicine</i> , <b>2018</b> , 57, 3213-3216	1.1	6
25	Cerebral Salt-wasting Syndrome and Inappropriate Antidiuretic Hormone Syndrome after Subarachnoid Hemorrhaging. <i>Internal Medicine</i> , <b>2017</b> , 56, 677-680	1.1	11
24	Caffeine intake enhances the benefits of sodium glucose transporter 2 inhibitor. <i>Diabetes/Metabolism Research and Reviews</i> , <b>2016</b> , 32, 694-699	7.5	3
23	Reply to: Serum high-sensitivity cardiac troponin T and coronary artery calcification. <i>Journal of Cardiovascular Computed Tomography</i> , <b>2016</b> , 10, e3-4	2.8	
22	Postprandial hyperglycemia was ameliorated by taking metformin 30 min before a meal than taking metformin with a meal; a randomized, open-label, crossover pilot study. <i>Endocrine</i> , <b>2016</b> , 52, 271-6	4	6
21	Which Measurement of Blood Pressure Is More Associated With Albuminuria in Patients With Type 2 Diabetes: Central Blood Pressure or Peripheral Blood Pressure?. <i>Journal of Clinical Hypertension</i> , <b>2016</b> , 18, 790-5	2.3	4
20	The PR interval and QRS duration could be predictors of renal function decline. <i>Atherosclerosis</i> , <b>2015</b> , 240, 105-9	3.1	4
19	Metabolically healthy obesity and risk of incident CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , <b>2015</b> , 10, 578-83	6.9	94
18	Association between Hemoglobin Concentration and the Progression or Development of Albuminuria in Diabetic Kidney Disease. <i>PLoS ONE</i> , <b>2015</b> , 10, e0129192	3.7	3
17	High-sensitivity cardiac troponin T is associated with coronary artery calcification. <i>Journal of Cardiovascular Computed Tomography</i> , <b>2015</b> , 9, 209-14	2.8	5
16	Senescence marker protein-30/gluconolactonase deficiency exacerbates diabetic nephropathy through tubular injury in a mouse model of type 2 diabetes. <i>Journal of Diabetes Investigation</i> , <b>2015</b> , 6, 35-43	3.9	13
15	A primary diffuse large B-cell lymphoma of the liver treated with R-CHOP regimen. <i>Okayama Igakkai Zasshi</i> , <b>2015</b> , 127, 209-212	0	
14	Low serum bilirubin concentration is a predictor of chronic kidney disease. <i>Atherosclerosis</i> , <b>2014</b> , 234, 421-5	3.1	31
13	The inter-arm difference in systolic blood pressure is a novel risk marker for subclinical atherosclerosis in patients with type 2 diabetes. <i>Hypertension Research</i> , <b>2014</b> , 37, 548-52	4.7	32
12	Low serum bilirubin concentration is a novel risk factor for the development of albuminuria in patients with type 2 diabetes. <i>Metabolism: Clinical and Experimental</i> , <b>2014</b> , 63, 409-14	12.7	19

11	A difference in systolic blood pressure between arms is a novel predictor of the development and progression of diabetic nephropathy in patients with type 2 diabetes. <i>Atherosclerosis</i> , <b>2013</b> , 230, 198-204	3.1	10
10	Testosterone deficiency induces markedly decreased serum triglycerides, increased small dense LDL, and hepatic steatosis mediated by dysregulation of lipid assembly and secretion in mice fed a high-fat diet. <i>Metabolism: Clinical and Experimental</i> , <b>2013</b> , 62, 851-60	12.7	44
9	A difference in systolic blood pressure between arms and between lower limbs is a novel risk marker for diabetic nephropathy in patients with type 2 diabetes. <i>Hypertension Research</i> , <b>2013</b> , 36, 403-407	4.7	19
8	Visit-to-visit blood pressure variability is a novel risk factor for the development and progression of diabetic nephropathy in patients with type 2 diabetes. <i>Diabetes Care</i> , <b>2013</b> , 36, 1908-12	14.6	60
7	Visit-to-visit variability in systolic blood pressure is a novel risk factor for the progression of coronary artery calcification. <i>Hypertension Research</i> , <b>2013</b> , 36, 996-9	4.7	15
6	Effects of liraglutide on postprandial insulin and glucagon responses in Japanese patients with type 2 diabetes. <i>Journal of Clinical Biochemistry and Nutrition</i> , <b>2013</b> , 53, 68-72	3.1	17
5	Lepr(db/db) Mice with senescence marker protein-30 knockout (Lepr(db/db)Smp30(Y/-)) exhibit increases in small dense-LDL and severe fatty liver despite being fed a standard diet. <i>PLoS ONE</i> , <b>2013</b> , 8, e65698	3.7	19
4	Visit-to-visit variability in systolic blood pressure is correlated with diabetic nephropathy and atherosclerosis in patients with type 2 diabetes. <i>Atherosclerosis</i> , <b>2012</b> , 220, 155-9	3.1	102
3	Pancreatic insulin release in vitamin C-deficient senescence marker protein-30/gluconolactonase knockout mice. <i>Journal of Clinical Biochemistry and Nutrition</i> , <b>2012</b> , 50, 114-8	3.1	12
2	Dipeptidyl-peptidase IV inhibitor is effective in patients with type 2 diabetes with high serum eicosapentaenoic acid concentrations. <i>Journal of Diabetes Investigation</i> , <b>2012</b> , 3, 498-502	3.9	18
1	Low insulin level is associated with aortic stiffness. <i>Hypertension Research</i> , <b>2011</b> , 34, 336-40	4.7	4