Nathalie Viguerie

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/3224228/nathalie-viguerie-publications-by-year.pdf

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

87	5,958	41	77
papers	citations	h-index	g-index
88 ext. papers	6,595 ext. citations	6.6 avg, IF	4.69 L-index

#	Paper	IF	Citations
87	Network analyses reveal negative link between changes in adipose tissue GDF15 and BMI during dietary induced weight loss. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 ,	5.6	1
86	Metabolic and cardiovascular adaptations to an 8-wk lifestyle weight loss intervention in younger and older obese men. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2021 , 321, E325-E	:337	О
85	Integrative phenotyping of glycemic responders upon clinical weight loss using multi-omics. <i>Scientific Reports</i> , 2020 , 10, 9236	4.9	9
84	Growth and differentiation factor 15 is secreted by skeletal muscle during exercise and promotes lipolysis in humans. <i>JCI Insight</i> , 2020 , 5,	9.9	37
83	Apolipoprotein M: new connections with diet, adipose tissue and metabolic syndrome. <i>Current Opinion in Lipidology</i> , 2020 , 31, 8-14	4.4	4
82	Atrial Natriuretic Peptide Orchestrates a Coordinated Physiological Response to Fuel Non-shivering Thermogenesis. <i>Cell Reports</i> , 2020 , 32, 108075	10.6	8
81	Nutrients and Gene Expression in Type 2 Diabetes 2020 , 441-445		
80	Genome-wide gene-based analyses of weight loss interventions identify a potential role for NKX6.3 in metabolism. <i>Nature Communications</i> , 2019 , 10, 540	17.4	11
79	Apolipoprotein M: a novel adipokine decreasing with obesity and upregulated by calorie restriction. <i>American Journal of Clinical Nutrition</i> , 2019 , 109, 1499-1510	7	20
78	FADS1 genotype is distinguished by human subcutaneous adipose tissue fatty acids, but not inflammatory gene expression. <i>International Journal of Obesity</i> , 2019 , 43, 1539-1548	5.5	8
77	Interaction between hormone-sensitive lipase and ChREBP in fat cells controls insulin sensitivity. Nature Metabolism, 2019, 1, 133-146	14.6	26
76	Niacin induces miR-502-3p expression which impairs insulin sensitivity in human adipocytes. <i>International Journal of Obesity</i> , 2019 , 43, 1485-1490	5.5	7
75	Caloric Restriction and Diet-Induced Weight Loss Do Not Induce Browning of Human Subcutaneous White Adipose Tissue in Women and Men with Obesity. <i>Cell Reports</i> , 2018 , 22, 1079-1089	10.6	40
74	Natriuretic peptides promote glucose uptake in a cGMP-dependent manner in human adipocytes. <i>Scientific Reports</i> , 2018 , 8, 1097	4.9	22
73	Multiple hot-deck imputation for network inference from RNA sequencing data. <i>Bioinformatics</i> , 2018 , 34, 1726-1732	7.2	3
72	Plasma metabolites and lipids predict insulin sensitivity improvement in obese, nondiabetic individuals after a 2-phase dietary intervention. <i>American Journal of Clinical Nutrition</i> , 2018 , 108, 13-23	7	15
71	Molecular Biomarkers for Weight Control in Obese Individuals Subjected to a Multiphase Dietary Intervention. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017 , 102, 2751-2761	5.6	19

(2013-2017)

70	and glycemic outcomes in obese, nondiabetic subjects. <i>American Journal of Clinical Nutrition</i> , 2017 , 106, 736-746	7	36
69	Protein quantitative trait locus study in obesity during weight-loss identifies a leptin regulator. <i>Nature Communications</i> , 2017 , 8, 2084	17.4	36
68	System model network for adipose tissue signatures related to weight changes in response to calorie restriction and subsequent weight maintenance. <i>PLoS Computational Biology</i> , 2015 , 11, e100404	47⁄7	28
67	Irf5 deficiency in macrophages promotes beneficial adipose tissue expansion and insulin sensitivity during obesity. <i>Nature Medicine</i> , 2015 , 21, 610-8	50.5	130
66	MAFB as a novel regulator of human adipose tissue inflammation. <i>Diabetologia</i> , 2015 , 58, 2115-23	10.3	17
65	Immune cell Toll-like receptor 4 mediates the development of obesity- and endotoxemia-associated adipose tissue fibrosis. <i>Cell Reports</i> , 2014 , 7, 1116-29	10.6	90
64	Pro-fibrotic activity of lysophosphatidic acid in adipose tissue: in vivo and in vitro evidence. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2014 , 1841, 88-96	5	17
63	Inferring Networks from Multiple Samples with Consensus LASSO. <i>Quality Technology and Quantitative Management</i> , 2014 , 11, 39-60	1.9	5
62	Effect of endurance training on skeletal muscle myokine expression in obese men: identification of apelin as a novel myokine. <i>International Journal of Obesity</i> , 2014 , 38, 707-13	5.5	111
61	Adipose tissue CIDEA is associated, independently of weight variation, to change in insulin resistance during a longitudinal weight control dietary program in obese individuals. <i>PLoS ONE</i> , 2014 , 9, e98707	3.7	7
60	Semaphorin 3C is a novel adipokine linked to extracellular matrix composition. <i>Diabetologia</i> , 2013 , 56, 1792-801	10.3	24
59	A role for adipocyte-derived lipopolysaccharide-binding protein in inflammation- and obesity-associated adipose tissue dysfunction. <i>Diabetologia</i> , 2013 , 56, 2524-37	10.3	75
58	Contribution of Dmics Approaches to Understand the Pathophysiology of Obesity 2013, 267-281		
57	Endurance exercise training up-regulates lipolytic proteins and reduces triglyceride content in skeletal muscle of obese subjects. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, 4863-71	5.6	57
56	Partial inhibition of adipose tissue lipolysis improves glucose metabolism and insulin sensitivity without alteration of fat mass. <i>PLoS Biology</i> , 2013 , 11, e1001485	9.7	143
55	Metabolic syndrome, circulating RBP4, testosterone, and SHBG predict weight regain at 6 months after weight loss in men. <i>Obesity</i> , 2013 , 21, 1997-2006	8	20
54	Influence of SNPs in nutrient-sensitive candidate genes and gene-diet interactions on blood lipids: the DiOGenes study. <i>British Journal of Nutrition</i> , 2013 , 110, 790-6	3.6	12
53	Weight loss improves the adipogenic capacity of human preadipocytes and modulates their secretory profile. <i>Diabetes</i> , 2013 , 62, 1990-5	0.9	37

52	Circulating ACE is a predictor of weight loss maintenance not only in overweight and obese women, but also in men. <i>International Journal of Obesity</i> , 2012 , 36, 1545-51	5.5	31
51	Determinants of human adipose tissue gene expression: impact of diet, sex, metabolic status, and cis genetic regulation. <i>PLoS Genetics</i> , 2012 , 8, e1002959	6	41
50	Multiple effects of a short-term dexamethasone treatment in human skeletal muscle and adipose tissue. <i>Physiological Genomics</i> , 2012 , 44, 141-51	3.6	17
49	Analyses of single nucleotide polymorphisms in selected nutrient-sensitive genes in weight-regain prevention: the DIOGENES study. <i>American Journal of Clinical Nutrition</i> , 2012 , 95, 1254-60	7	32
48	Natriuretic peptides enhance the oxidative capacity of human skeletal muscle. <i>Journal of Clinical Investigation</i> , 2012 , 122, 4675-9	15.9	127
47	Macrophage gene expression is related to obesity and the metabolic syndrome in human subcutaneous fat as well as in visceral fat. <i>Diabetologia</i> , 2011 , 54, 876-87	10.3	60
46	A distinct adipose tissue gene expression response to caloric restriction predicts 6-mo weight maintenance in obese subjects. <i>American Journal of Clinical Nutrition</i> , 2011 , 94, 1399-409	7	48
45	Worsening of obesity and metabolic status yields similar molecular adaptations in human subcutaneous and visceral adipose tissue: decreased metabolism and increased immune response. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011 , 96, E73-82	5.6	82
44	Impact of a mechanical massage on gene expression profile and lipid mobilization in female gluteofemoral adipose tissue. <i>Obesity Facts</i> , 2011 , 4, 121-9	5.1	19
43	Peroxisome proliferator-activated receptor-alpha control of lipid and glucose metabolism in human white adipocytes. <i>Endocrinology</i> , 2010 , 151, 123-33	4.8	79
42	Adipose tissue transcriptome reflects variations between subjects with continued weight loss and subjects regaining weight 6 mo after caloric restriction independent of energy intake. <i>American Journal of Clinical Nutrition</i> , 2010 , 92, 975-84	7	52
41	Macrophages and adipocytes in human obesity: adipose tissue gene expression and insulin sensitivity during calorie restriction and weight stabilization. <i>Diabetes</i> , 2009 , 58, 1558-67	0.9	142
40	Visfatin expression in subcutaneous adipose tissue of pre-menopausal women: relation to hormones and weight reduction. <i>European Journal of Clinical Investigation</i> , 2008 , 38, 516-22	4.6	20
39	Adipose tissue transcriptomic signature highlights the pathological relevance of extracellular matrix in human obesity. <i>Genome Biology</i> , 2008 , 9, R14	18.3	300
38	Contribution of energy restriction and macronutrient composition to changes in adipose tissue gene expression during dietary weight-loss programs in obese women. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008 , 93, 4315-22	5.6	64
37	Gene expression profiling of human skeletal muscle in response to stabilized weight loss. <i>American Journal of Clinical Nutrition</i> , 2008 , 88, 125-32	7	24
36	Adipose tissue 2008 , 51-58		
35	Adipose gene expression prior to weight loss can differentiate and weakly predict dietary responders. <i>PLoS ONE</i> , 2007 , 2, e1344	3.7	42

(2004-2007)

34	An intervention study of the effects of calcium intake on faecal fat excretion, energy metabolism and adipose tissue mRNA expression of lipid-metabolism related proteins. <i>International Journal of Obesity</i> , 2007 , 31, 1704-12	5.5	67
33	Plasma levels and adipose tissue messenger ribonucleic acid expression of retinol-binding protein 4 are reduced during calorie restriction in obese subjects but are not related to diet-induced changes in insulin sensitivity. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007 , 92, 2330-5	5.6	82
32	The transcriptional coactivator peroxisome proliferator activated receptor (PPAR)gamma coactivator-1 alpha and the nuclear receptor PPAR alpha control the expression of glycerol kinase and metabolism genes independently of PPAR gamma activation in human white adipocytes. Diabetes, 2007, 56, 2467-75	0.9	70
31	Profiling of adipokines secreted from human subcutaneous adipose tissue in response to PPAR agonists. <i>Biochemical and Biophysical Research Communications</i> , 2007 , 358, 897-902	3.4	28
30	Dynamic strength training improves insulin sensitivity without altering plasma levels and gene expression of adipokines in subcutaneous adipose tissue in obese men. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2006 , 91, 5107-12	5.6	111
29	Effect of aerobic training on plasma levels and subcutaneous abdominal adipose tissue gene expression of adiponectin, leptin, interleukin 6, and tumor necrosis factor alpha in obese women. <i>Metabolism: Clinical and Experimental</i> , 2006 , 55, 1375-81	12.7	150
28	Role of adipokines in the control of energy metabolism: focus on adiponectin. <i>Current Opinion in Pharmacology</i> , 2006 , 6, 580-5	5.1	70
27	Microarray profiling of human white adipose tissue after exogenous leptin injection. <i>European Journal of Clinical Investigation</i> , 2006 , 36, 153-63	4.6	19
26	The effects of increasing serum calcitriol on energy and fat metabolism and gene expression. <i>Obesity</i> , 2006 , 14, 1739-46	8	41
25	Transcriptomics applied to obesity and caloric restriction. <i>Biochimie</i> , 2005 , 87, 117-23	4.6	57
24	Reduction of macrophage infiltration and chemoattractant gene expression changes in white adipose tissue of morbidly obese subjects after surgery-induced weight loss. <i>Diabetes</i> , 2005 , 54, 2277-8	36 ^{0.9}	870
23	Adipose tissue gene expression in obese subjects during low-fat and high-fat hypocaloric diets. <i>Diabetologia</i> , 2005 , 48, 123-31	10.3	113
22	Serum amyloid A: production by human white adipocyte and regulation by obesity and nutrition. <i>Diabetologia</i> , 2005 , 48, 519-28	10.3	139
21	Dynamic strength training improves insulin sensitivity and functional balance between adrenergic alpha 2A and beta pathways in subcutaneous adipose tissue of obese subjects. <i>Diabetologia</i> , 2005 , 48, 2631-40	10.3	37
20	Effects of 3 diets with various calcium contents on 24-h energy expenditure, fat oxidation, and adipose tissue message RNA expression of lipid metabolism-related proteins. <i>American Journal of Clinical Nutrition</i> , 2005 , 82, 1244-52	7	37
19	Cathepsin S, a novel biomarker of adiposity: relevance to atherogenesis. FASEB Journal, 2005, 19, 1540-	-2 0.9	119
18	Adipocyte lipases and defect of lipolysis in human obesity. <i>Diabetes</i> , 2005 , 54, 3190-7	0.9	293
17	Effect of endurance training on adrenergic control of lipolysis in adipose tissue of obese women. Journal of Clinical Endocrinology and Metabolism, 2004 , 89, 1325-31	5.6	37

16	Adiponutrin: A new gene regulated by energy balance in human adipose tissue. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2004 , 89, 2684-9	5.6	79
15	Effects of different hypocaloric diets on protein secretion from adipose tissue of obese women. <i>Diabetes</i> , 2004 , 53, 1966-71	0.9	112
14	Weight loss regulates inflammation-related genes in white adipose tissue of obese subjects. <i>FASEB Journal</i> , 2004 , 18, 1657-69	0.9	506
13	Adiponectin gene expression and plasma values in obese women during very-low-calorie diet. Relationship with cardiovascular risk factors and insulin resistance. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2004 , 89, 756-60	5.6	60
12	In vivo epinephrine-mediated regulation of gene expression in human skeletal muscle. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2004 , 89, 2000-14	5.6	48
11	. Current Opinion in Clinical Nutrition and Metabolic Care, 2003 , 6, 377-381	3.8	8
10	Effect of thyroid hormone on gene expression. <i>Current Opinion in Clinical Nutrition and Metabolic Care</i> , 2003 , 6, 377-81	3.8	37
9	Semicarbazide-sensitive amine oxidase/vascular adhesion protein-1 activity exerts an antidiabetic action in Goto-Kakizaki rats. <i>Diabetes</i> , 2003 , 52, 1004-13	0.9	52
8	Adiponectin gene expression in subcutaneous adipose tissue of obese women in response to short-term very low calorie diet and refeeding. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2003 , 88, 5881-6	5.6	56
7	Regulation of human adipocyte gene expression by thyroid hormone. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2002 , 87, 630-4	5.6	79
6	In vivo regulation of human skeletal muscle gene expression by thyroid hormone. <i>Genome Research</i> , 2002 , 12, 281-91	9.7	124
5	Hypocaloric Diet Reduces Exercise-Induced 12-Adrenergic Antilipolytic Effect and 12-Adrenergic Receptor mRNA Levels in Adipose Tissue of Obese Women. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2002 , 87, 1274-1281	5.6	25
4	Triiodothyronine-mediated up-regulation of UCP2 and UCP3 mRNA expression in human skeletal muscle without coordinated induction of mitochondrial respiratory chain genes. <i>FASEB Journal</i> , 2001 , 15, 13-15	0.9	87
3	Increase in uncoupling protein-2 mRNA expression by BRL49653 and bromopalmitate in human adipocytes. <i>Biochemical and Biophysical Research Communications</i> , 1999 , 256, 138-41	3.4	48
2	Dexamethasone effects on somatostatin receptors in pancreatic acinar AR4-2J cells. <i>Biochemical and Biophysical Research Communications</i> , 1987 , 147, 942-8	3.4	30
1	Solubilization and characterization of guinea-pig pancreatic somatostatin receptors. <i>FEBS Journal</i> , 1987 , 164, 667-73		11