A Nedret Koc

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3227092/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Antifungal Activity of the Honeybee Products Against <i>Candida</i> spp. and <i>Trichosporon</i> spp Journal of Medicinal Food, 2011, 14, 128-134.	1.5	61
2	Antifungal Activity of Turkish Honey against <i>Candida</i> spp. and <i>Trichosporon</i> spp: an <i>in vitro</i> evaluation. Medical Mycology, 2009, 47, 707-712.	0.7	51
3	Investigation of the relationship between virulence factors and genotype of Candida spp. isolated from blood cultures. Journal of Infection in Developing Countries, 2015, 9, 857-864.	1.2	29
4	Recent Increase in the Prevalence of Fluconazole-Non-susceptible Candida tropicalis Blood Isolates in Turkey: Clinical Implication of Azole-Non-susceptible and Fluconazole Tolerant Phenotypes and Genotyping. Frontiers in Microbiology, 2020, 11, 587278.	3.5	21
5	Outbreak of nosocomial fungemia caused byCandida glabrata. Mycoses, 2002, 45, 470-475.	4.0	19
6	Comparison of Etest with the broth microdilution method in susceptibility testing of yeast isolates against four antifungals. Mycoses, 2000, 43, 293-297.	4.0	16
7	Antifungal Activity of Olive Oil and Ozonated Olive Oil Against <i>Candida</i> Spp. and <i>Saprochaete</i> Spp Ozone: Science and Engineering, 2017, 39, 462-470.	2.5	15
8	Molecular epidemiology and antifungal susceptibility of Saprochaete capitata (Blastoschizomyces) Tj ETQq0 0 0 r 596-603.	gBT /Overl 2.8	lock 10 Tf 50 10
9	Conventional Morphology Versus PCR Sequencing, repâ€PCR, and MALDIâ€TOFâ€MS for Identification of Clinical <i>Aspergillus</i> Isolates Collected Over a 2â€Year Period in a University Hospital at Kayseri, Turkey. Journal of Clinical Laboratory Analysis, 2016, 30, 745-750.	2.1	9
10	Genotypes and virulence factors of Candida species isolated from oralcavities of patients with type 2 diabetes mellitus. Turkish Journal of Medical Sciences, 2016, 46, 18-27.	0.9	7
11	Mucormycosis experience through the eyes of the laboratory. Infectious Diseases, 2019, 51, 730-737.	2.8	7
12	Gamma-glutamyl transpeptidase–platelet ratio, systemic immune inflammation index, and system inflammation response index in invasive Aspergillosis. Revista Da Associação Médica Brasileira, 2021, 67, 1021-1025.	0.7	7
13	Prevalence and risk factors of tinea capitis and tinea pedis in school children in Turkey. JPMA the Journal of the Pakistan Medical Association, 2014, 64, 514-8.	0.2	6
14	Molecular epidemiology, antifungal susceptibility and virulence factors of Candida glabrata complex strains in Kayseri/Turkey. Microbial Pathogenesis, 2021, 154, 104870.	2.9	5
15	Identification and molecular epidemiology of dermatophyte isolates by repetitiveâ€sequenceâ€ <scp>PCR</scp> â€based <scp>DNA</scp> fingerprinting using the DiversiLab system in Turkey. Mycoses, 2017, 60, 348-354.	4.0	3
16	Usefulness of (1→3)β-D glucan in early diagnosing Pneumocystis jirovecii pneumonia: a case report. Infezioni in Medicina, 2014, 22, 57-61.	1.1	2
17	Utility of the Aspergillus galactomannan antigen testing for neutropenic paediatric patients. Infezioni in Medicina, 2017, 25, 38-44.	1.1	2
18	Studies on the Effectiveness of Ozone Therapy on the Treatment of Experimentally Induced Keratitis with Candida albicans in Rabbits. Seminars in Ophthalmology, 2021, , 1-12.	1.6	0