Chung-Ping Yu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3229663/publications.pdf

Version: 2024-02-01

759233 794594 21 367 12 19 citations h-index g-index papers 21 21 21 609 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Bidirectional Influences of Cranberry on the Pharmacokinetics and Pharmacodynamics of Warfarin with Mechanism Elucidation. Nutrients, 2021, 13, 3219.	4.1	4
2	Folium Sennae Increased the Bioavailability of Methotrexate through Modulation on MRP 2 and BCRP. Pharmaceuticals, 2021, 14, 1036.	3.8	2
3	Magnolol and Honokiol Inhibited the Function and Expression of BCRP with Mechanism Exploration. Molecules, 2021, 26, 7390.	3.8	10
4	Effects of antibiotics on the pharmacokinetics of indoxyl sulfate, a nephro-cardiovascular toxin. Xenobiotica, 2020, 50, 588-592.	1.1	6
5	Resveratrol stereoselectively affected ($\hat{A}\pm$)warfarin pharmacokinetics and enhanced the anticoagulation effect. Scientific Reports, 2020, 10, 15910.	3.3	10
6	Activation of P-glycoprotein and CYP 3A by Coptidis Rhizoma inÂvivo: Using cyclosporine as a probe substrate in rats. Journal of Food and Drug Analysis, 2018, 26, S125-S132.	1.9	25
7	Effects of nonsteroidal anti-inflammatory drugs on the renal excretion of indoxyl sulfate, a nephro-cardiovascular toxin, in rats. European Journal of Pharmaceutical Sciences, 2017, 101, 66-70.	4.0	20
8	Aloe activated P-glycoprotein and CYP 3A: a study on the serum kinetics of aloe and its interaction with cyclosporine in rats. Food and Function, 2017, 8, 315-322.	4.6	17
9	Increased Systemic Exposure of Methotrexate by a Polyphenol-Rich Herb via Modulation on Efflux Transporters Multidrug Resistance–Associated Protein 2 and Breast Cancer Resistance Protein. Journal of Pharmaceutical Sciences, 2016, 105, 343-349.	3.3	26
10	Analysis of the pharmacokinetics and metabolism of aloeâ€emodin following intravenous and oral administrations in rats. Biomedical Chromatography, 2016, 30, 1641-1647.	1.7	19
11	Rhubarb decreased the systemic exposure of cyclosporine, a probe substrate of P-glycoprotein and CYP 3A. Xenobiotica, 2016, 46, 677-682.	1.1	14
12	Green tea inhibited the elimination of nephro-cardiovascular toxins and deteriorated the renal function in rats with renal failure. Scientific Reports, 2015, 5, 16226.	3.3	14
13	Oral intake of curcumin markedly activated CYP 3A4: in vivo and ex-vivo studies. Scientific Reports, 2015, 4, 6587.	3.3	39
14	Serum Concentrations of Anthraquinones after Intake of Folium Sennae and Potential Modulation on P-glycoprotein. Planta Medica, 2014, 80, 1291-1297.	1.3	11
15	Potential modulation on BCRP and MRP 4 by onion: in vivo and ex-vivo studies. Journal of Functional Foods, 2014, 8, 243-251.	3.4	3
16	A Chinese herb formula decreases the monocarboxylate transporter-mediated absorption of valproic acid in rats. Phytomedicine, 2013, 20, 648-653.	5.3	2
17	Inhibition of Monocarboxylate Transporter-Mediated Absorption of Valproic Acid by Gegen-Qinlian-Tang. The American Journal of Chinese Medicine, 2013, 41, 369-378.	3.8	8
18	Pharmacokinetics and Relative Bioavailability of Flavonoids between Two Dosage Forms of Gegen-Qinlian-Tang in Rats. Evidence-based Complementary and Alternative Medicine, 2012, 2012, 1-8.	1.2	20

#	Article	lF	CITATIONS
19	Quercetin and Rutin Reduced the Bioavailability of Cyclosporine from Neoral, an Immunosuppressant, through Activating P-Glycoprotein and CYP 3A4. Journal of Agricultural and Food Chemistry, 2011, 59, 4644-4648.	5.2	83
20	Comparison of Puerariae Radix and Its Hydrolysate on Stimulation of Hyaluronic Acid Production in NHEK Cells. The American Journal of Chinese Medicine, 2010, 38, 143-155.	3.8	18
21	Metabolic Transformation of Sesamol and ex Vivo Effect on 2,2′-Azo-bis(2-amidinopropane)dihydrochloride-Induced Hemolysis. Journal of Agricultural and Food Chemistry, 2008, 56, 9636-9640.	5.2	16