Fangxu Tang

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/3239780/fangxu-tang-publications-by-year.pdf
Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

19 papers	273 citations	8 h-index	16 g-index
19 ext. papers	323 ext. citations	4 avg, IF	3.11 L-index

#	Paper	IF	Citations
19	6 Circulating miRNAs can be used as Non-invasive Biomarkers for the Detection of Cervical Lesions. *Journal of Cancer*, **2021**, 12, 5106-5113	4.5	1
18	Risk Factors for Mortality in 244 Older Adults With COVID-19 in Wuhan, China: A Retrospective Study. *Journal of the American Geriatrics Society*, **2020**, 68, E19-E23	5.6	115
17	Comparison of clinical and microbiological diagnoses for older adults with COVID-19 in Wuhan: a retrospective study. *Aging Clinical and Experimental Research*, **2020**, 32, 1889-1895	4.8	1
16	A case of SLE with COVID-19 and multiple infections. *Open Medicine (Poland)*, **2020**, 15, 1054-1060	2.2	6
15	Reply from Sun et al. *Journal of the American Geriatrics Society*, **2020**, 68, 2195-2196	5.6	
14	Adjuvant chemotherapy after surgery can improve clinical outcomes for patients with IB2-IIB cervical cancer with neoadjuvant chemotherapy followed by radical surgery. *Scientific Reports*, **2018**, 8, 6443	4.9	8
13	Genome-wide association study identifies four SNPs associated with response to platinum-based neoadjuvant chemotherapy for cervical cancer. *Scientific Reports*, **2017**, 7, 41103	4.9	11
12	Whole-exome sequencing reveals genetic variants in ERC1 and KCNG4 associated with complete hydatidiform mole in Chinese Han women. *Oncotarget*, **2017**, 8, 75264-75271	3.3	1
11	Young Cervical Cancer Patients May Be More Responsive than Older Patients to Neoadjuvant Chemotherapy Followed by Radical Surgery. *PLoS ONE*, **2016**, 11, e0149534	3.7	4
10	Risk model in stage IB1-IIB cervical cancer with positive node after radical hysterectomy. *OncoTargets and Therapy*, **2016**, 9, 3171-9	4.4	4
9	Development and validation of a surgical-pathologic staging and scoring system for cervical cancer. *Oncotarget*, **2016**, 7, 21054-63	3.3	6
8	Association between vascular endothelial growth factor expression and lymph node metastasis in cervical cancer: A meta-analysis. *Journal of Obstetrics and Gynaecology Research*, **2016**, 42, 1310-1316	1.9	9
7	Prognostic risk model development and prospective validation among patients with cervical cancer stage IB2 to IIB submitted to neoadjuvant chemotherapy. *Scientific Reports*, **2016**, 6, 27568	4.9	3
6	Optimal pathological response indicated better long-term outcome among patients with stage IB2 to IIB cervical cancer submitted to neoadjuvant chemotherapy. *Scientific Reports*, **2016**, 6, 28278	4.9	9
5	Association of 42 SNPs with genetic risk for cervical cancer: an extensive meta-analysis. *BMC Medical Genetics*, **2015**, 16, 25	2.1	12
4	The predictive value of serum squamous cell carcinoma antigen in patients with cervical cancer who receive neoadjuvant chemotherapy followed by radical surgery: a single-institute study. *PLoS ONE*, **2015**, 10, e0122361	3.7	20
3	Could the extent of lymphadenectomy be modified by neoadjuvant chemotherapy in cervical cancer? A large-scale retrospective study. *PLoS ONE*, **2015**, 10, e0123539	3.7	2

2	Prospective cohort study to evaluate the efficacy of taxane plus platinum and CPT-11plus platinum regimes and to identify prognostic risk factors in cervical cancer patients. *International Journal of Clinical and Experimental Medicine*, **2015**, 8, 15018-29		1
1	Changes in prevalence and clinical characteristics of cervical cancer in the Peopleæs Republic of China: a study of 10,012 cases from a nationwide working group. *Oncologist*, **2013**, 18, 1101-7	5.7	60