

Ming-Li He

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3244223/publications.pdf>

Version: 2024-02-01

49
papers

1,457
citations

516710

16
h-index

330143

37
g-index

50
all docs

50
docs citations

50
times ranked

1910
citing authors

#	ARTICLE	IF	CITATIONS
1	Blood Pressure Rhythm and Blood Pressure Variability as Risk Factors for White Matter Lesions: A Cross-Sectional Study. <i>Medical Science Monitor</i> , 2022, 28, e933880.	1.1	1
2	Blood pressure undulation of peripheral thrombolysis period in acute ischemic stroke is associated with prognosis. <i>Journal of Hypertension</i> , 2022, 40, 749-757.	0.5	4
3	Optimal blood pressure levels in different phases of peripheral thrombolysis period in acute ischemic stroke. <i>Journal of Hypertension</i> , 2021, 39, 1453-1461.	0.5	4
4	Association of Blood Pressure, White Matter Lesions, and Regional Cerebral Blood Flow. <i>Medical Science Monitor</i> , 2021, 27, e929958.	1.1	5
5	AKR1C3 and Its Transcription Factor HOXB4 Are Promising Diagnostic Biomarkers for Acute Myocardial Infarction. <i>Frontiers in Cardiovascular Medicine</i> , 2021, 8, 694238.	2.4	12
6	Focus on blood pressure levels and variability in the early phase of acute ischemic stroke with hypertension and carotid stenosis. <i>Journal of Clinical Hypertension</i> , 2021, 23, 2089-2099.	2.0	3
7	Electrocardiographic-left ventricular hypertrophy and incident stroke among Chinese hypertensive adults. <i>Journal of Human Hypertension</i> , 2020, 34, 286-292.	2.2	2
8	The role of carotid stenosis ultrasound scale in the prediction of ischemic stroke. <i>Neurological Sciences</i> , 2020, 41, 1193-1199.	1.9	10
9	A U-shaped association between serum betaine and incident risk of first ischemic stroke in hypertensive patients. <i>Clinical Nutrition</i> , 2020, 39, 2517-2524.	5.0	11
10	Higher uric acid is associated with better discharge recovery and short-term outcome in stroke patients treated with thrombolysis. <i>Neurological Sciences</i> , 2020, 42, 3225-3231.	1.9	8
11	Age and Diastolic Blood Pressure Play an Important Role in the Progression of White Matter Lesions: A Meta-Analysis. <i>European Neurology</i> , 2020, 83, 351-359.	1.4	6
12	Clinical and imaging features of reversible splenial lesion syndrome with language disorder. <i>Translational Neuroscience</i> , 2020, 11, 210-214.	1.4	1
13	<i>MTHFR</i> Gene and Serum Folate Interaction on Serum Homocysteine Lowering. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2018, 38, 679-685.	2.4	29
14	Effect of Smoking and Folate Levels on the Efficacy of Folic Acid Therapy in Prevention of Stroke in Hypertensive Men. <i>Stroke</i> , 2018, 49, 114-120.	2.0	13
15	Impact of achieved blood pressure on renal function decline and first stroke in hypertensive patients with chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, 409-417.	0.7	18
16	Effect of long-term low-dose folic acid supplementation on degree of total homocysteine-lowering: major effect modifiers. <i>British Journal of Nutrition</i> , 2018, 120, 1122-1130.	2.3	9
17	Serum Trimethylamine N-Oxide Concentration Is Positively Associated With First Stroke in Hypertensive Patients. <i>Stroke</i> , 2018, 49, 2021-2028.	2.0	119
18	Blood pressures immediately following ischemic strokes are associated with cerebral perfusion and neurologic function. <i>Journal of Clinical Hypertension</i> , 2018, 20, 1008-1015.	2.0	9

#	ARTICLE	IF	CITATIONS
19	Corpus callosum infarction with cognitive dysfunction: two case reports and literature review. <i>Neuropsychiatric Disease and Treatment</i> , 2018, Volume 14, 511-515.	2.2	5
20	Community-based stroke system of care improves patient outcomes in Chinese rural areas. <i>Journal of Epidemiology and Community Health</i> , 2018, 72, 630-635.	3.7	5
21	Effects of Blood Pressure in the Early Phase of Ischemic Stroke and Stroke Subtype on Poststroke Cognitive Impairment. <i>Stroke</i> , 2018, 49, 1610-1617.	2.0	23
22	Antihypertensive Therapy in the Acute Phase of Lacunar Infarcts. <i>Medical Science Monitor</i> , 2018, 24, 6688-6694.	1.1	1
23	Folic acid therapy reduces serum uric acid in hypertensive patients: a substudy of the China Stroke Primary Prevention Trial (CSPPT). <i>American Journal of Clinical Nutrition</i> , 2017, 105, 882-889.	4.7	43
24	Homocysteine and all-cause mortality in hypertensive adults without pre-existing cardiovascular conditions. <i>Medicine (United States)</i> , 2017, 96, e5862.	1.0	20
25	Mid-Term Blood Pressure Variability Is Associated With Clinical Outcome After Ischemic Stroke. <i>American Journal of Hypertension</i> , 2017, 30, 968-977.	2.0	21
26	Prospective study of serum uric acid levels and stroke in a Chinese hypertensive cohort. <i>Clinical and Experimental Hypertension</i> , 2017, 39, 527-531.	1.3	9
27	Effects of Folic Acid Therapy on the New-Onset Proteinuria in Chinese Hypertensive Patients. <i>Hypertension</i> , 2017, 70, 300-306.	2.7	15
28	Homocysteine and Stroke Risk. <i>Stroke</i> , 2017, 48, 1183-1190.	2.0	89
29	Impact of Achieved Blood Pressure on First Stroke in Uncomplicated Grade 1 Hypertension. <i>Journal of the American Heart Association</i> , 2017, 6, .	3.7	6
30	Higher Adiposity Is Associated With Slower Cognitive Decline in Hypertensive Patients: Secondary Analysis of the China Stroke Primary Prevention Trial. <i>Journal of the American Heart Association</i> , 2017, 6, .	3.7	11
31	Association between percent decline in serum total homocysteine and risk of first stroke. <i>Neurology</i> , 2017, 89, 2101-2107.	1.1	60
32	Midterm Blood Pressure Variability Is Associated with Poststroke Cognitive Impairment: A Prospective Cohort Study. <i>Frontiers in Neurology</i> , 2017, 8, 365.	2.4	12
33	Distinct Patterns of Cognitive Aging Modified by Education Level and Gender among Adults with Limited or No Formal Education: A Normative Study of the Mini-Mental State Examination. <i>Journal of Alzheimer's Disease</i> , 2016, 49, 961-969.	2.6	18
34	Effect of folic acid supplementation on risk of new-onset diabetes in adults with hypertension in China: Findings from the China Stroke Primary Prevention Trial (CSPPT). <i>Journal of Diabetes</i> , 2016, 8, 286-294.	1.8	32
35	Folic Acid Therapy Reduces the First Stroke Risk Associated With Hypercholesterolemia Among Hypertensive Patients. <i>Stroke</i> , 2016, 47, 2805-2812.	2.0	59
36	A Reasonable Blood Pressure Level for Good Clinical Outcome After the Acute Phase of Ischemic Stroke. <i>Journal of Clinical Hypertension</i> , 2016, 18, 536-542.	2.0	7

#	ARTICLE	IF	CITATIONS
37	Proteinuria Is an Independent Risk Factor for First Incident Stroke in Adults Under Treatment for Hypertension in China. <i>Journal of the American Heart Association</i> , 2015, 4, .	3.7	15
38	Prestroke glycemic status is associated with the functional outcome in spontaneous intracerebral hemorrhage. <i>Neurological Sciences</i> , 2015, 36, 927-934.	1.9	18
39	Efficacy of Folic Acid Therapy in Primary Prevention of Stroke Among Adults With Hypertension in China. <i>JAMA - Journal of the American Medical Association</i> , 2015, 313, 1325.	7.4	577
40	Community-Based Stroke System of Care for Chinese Rural Areas. <i>Stroke</i> , 2014, 45, 2385-2390.	2.0	7
41	Effects of Organized Education by Video on Knowledge of Early Stroke Symptoms among a Chinese Rural Population. <i>International Journal of Stroke</i> , 2014, 9, E41-E41.	5.9	1
42	Prevalence of obesity, abdominal obesity and associated factors in hypertensive adults aged 45-75 years. <i>Clinical Nutrition</i> , 2013, 32, 361-367.	5.0	32
43	Prevalence of hyperhomocysteinaemia and its major determinants in rural Chinese hypertensive patients aged 45-75 years. <i>British Journal of Nutrition</i> , 2013, 109, 1284-1293.	2.3	45
44	Methylenetetrahydrofolate reductase C677T polymorphism is associated with estimated glomerular filtration rate in hypertensive Chinese males. <i>BMC Medical Genetics</i> , 2012, 13, 74.	2.1	7
45	Prevalence of Unrecognized Lower Extremity Peripheral Arterial Disease and the Associated Factors in Chinese Hypertensive Adults. <i>American Journal of Cardiology</i> , 2012, 110, 1692-1698.	1.6	28
46	Treatment of depression using sleep electroencephalogram modulated repetitive transcranial magnetic stimulation. <i>Chinese Medical Journal</i> , 2011, 124, 1779-83.	2.3	3
47	Effects of repetitive transcranial magnetic stimulation on hypothalamic-pituitary-adrenal axis of patients with depression. <i>Journal of Medical Colleges of PLA</i> , 2009, 24, 337-345.	0.1	7
48	An HPLC method for the determination of ng mifepristone in human plasma. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2006, 832, 181-184.	2.3	13
49	Red blood cell distribution width in different time-points of peripheral thrombolysis period in acute ischemic stroke is associated with prognosis. <i>Aging</i> , 0, , .	3.1	4