## Kei Nakajima

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3251735/publications.pdf

Version: 2024-02-01

		1683354	1125271	
18	184	5	13	
papers	citations	h-index	g-index	
23	23	23	455	
all docs	docs citations	times ranked	citing authors	

#	Article	IF	Citations
1	High Aspartate Aminotransferase/Alanine Aminotransferase Ratio May Be Associated with All-Cause Mortality in the Elderly: A Retrospective Cohort Study Using Artificial Intelligence and Conventional Analysis. Healthcare (Switzerland), 2022, 10, 674.	1.0	4
2	Elevated All-Cause Mortality among Overweight Older People: Al Predicts a High Normal Weight Is Optimal. Geriatrics (Switzerland), 2022, 7, 68.	0.6	1
3	Association between extremely high high-density lipoprotein-cholesterol and hypertensive retinopathy: results of a cross-sectional study from Kanagawa Investigation of Total Checkup Data from the National Database-6 (KITCHEN-6). BMJ Open, 2021, 11, e043677.	0.8	3
4	Potential Role of Artificial Intelligence for the Previous Study Using Traditional Analysis. Journal of Clinical Medicine Research, 2021, 13, 409-411.	0.6	0
5	Unexpectedly High Prevalence of Breakfast Skipping in Low Body-Weight Middle-Aged Men: Results of the Kanagawa Investigation of Total Checkup Data from the National Data Base-7 (KITCHEN-7). Nutrients, 2021, 13, 102.	1.7	2
6	Association of Serum High-Density Lipoprotein Cholesterol with High Blood Pressures at Checkup: Results of Kanagawa Investigation of Total Checkup Data from the National Database-9 (KITCHEN-9). Journal of Clinical Medicine, 2021, 10, 5118.	1.0	4
7	High Serum Aspartate Aminotransferase, Underweight, and Weight Loss in Older People: Results of the KITCHEN-4. Healthcare (Switzerland), 2020, 8, 69.	1.0	2
8	The association of low serum salivary and pancreatic amylases with the increased use of lipids as an energy source in non-obese healthy women. BMC Research Notes, 2020, 13, 237.	0.6	6
9	High Concentration of Serum Aspartate Aminotransferase in Older Underweight People: Results of the Kanagawa Investigation of the Total Check-Up Data from the National Database-2 (KITCHEN-2). Journal of Clinical Medicine, 2019, 8, 1282.	1.0	11
10	High Incidence of Diabetes in People with Extremely High High-Density Lipoprotein Cholesterol: Results of the Kanagawa Investigation of Total Checkup Data from the National Database-1 (KITCHEN-1). Journal of Clinical Medicine, 2019, 8, 381.	1.0	6
11	Impaired Glucose Metabolism in People with Extremely Elevated High-Density Lipoprotein Cholesterol and Low Alcohol Consumption: Results of the Kanagawa Investigation of Total Checkup Data from the National Database-3 (KITCHEN-3). Journal of Clinical Medicine, 2019, 8, 1825.	1.0	2
12	Adjustment for Waist Circumference Reveals a U-Shaped Association Between Glycated Hemoglobin Levels and Body Mass Index in Young Adults. Canadian Journal of Diabetes, 2019, 43, 201-206.	0.4	2
13	Unexpected association between subclinical hearing loss and restorative sleep in a middle-aged and elderly Japanese population. BMC Research Notes, 2018, 11, 195.	0.6	3
14	Ketonuria may be associated with low serum amylase independent of body weight and glucose metabolism. Archives of Physiology and Biochemistry, 2017, 123, 293-296.	1.0	5
15	Association between positive fecal occult blood test and diabetes in a population undergoing health screening. Clinical Biochemistry, 2017, 50, 97-100.	0.8	8
16	Potential Association between Breakfast Skipping and Concomitant Late-Night-Dinner Eating with Metabolic Syndrome and Proteinuria in the Japanese Population. Scientifica, 2014, 2014, 1-9.	0.6	99
17	Low Vital Capacity and Electrocardiographic ST-T Abnormalities in Asymptomatic Adults. Pulmonary Medicine, 2012, 2012, 1-7.	0.5	1
18	Coffee Consumption and Cystatin-C-Based Estimated Glomerular Filtration Rates in Healthy Young Adults: Results of a Clinical Trial. Journal of Nutrition and Metabolism, 2011, 2011, 1-7.	0.7	24