

Luis A Videla

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3251888/publications.pdf>

Version: 2024-02-01

141
papers

7,341
citations

47006

47
h-index

62596

80
g-index

142
all docs

142
docs citations

142
times ranked

7867
citing authors

#	ARTICLE	IF	CITATIONS
1	Perspectives in liver redox imbalance: Toxicological and pharmacological aspects underlying iron overloading, nonalcoholic fatty liver disease, and thyroid hormone action. <i>BioFactors</i> , 2022, 48, 400-415.	5.4	44
2	Standpoints in mitochondrial dysfunction: Underlying mechanisms in search of therapeutic strategies. <i>Mitochondrion</i> , 2022, 63, 9-22.	3.4	9
3	N-3 PUFAs and their specialized pro-resolving lipid mediators on airway inflammatory response: beneficial effects in the prevention and treatment of respiratory diseases. <i>Food and Function</i> , 2022, 13, 4260-4272.	4.6	13
4	Influence of the nutritional status and oxidative stress in the desaturation and elongation of n-3 and n-6 polyunsaturated fatty acids: Impact on non-alcoholic fatty liver disease. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2022, 181, 102441.	2.2	33
5	Beneficial effects of natural compounds on experimental liver ischemia-reperfusion injury. <i>Food and Function</i> , 2021, 12, 3787-3798.	4.6	9
6	Effects of acute iron overload on Nrf2-related glutathione metabolism in rat brain. <i>BioMetals</i> , 2021, 34, 1017-1027.	4.1	9
7	N-3 Polyunsaturated Fatty Acids and Their Lipid Mediators as A Potential Immuneâ€“Nutritional Intervention: A Molecular and Clinical View in Hepatic Disease and Other Non-Communicable Illnesses. <i>Nutrients</i> , 2021, 13, 3384.	4.1	7
8	Targeting resolvins in cholestatic liver injury. <i>Hepatobiliary Surgery and Nutrition</i> , 2021, 10, 689-691.	1.5	0
9	Protective Effects of Eicosapentaenoic Acid Plus Hydroxytyrosol Supplementation Against White Adipose Tissue Abnormalities in Mice Fed a High-Fat Diet. <i>Molecules</i> , 2020, 25, 4433.	3.8	17
10	Impact of Mitophagy and Mitochondrial Unfolded Protein Response as New Adaptive Mechanisms Underlying Old Pathologies: Sarcopenia and Non-Alcoholic Fatty Liver Disease. <i>International Journal of Molecular Sciences</i> , 2020, 21, 7704.	4.1	37
11	The metabolic dysfunction of white adipose tissue induced in mice by a high-fat diet is abrogated by co-administration of docosahexaenoic acid and hydroxytyrosol. <i>Food and Function</i> , 2020, 11, 9086-9102.	4.6	25
12	Suppression of high-fat diet-induced obesity-associated liver mitochondrial dysfunction by docosahexaenoic acid and hydroxytyrosol co-administration. <i>Digestive and Liver Disease</i> , 2020, 52, 895-904.	0.9	78
13	Impact of the Co-Administration of N-3 Fatty Acids and Olive Oil Components in Preclinical Nonalcoholic Fatty Liver Disease Models: A Mechanistic View. <i>Nutrients</i> , 2020, 12, 499.	4.1	70
14	Iron-induced derangement in hepatic Δ^5 and Δ^6 desaturation capacity and fatty acid profile leading to steatosis: Impact on extrahepatic tissues and prevention by antioxidant-rich extra virgin olive oil. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2020, 153, 102058.	2.2	13
15	Docosahexaenoic acid and hydroxytyrosol co-administration fully prevents liver steatosis and related parameters in mice subjected to high-fat diet: A molecular approach. <i>BioFactors</i> , 2019, 45, 930-943.	5.4	42
16	High-fat diet induces mouse liver steatosis with a concomitant decline in energy metabolism: attenuation by eicosapentaenoic acid (EPA) or hydroxytyrosol (HT) supplementation and the additive effects upon EPA and HT co-administration. <i>Food and Function</i> , 2019, 10, 6170-6183.	4.6	62
17	Combined docosahexaenoic acid and thyroid hormone supplementation as a protocol supporting energy supply to precondition and afford protection against metabolic stress situations. <i>IUBMB Life</i> , 2019, 71, 1211-1220.	3.4	11
18	Docosahexaenoic acidâ€“thyroid hormone combined protocol as a novel approach to metabolic stress disorders: Relation to mitochondrial adaptation via liver PGC1 α and sirtuin1 activation. <i>BioFactors</i> , 2019, 45, 271-278.	5.4	10

#	ARTICLE	IF	CITATIONS
19	Hydroxytyrosol supplementation ameliorates the metabolic disturbances in white adipose tissue from mice fed a high-fat diet through recovery of transcription factors Nrf2, SREBP-1c, PPAR- α and NF- κ B. <i>Biomedicine and Pharmacotherapy</i> , 2019, 109, 2472-2481.	5.6	106
20	Combined administration of docosahexaenoic acid and thyroid hormone synergistically enhances rat liver levels of resolvins RvD1 and RvD2. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2019, 140, 42-46.	2.2	12
21	Reduction of high-fat diet-induced liver proinflammatory state by eicosapentaenoic acid plus hydroxytyrosol supplementation: involvement of resolvins RvE1/2 and RvD1/2. <i>Journal of Nutritional Biochemistry</i> , 2019, 63, 35-43.	4.2	80
22	Crosstalk mechanisms in hepatoprotection: Thyroid hormone-docosahexaenoic acid (DHA) and DHA-extra virgin olive oil combined protocols. <i>Pharmacological Research</i> , 2018, 132, 168-175.	7.1	26
23	Thyroid Hormone-Induced Expression of the Hepatic Scaffold Proteins Sestrin2, β -Klotho, and FRS2 α in Relation to FGF21-AMPK Signaling. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2018, 126, 182-186.	1.2	5
24	Iron-induced pro-oxidant and pro-lipogenic responses in relation to impaired synthesis and accretion of long-chain polyunsaturated fatty acids in rat hepatic and extrahepatic tissues. <i>Nutrition</i> , 2018, 45, 49-58.	2.4	36
25	Molecular mechanisms related to the hepatoprotective effects of antioxidant-rich extra virgin olive oil supplementation in rats subjected to short-term iron administration. <i>Free Radical Biology and Medicine</i> , 2018, 126, 313-321.	2.9	39
26	Attenuation of High-Fat Diet-Induced Rat Liver Oxidative Stress and Steatosis by Combined Hydroxytyrosol- (HT-) Eicosapentaenoic Acid Supplementation Mainly Relies on HT. <i>Oxidative Medicine and Cellular Longevity</i> , 2018, 2018, 1-13.	4.0	51
27	Anti-oxidative and anti-inflammatory effects of Rosa Mosqueta oil supplementation in rat liver ischemia-reperfusion. <i>Food and Function</i> , 2018, 9, 4847-4857.	4.6	15
28	Nutrients and Diet: A Relationship between Oxidative Stress, Aging, Obesity, and Related Noncommunicable Diseases. <i>Oxidative Medicine and Cellular Longevity</i> , 2018, 2018, 1-1.	4.0	8
29	Thyroid hormone suppresses ischemia-reperfusion-induced liver NLRP3 inflammasome activation: Role of AMP-activated protein kinase. <i>Immunology Letters</i> , 2017, 184, 92-97.	2.5	26
30	Hydroxytyrosol prevents reduction in liver activity of Δ ⁵ and Δ ⁶ desaturases, oxidative stress, and depletion in long chain polyunsaturated fatty acid content in different tissues of high-fat diet fed mice. <i>Lipids in Health and Disease</i> , 2017, 16, 64.	3.0	73
31	Molecular adaptations underlying the beneficial effects of hydroxytyrosol in the pathogenic alterations induced by a high-fat diet in mouse liver: PPAR- α and Nrf2 activation, and NF- κ B down-regulation. <i>Food and Function</i> , 2017, 8, 1526-1537.	4.6	109
32	A combined docosahexaenoic acid+thyroid hormone protocol upregulates rat liver β -Klotho expression and downstream components of FGF21 signaling as a potential novel approach to metabolic stress conditions. <i>Food and Function</i> , 2017, 8, 3980-3988.	4.6	7
33	Supplementation with Docosahexaenoic Acid and Extra Virgin Olive Oil Prevents Liver Steatosis Induced by a High-Fat Diet in Mice through PPAR- α and Nrf2 Upregulation with Concomitant SREBP-1c and NF- κ B Downregulation. <i>Molecular Nutrition and Food Research</i> , 2017, 61, 1700479.	3.3	106
34	Liver Protective Effects of Extra Virgin Olive Oil: Interaction between Its Chemical Composition and the Cell-signaling Pathways Involved in Protection. <i>Endocrine, Metabolic and Immune Disorders - Drug Targets</i> , 2017, 18, 75-84.	1.2	62
35	Hydroxytyrosol and Cytoprotection: A Projection for Clinical Interventions. <i>International Journal of Molecular Sciences</i> , 2017, 18, 930.	4.1	90
36	Upregulation of rat liver PPAR- α -FGF21 signaling by a docosahexaenoic acid and thyroid hormone combined protocol. <i>BioFactors</i> , 2016, 42, 638-646.	5.4	20

#	ARTICLE	IF	CITATIONS
37	Supplementation with antioxidant-rich extra virgin olive oil prevents hepatic oxidative stress and reduction of desaturation capacity in mice fed a high-fat diet: Effects on fatty acid composition in liver and extrahepatic tissues. <i>Nutrition</i> , 2016, 32, 1254-1267.	2.4	65
38	Long-chain polyunsaturated fatty acids regulation of PPARs, signaling: Relationship to tissue development and aging. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2016, 114, 28-34.	2.2	150
39	Hormone resuscitation therapy for brain-dead donors is insulin beneficial or detrimental?. <i>Clinical Transplantation</i> , 2016, 30, 754-759.	1.6	13
40	Thyroid hormone therapy and procurement of livers from brain-dead donors. <i>Endocrine Research</i> , 2016, 41, 270-273.	1.2	9
41	Sub-chronic iron overload triggers oxidative stress development in rat brain: implications for cell protection. <i>BioMetals</i> , 2016, 29, 119-130.	4.1	19
42	Anti-steatotic effects of an n-3 LCPUFA and extra virgin olive oil mixture in the liver of mice subjected to high-fat diet. <i>Food and Function</i> , 2016, 7, 140-150.	4.6	32
43	Thyroid hormone in the frontier of cell protection, survival and functional recovery. <i>Expert Reviews in Molecular Medicine</i> , 2015, 17, e10.	3.9	14
44	Relevant Aspects of Nutritional and Dietary Interventions in Non-Alcoholic Fatty Liver Disease. <i>International Journal of Molecular Sciences</i> , 2015, 16, 25168-25198.	4.1	99
45	Reduction in the desaturation capacity of the liver in mice subjected to high fat diet: Relation to LCPUFA depletion in liver and extrahepatic tissues. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 2015, 98, 7-14.	2.2	79
46	A combined iron and thyroid hormone protocol suppresses ischemia-reperfusion injury in rat livers. <i>RSC Advances</i> , 2015, 5, 26209-26217.	3.6	1
47	Causal role of oxidative stress in unfolded protein response development in the hyperthyroid state. <i>Free Radical Biology and Medicine</i> , 2015, 89, 401-408.	2.9	16
48	Nrf2 activation in the liver of rats subjected to a preconditioning sub-chronic iron protocol. <i>Food and Function</i> , 2014, 5, 243-250.	4.6	9
49	N-3 long-chain PUFA supplementation prevents high fat diet induced mouse liver steatosis and inflammation in relation to PPAR- α upregulation and NF- κ B DNA binding abrogation. <i>Molecular Nutrition and Food Research</i> , 2014, 58, 1333-1341.	3.3	72
50	Reversal of high-fat diet-induced hepatic steatosis by n-3 LCPUFA: role of PPAR- α and SREBP-1c. <i>Journal of Nutritional Biochemistry</i> , 2014, 25, 977-984.	4.2	47
51	T ₃ -induced liver AMP-activated protein kinase signaling: Redox dependency and upregulation of downstream targets. <i>World Journal of Gastroenterology</i> , 2014, 20, 17416.	3.3	16
52	Nrf2-regulated phase II detoxification enzymes and phase III transporters are induced by thyroid hormone in rat liver. <i>BioFactors</i> , 2013, 39, 514-521.	5.4	25
53	Reestablishment of Ischemia-Reperfusion Liver Injury by N-Acetylcysteine Administration prior to a Preconditioning Iron Protocol. <i>Scientific World Journal</i> , The, 2013, 2013, 1-9.	2.1	6
54	Thyroid hormone activates rat liver adenosine 5,-monophosphate-activated protein kinase: relation to CaMKKb, TAK1 and LKB1 expression and energy status. <i>Journal of Biological Regulators and Homeostatic Agents</i> , 2013, 27, 989-99.	0.7	4

#	ARTICLE	IF	CITATIONS
55	The oleic acid esterification of policosanol increases its bioavailability and hypocholesterolemic action in rats. <i>Grasas Y Aceites</i> , 2012, 63, 345-354.	0.9	9
56	Misregulation of PPAR Functioning and Its Pathogenic Consequences Associated with Nonalcoholic Fatty Liver Disease in Human Obesity. <i>PPAR Research</i> , 2012, 2012, 1-14.	2.4	63
57	Colloidal carbon stimulation of Kupffer cells triggers Nrf2 activation in the isolated perfused rat liver. <i>Journal of Molecular Histology</i> , 2012, 43, 343-349.	2.2	0
58	Vanillin suppresses Kupffer cell-related colloidal carbon-induced respiratory burst activity in isolated perfused rat liver: anti-inflammatory implications. <i>Food and Function</i> , 2012, 3, 1319.	4.6	13
59	Prevention of liver ischemia reperfusion injury by a combined thyroid hormone and fish oil protocol. <i>Journal of Nutritional Biochemistry</i> , 2012, 23, 1113-1120.	4.2	32
60	N-3 Long-Chain Polyunsaturated Fatty Acid Supplementation Significantly Reduces Liver Oxidative Stress in High Fat Induced Steatosis. <i>PLoS ONE</i> , 2012, 7, e46400.	2.5	88
61	Metabolic Basis for Thyroid Hormone Liver Preconditioning: Upregulation of AMP-Activated Protein Kinase Signaling. <i>Scientific World Journal</i> , The, 2012, 2012, 1-10.	2.1	7
62	Thyroid Hormone-Induced Cytosol-to-Nuclear Translocation of Rat Liver Nrf2 Is Dependent on Kupffer Cell Functioning. <i>Scientific World Journal</i> , The, 2012, 2012, 1-10.	2.1	4
63	Recent advances in liver preconditioning: Thyroid hormone, n-3 long-chain polyunsaturated fatty acids and iron. <i>World Journal of Hepatology</i> , 2012, 4, 119.	2.0	27
64	The importance of the long-chain polyunsaturated fatty acid n-6/n-3 ratio in development of non-alcoholic fatty liver associated with obesity. <i>Food and Function</i> , 2011, 2, 644.	4.6	146
65	Liver preconditioning induced by iron in a rat model of ischemia/reperfusion. <i>Life Sciences</i> , 2011, 89, 221-228.	4.3	22
66	Thyroid Hormone Administration Induces Rat Liver Nrf2 Activation: Suppression by N-Acetylcysteine Pretreatment. <i>Thyroid</i> , 2011, 21, 655-662.	4.5	41
67	Up-Regulation of PPAR- δ mRNA Expression in the Liver of Obese Patients: an Additional Reinforcing Lipogenic Mechanism to SREBP-1c Induction. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011, 96, 1424-1430.	3.6	288
68	N-3 PUFA Supplementation Triggers PPAR- δ Activation and PPAR- δ /NF- κ B Interaction: Anti-Inflammatory Implications in Liver Ischemia-Reperfusion Injury. <i>PLoS ONE</i> , 2011, 6, e28502.	2.5	167
69	Liver NF- κ B and AP-1 activation and PPAR- δ expression are negatively correlated in obese patients: Pro-inflammatory implications. <i>Clinical Nutrition</i> , 2010, 29, 687-688.	5.0	18
70	Homeostatic responses of thyroid hormone calorogenesis in the liver: Association with oxidative stress. <i>IUBMB Life</i> , 2010, 62, 460-466.	3.4	16
71	Decreased Liver Fatty Acid Δ^6 and Δ^5 Desaturase Activity in Obese Patients. <i>Obesity</i> , 2010, 18, 1460-1463.	3.0	104
72	Cytoprotective and suicidal signaling in oxidative stress. <i>Biological Research</i> , 2010, 43, .	3.4	8

#	ARTICLE	IF	CITATIONS
73	Protection against <i>in vivo</i> liver ischemia-reperfusion injury by n-3 long-chain polyunsaturated fatty acids in the rat. <i>Free Radical Research</i> , 2010, 44, 854-863.	3.3	44
74	Kupffer-cell activity is essential for thyroid hormone rat liver preconditioning. <i>Molecular and Cellular Endocrinology</i> , 2010, 323, 292-297.	3.2	13
75	Biosynthesis and bioavailability of long-chain polyunsaturated fatty acids in non-alcoholic fatty liver disease. <i>Progress in Lipid Research</i> , 2010, 49, 407-419.	11.6	42
76	Delayed Ischemic Preconditioning Protects Against Liver Ischemia Reperfusion Injury In Vivo. <i>Transplantation Proceedings</i> , 2010, 42, 1569-1575.	0.6	14
77	Cytoprotective and suicidal signaling in oxidative stress. <i>Biological Research</i> , 2010, 43, 363-9.	3.4	9
78	Oxidative stress signaling underlying liver disease and hepatoprotective mechanisms. <i>World Journal of Hepatology</i> , 2009, 1, 72.	2.0	120
79	Liver NF- κ B and AP-1 DNA Binding in Obese Patients. <i>Obesity</i> , 2009, 17, 973-979.	3.0	85
80	Trace quantification of 1-octacosanol and 1-triacontanol and their main metabolites in plasma by liquid-liquid extraction coupled with gas chromatography-mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2009, 877, 4154-4158.	2.3	15
81	Enhancement in liver SREBP-1c/PPAR- α ratio and steatosis in obese patients: Correlations with insulin resistance and n-3 long-chain polyunsaturated fatty acid depletion. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2009, 1792, 1080-1086.	3.8	183
82	Upregulation of liver inducible nitric oxide synthase following thyroid hormone preconditioning: suppression by N-acetylcysteine. <i>Biological Research</i> , 2009, 42, .	3.4	18
83	Causal role of oxidative stress in liver preconditioning by thyroid hormone in rats. <i>Free Radical Biology and Medicine</i> , 2008, 44, 1724-1731.	2.9	35
84	Study of Cytochrome P450 2E1 and its allele Variants in Liver Injury of Nondiabetic, Nonalcoholic Steatohepatitis Obese Women. <i>Biological Research</i> , 2008, 41, .	3.4	33
85	Effects of weight loss on liver and erythrocyte polyunsaturated fatty acid pattern and oxidative stress status in obese patients with non-alcoholic fatty liver disease. <i>Biological Research</i> , 2008, 41, .	3.4	43
86	Involvement of Kupffer cell-dependent signaling in T ₃ -induced hepatocyte proliferation <i>in vivo</i> . <i>Biological Chemistry</i> , 2007, 388, 831-837.	2.5	22
87	Hepatoprotective role of nitric oxide in an experimental model of chronic iron overload. <i>Nitric Oxide - Biology and Chemistry</i> , 2007, 16, 143-149.	2.7	5
88	Thyroid hormone preconditioning: Protection against ischemia-reperfusion liver injury in the rat. <i>Hepatology</i> , 2007, 45, 170-177.	7.3	67
89	Polyunsaturated Fatty Acid Pattern in Liver and Erythrocyte Phospholipids from Obese Patients. <i>Obesity</i> , 2007, 15, 24-31.	3.0	109
90	Thyroid hormone calorigenesis and mitochondrial redox signaling: upregulation of gene expression. <i>Frontiers in Bioscience - Landmark</i> , 2007, 12, 1220.	3.0	25

#	ARTICLE	IF	CITATIONS
91	Thyroid hormone-induced oxidative stress in rodents and humans: A comparative view and relation to redox regulation of gene expression. <i>Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology</i> , 2006, 142, 231-239.	2.6	47
92	Increased production of IL-1 β and TNF- α in lipopolysaccharide-stimulated blood from obese patients with non-alcoholic fatty liver disease. <i>Cytokine</i> , 2006, 33, 252-257.	3.2	72
93	Insulin resistance and oxidative stress interdependency in non-alcoholic fatty liver disease. <i>Trends in Molecular Medicine</i> , 2006, 12, 555-558.	6.7	141
94	Relationship between in vivo chlorzoxazone hydroxylation, hepatic cytochrome P450 2E1 content and liver injury in obese non-alcoholic fatty liver disease patients. <i>Hepatology Research</i> , 2006, 34, 57-63.	3.4	67
95	The acute-phase response of the liver in relation to thyroid hormone-induced redox signaling. <i>Free Radical Biology and Medicine</i> , 2006, 40, 1628-1635.	2.9	33
96	Redox up-regulated expression of rat liver manganese superoxide dismutase and Bcl-2 by thyroid hormone is associated with inhibitor of $\text{I}\kappa\text{B}$ phosphorylation and nuclear factor- $\text{I}\kappa\text{B}$ activation. <i>Journal of Endocrinology</i> , 2005, 186, 539-547.	2.6	47
97	Redox regulation of thyroid hormone-induced Kupffer cell-dependent $\text{I}\kappa\text{B}$ phosphorylation in relation to inducible nitric oxide synthase expression. <i>Free Radical Research</i> , 2005, 39, 411-418.	3.3	26
98	Chronic iron overload enhances inducible nitric oxide synthase expression in rat liver. <i>Nitric Oxide - Biology and Chemistry</i> , 2005, 13, 54-61.	2.7	50
99	Effects of Acute I^3 -Hexachlorocyclohexane Intoxication in Relation to the Redox Regulation of Nuclear Factor- $\text{I}\kappa\text{B}$, Cytokine Gene Expression, and Liver Injury in the Rat. <i>Antioxidants and Redox Signaling</i> , 2004, 6, 471-480.	5.4	20
100	Oxidative stress and depletion of hepatic long-chain polyunsaturated fatty acids may contribute to nonalcoholic fatty liver disease. <i>Free Radical Biology and Medicine</i> , 2004, 37, 1499-1507.	2.9	215
101	Tri-iodothyronine differentially induces Kupffer cell ED1/ED2 subpopulations. <i>Molecular Aspects of Medicine</i> , 2004, 25, 183-190.	6.4	18
102	Oxidative stress-related parameters in the liver of non-alcoholic fatty liver disease patients. <i>Clinical Science</i> , 2004, 106, 261-268.	4.3	426
103	Increase in long-chain polyunsaturated fatty acid n \sim 6/n \sim 3 ratio in relation to hepatic steatosis in patients with non-alcoholic fatty liver disease. <i>Clinical Science</i> , 2004, 106, 635-643.	4.3	576
104	Oxidative stress-mediated hepatotoxicity of iron and copper: role of Kupffer cells. <i>BioMetals</i> , 2003, 16, 103-111.	4.1	117
105	Thyroid hormone-induced oxidative stress triggers nuclear factor- $\text{I}\kappa\text{B}$ activation and cytokine gene expression in rat liver. <i>Free Radical Biology and Medicine</i> , 2003, 35, 257-265.	2.9	57
106	Kupffer cell stimulation in the isolated perfused rat liver triggers nuclear factor- $\text{I}\kappa\text{B}$ DNA binding activity. <i>Redox Report</i> , 2003, 8, 341-346.	4.5	7
107	Effects of g-hexachlorocyclohexane and L-3,3',5- triiodothyronine on rat liver cytochrome P4502E1-dependent activity and content in relation to microsomal superoxide radical generation. <i>Biological Research</i> , 2003, 36, 359-65.	3.4	17
108	Enhancement of Lindane-induced Liver Oxidative Stress and Hepatotoxicity by Thyroid Hormone is Reduced by Gadolinium Chloride. <i>Free Radical Research</i> , 2002, 36, 1033-1039.	3.3	44

#	ARTICLE	IF	CITATIONS
109	Increases in Tumor Necrosis Factor- $\hat{I}\alpha$ in Response to Thyroid Hormone-induced Liver Oxidative Stress in the Rat. <i>Free Radical Research</i> , 2002, 36, 719-725.	3.3	26
110	Iron-induced changes in nitric oxide and superoxide radical generation in rat liver after lindane or thyroid hormone treatment. <i>Toxicology Letters</i> , 2001, 119, 87-93.	0.8	30
111	Effects of Iron Overload and Lindane Intoxication in Relation to Oxidative Stress, Kupffer Cell Function, and Liver Injury in the Rat. <i>Toxicology and Applied Pharmacology</i> , 2001, 170, 23-28.	2.8	26
112	Energy metabolism, thyroid calorigenesis, and oxidative stress: functional and cytotoxic consequences. <i>Redox Report</i> , 2000, 5, 265-275.	4.5	73
113	Derangement of Kupffer cell functioning and hepatotoxicity in hyperthyroid rats subjected to acute iron overload. <i>Redox Report</i> , 1999, 4, 243-250.	4.5	12
114	Influence of copper-(II) on colloidal carbon-induced kupffer cell-dependent oxygen uptake in rat liver: Relation to hepatotoxicity. <i>Free Radical Research</i> , 1999, 30, 489-498.	3.3	10
115	Protein oxidation in thyroid hormone-induced liver oxidative stress: relation to lipid peroxidation. <i>Toxicology Letters</i> , 1999, 106, 209-214.	0.8	54
116	Liver microsomal parameters related to oxidative stress and antioxidant systems in hyperthyroid rats subjected to acute lindane treatment. <i>Free Radical Research</i> , 1998, 29, 35-42.	3.3	27
117	Dose-dependent effects of acute lindane treatment on Kupffer cell function assessed in the isolated perfused rat liver. <i>Xenobiotica</i> , 1997, 27, 747-757.	1.1	13
118	Acetaminophen-induced liver oxidative stress and hepatotoxicity: influence of Kupffer cell activity assessed in the isolated perfused rat liver. <i>Redox Report</i> , 1997, 3, 213-218.	4.5	7
119	Kupffer Cell Function in Thyroid Hormone-Induced Liver Oxidative Stress in the Rat. <i>Free Radical Research</i> , 1997, 26, 267-279.	3.3	59
120	Influence of Hyperthyroidism on the Activity of Liver Nitric Oxide Synthase in the Rat. <i>Nitric Oxide - Biology and Chemistry</i> , 1997, 1, 463-468.	2.7	47
121	Potentialiation of Ischemia-Reperfusion Liver Injury by Hyperthyroidism in the Rat. <i>Free Radical Biology and Medicine</i> , 1997, 23, 19-25.	2.9	27
122	Regression of morphological alterations and oxidative stress-related parameters after acute lindane-induced hepatotoxicity in rats. <i>Toxicology</i> , 1997, 117, 199-205.	4.2	30
123	Effect of aliphatic aldehydes on the lipid peroxidation and chemiluminescence of biological systems under oxidative stress. <i>Luminescence</i> , 1997, 12, 141-148.	0.0	1
124	Hepatic glutathione biosynthetic capacity in hyperthyroid rat. <i>Toxicology Letters</i> , 1996, 89, 85-89.	0.8	25
125	Effect of hyperthyroidism on the biliary release of thiobarbituric acid reactants in the rat. <i>Toxicology Letters</i> , 1996, 84, 149-153.	0.8	22
126	On the mechanism of thyroid hormone-induced respiratory burst activity in rat polymorphonuclear leukocytes. <i>Free Radical Biology and Medicine</i> , 1995, 19, 359-363.	2.9	38

#	ARTICLE	IF	CITATIONS
127	Influence of hyperthyroidism on lindane-induced hepatotoxicity in the rat. <i>Biochemical Pharmacology</i> , 1995, 50, 1557-1565.	4.4	38
128	Zymosan-induced luminol-amplified chemiluminescence of whole blood phagocytes in experimental and human hyperthyroidism. <i>Free Radical Biology and Medicine</i> , 1993, 14, 669-675.	2.9	33
129	3,3',5-Triiodothyronine-induced hepatic respiration: Effects of desferrioxamine and allopurinol in the isolated perfused rat liver. <i>Toxicology Letters</i> , 1993, 69, 205-210.	0.8	18
130	Influence of Hyperthyroidism on Superoxide Radical and Hydrogen Peroxide Production by Rat Liver Submitochondrial Particles. <i>Free Radical Research Communications</i> , 1993, 18, 329-335.	1.8	85
131	Is spontaneous urinary visible chemiluminescence a reflection of in vivo oxidative stress?. <i>Free Radical Biology and Medicine</i> , 1992, 12, 317-322.	2.9	29
132	Effects of Hyperthyroidism on Rat Liver Glutathione Metabolism: Related Enzymes' Activities, Efflux, and Turnover*. <i>Endocrinology</i> , 1991, 129, 85-91.	2.8	62
133	Free Radical Scavenging Activity of Carnosine. <i>Free Radical Research Communications</i> , 1991, 14, 263-270.	1.8	27
134	Lindane-induced liver oxidative stress. <i>Free Radical Biology and Medicine</i> , 1990, 9, 169-179.	2.9	91
135	Lindane-induced liver oxidative stress: Respiratory alterations and the effect of desferrioxamine in the isolated perfused rat liver. <i>Cell Biochemistry and Function</i> , 1989, 7, 179-183.	2.9	11
136	Increased Lipid Peroxidation in Hyperthyroid Patients: Suppression by Propylthiouracil Treatment. <i>Free Radical Research Communications</i> , 1988, 5, 1-10.	1.8	59
137	Chemiluminescent and Respiratory Responses Related to Thyroid Hormone-Induced Liver Oxidative Stress. <i>Free Radical Research Communications</i> , 1988, 5, 77-84.	1.8	72
138	Lindane-Induced Oxidative Stress. II. Time Course of Changes in Hepatic Glutathione Status. <i>Xenobiotica</i> , 1988, 18, 1305-1310.	1.1	32
139	Lindane-Induced Oxidative Stress. I. Time Course of Changes in Hepatic Microsomal Parameters, Antioxidant Enzymes, Lipid Peroxidative Indices and Morphological Characteristics. <i>Xenobiotica</i> , 1988, 18, 1297-1304.	1.1	66
140	Dose-dependent study of the effects of acute lindane administration on rat liver superoxide anion production, antioxidant enzyme activities and lipid peroxidation. <i>Toxicology</i> , 1986, 41, 193-204.	4.2	75
141	Superoxide Radical Generation, NADPH Oxidase Activity, and Cytochrome P-450 Content of Rat Liver Microsomal Fractions in an Experimental Hyperthyroid State: Relation to Lipid Peroxidation*. <i>Endocrinology</i> , 1985, 117, 496-501.	2.8	146