

# Masato Uehara

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3259342/publications.pdf>

Version: 2024-02-01

13  
papers

88  
citations

1478505

6  
h-index

1588992

8  
g-index

14  
all docs

14  
docs citations

14  
times ranked

96  
citing authors

#	ARTICLE	IF	CITATIONS
1	Life-history traits of a widely distributed coral reef fish, <i>Mulloidichthys flavolineatus</i> (Mullidae), Okinawa Island, southern Japan. <i>Environmental Biology of Fishes</i> , 2021, 104, 1559-1574.	1.0	3
2	Early development and occurrence patterns of the Pagrinae seabream <i>Argyrops bleekeri</i> (Perciformes: Tj ETQq0 0 0 rgBT /Overlock 10 T 2020, 96, 631-641.	1.6	3
3	Comparative age-specific demography of four commercially important deep-water snappers: implication for fishery management of a long-lived lutjanid. <i>Journal of Fish Biology</i> , 2020, 97, 121-136.	1.6	10
4	Effectiveness of deepwater marine protected areas: Implication for Okinawan demersal fisheries management. <i>Fisheries Research</i> , 2019, 215, 123-130.	1.7	7
5	Reproductive traits of deep-sea snappers (Lutjanidae): Implication for Okinawan bottomfish fisheries management. <i>Regional Studies in Marine Science</i> , 2018, 17, 112-126.	0.7	9
6	Life history traits of the pagrinae sparid <i>Argyrops bleekeri</i> inhabiting bays in Okinawa Island, Ryukyu Archipelago, southwestern Japan. <i>Regional Studies in Marine Science</i> , 2017, 10, 97-106.	0.7	4
7	Age-based demography and reproductive biology of three <i>Epinephelus</i> groupers, <i>E. polyphekadion</i> , <i>E. tauvina</i> , and <i>E. howlandi</i> (Serranidae), inhabiting coral reefs in Okinawa. <i>Environmental Biology of Fishes</i> , 2017, 100, 1451-1467.	1.0	9
8	Life history variables, annual change in sex ratios with age, and total mortality observed on commercial catch on Pacific steephead parrotfish, <i>Chlorurus microrhinos</i> in waters off the Okinawa Island, southwestern Japan. <i>Regional Studies in Marine Science</i> , 2016, 8, 65-76.	0.7	9
9	Changes of group construction accompanying with growth and maturity in blue-barred parrotfish ( <i>Scarus ghobban</i> ), and influences of the fishing targeting the immature group to the stock. <i>Regional Studies in Marine Science</i> , 2016, 7, 32-42.	0.7	7
10	Reproductive traits of the western Pacific gizzard shad <i>Nematalosa come</i> : implications for conservation and management in a population at its distributional margin. <i>Environmental Biology of Fishes</i> , 2015, 98, 1-10.	1.0	6
11	Reproductive biology of Japanese gizzard shad ( <i>Nematalosa japonica</i> ) in coastal waters around Okinawa Island, Ryukyu Archipelago, southwestern Japan. <i>Ichthyological Research</i> , 2012, 59, 314-322.	0.8	6
12	Biological traits of naturally induced hybrid individuals of two gizzard shads, <i>Nematalosa come</i> and <i>N. japonica</i> , in coastal waters around Okinawa Island, Ryukyu Archipelago, southwestern Japan. <i>Ichthyological Research</i> , 2011, 58, 344-349.	0.8	4
13	Age and growth of two gizzard shads, <i>Nematalosa come</i> and <i>N. japonica</i> , in coastal waters around Okinawa Island, southwestern Japan. <i>Fisheries Science</i> , 2009, 75, 295-301.	1.6	11