

Paul R Albert

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3261921/publications.pdf>

Version: 2024-02-01

135
papers

8,797
citations

57631

44
h-index

46693

89
g-index

146
all docs

146
docs citations

146
times ranked

9250
citing authors

#	ARTICLE	IF	CITATIONS
1	Depression, dementia and immune dysregulation. <i>Brain</i> , 2021, 144, 746-760.	3.7	81
2	Rewiring of the Serotonin System in Major Depression. <i>Frontiers in Psychiatry</i> , 2021, 12, 802581.	1.3	28
3	Influence of functional gene polymorphisms on human behaviour: the case of <i>CCR5</i> . <i>Journal of Psychiatry and Neuroscience</i> , 2021, 46, E659-E662.	1.4	0
4	Fluoxetine-induced recovery of serotonin and norepinephrine projections in a mouse model of post-stroke depression. <i>Translational Psychiatry</i> , 2020, 10, 334.	2.4	21
5	Orphans to the rescue: orphan G-protein coupled receptors as new antidepressant targets. <i>Journal of Psychiatry and Neuroscience</i> , 2020, 45, 301-303.	1.4	1
6	Orphans to the rescue: orphan G-protein coupled receptors as new antidepressant targets. <i>Journal of Psychiatry and Neuroscience</i> , 2020, 45, 301-303.	1.4	3
7	Genetic, epigenetic and posttranscriptional mechanisms for treatment of major depression: the 5-HT1A receptor gene as a paradigm. <i>Journal of Psychiatry and Neuroscience</i> , 2019, 44, 164-176.	1.4	41
8	Targeting Homer1a for Rapid Antidepressant Effects. <i>Neuron</i> , 2019, 104, 182-183.	3.8	4
9	The Transcription Factor Deaf1 Modulates Engrailed-1 Expression to Regulate Skin Appendage Fate. <i>Journal of Investigative Dermatology</i> , 2019, 139, 2378-2381.e4.	0.3	9
10	Biased signaling of G protein coupled receptors (GPCRs): Molecular determinants of GPCR/transducer selectivity and therapeutic potential. , 2019, 200, 148-178.		100
11	Overcoming Resistance to Selective Serotonin Reuptake Inhibitors: Targeting Serotonin, Serotonin-1A Receptors and Adult Neuroplasticity. <i>Frontiers in Neuroscience</i> , 2019, 13, 404.	1.4	29
12	The 5-HT1A receptor: Signaling to behavior. <i>Biochimie</i> , 2019, 161, 34-45.	1.3	114
13	Loss of Adult 5-HT1A Autoreceptors Results in a Paradoxical Anxiogenic Response to Antidepressant Treatment. <i>Journal of Neuroscience</i> , 2019, 39, 1334-1346.	1.7	19
14	Adult neuroplasticity: A new "cure" for major depression?. <i>Journal of Psychiatry and Neuroscience</i> , 2019, 44, 147-150.	1.4	22
15	Loss of MeCP2 in adult 5-HT neurons induces 5-HT1A autoreceptors, with opposite sex-dependent anxiety and depression phenotypes. <i>Scientific Reports</i> , 2018, 8, 5788.	1.6	28
16	A Novel Alternative Splicing Mechanism That Enhances Human 5-HT1A Receptor RNA Stability Is Altered in Major Depression. <i>Journal of Neuroscience</i> , 2018, 38, 8200-8210.	1.7	30
17	Is poststroke depression the same as major depression?. <i>Journal of Psychiatry and Neuroscience</i> , 2018, 43, 76-77.	1.4	9
18	Chronic Fluoxetine Induces Activity Changes in Recovery From Poststroke Anxiety, Depression, and Cognitive Impairment. <i>Neurotherapeutics</i> , 2018, 15, 200-215.	2.1	21

#	ARTICLE	IF	CITATIONS
19	Recruitment by the Repressor Freud-1 of Histone Deacetylase-Brg1 Chromatin Remodeling Complexes to Strengthen HTR1A Gene Repression. <i>Molecular Neurobiology</i> , 2017, 54, 8263-8277.	1.9	4
20	Length of axons expressing the serotonin transporter in orbitofrontal cortex is lower with age in depression. <i>Neuroscience</i> , 2017, 359, 30-39.	1.1	21
21	Abrogated Freud-1/Cc2d1a Repression of 5-HT1A Autoreceptors Induces Fluoxetine-Resistant Anxiety/Depression-Like Behavior. <i>Journal of Neuroscience</i> , 2017, 37, 11967-11978.	1.7	35
22	The adaptive brain in mental health: overcoming inherited risk factors. <i>Journal of Psychiatry and Neuroscience</i> , 2017, 42, 3-5.	1.4	1
23	The functional serotonin 1a receptor promoter polymorphism, rs6295, is associated with psychiatric illness and differences in transcription. <i>Translational Psychiatry</i> , 2016, 6, e746-e746.	2.4	49
24	Persistent post-stroke depression in mice following unilateral medial prefrontal cortical stroke. <i>Translational Psychiatry</i> , 2016, 6, e863-e863.	2.4	69
25	Sex-dependent adaptive changes in serotonin-1A autoreceptor function and anxiety in Deaf1-deficient mice. <i>Molecular Brain</i> , 2016, 9, 77.	1.3	22
26	Concentration-Dependent Dual Mode of Zn Action at Serotonin 5-HT1A Receptors: In Vitro and In Vivo Studies. <i>Molecular Neurobiology</i> , 2016, 53, 6869-6881.	1.9	30
27	COMT polymorphism modulates the resting-state EEG alpha oscillatory response to acute nicotine in male non-smokers. <i>Genes, Brain and Behavior</i> , 2015, 14, 466-476.	1.1	7
28	Why is depression more prevalent in women?. <i>Journal of Psychiatry and Neuroscience</i> , 2015, 40, 219-221.	1.4	1,007
29	Evidence Revealing Deregulation of The KLF11-Mao A Pathway in Association with Chronic Stress and Depressive Disorders. <i>Neuropsychopharmacology</i> , 2015, 40, 1373-1382.	2.8	35
30	Chronic mild stress and antidepressant treatment alter 5-HT1A receptor expression by modifying DNA methylation of a conserved Sp4 site. <i>Neurobiology of Disease</i> , 2015, 82, 332-341.	2.1	53
31	Requirement of a Blocking Step in Affinity Purification of Polyclonal Antibodies. <i>International Journal of Molecular and Cellular Medicine</i> , 2015, 4, 196-8.	1.1	0
32	Serotonin-prefrontal cortical circuitry in anxiety and depression phenotypes: pivotal role of pre- and post-synaptic 5-HT1A receptor expression. <i>Frontiers in Behavioral Neuroscience</i> , 2014, 8, 199.	1.0	222
33	Light up your life: Optogenetics for depression?. <i>Journal of Psychiatry and Neuroscience</i> , 2014, 39, 3-5.	1.4	19
34	Stress-induced alterations in 5-HT1A receptor transcriptional modulators NUDR and Freud-1. <i>International Journal of Neuropsychopharmacology</i> , 2014, 17, 1763-1775.	1.0	24
35	Editorial. <i>International Journal of Neuropsychopharmacology</i> , 2014, 17, 1727-1728.	1.0	0
36	The Expression of KLF11 (TIEG2), a Monoamine Oxidase B Transcriptional Activator in the Prefrontal Cortex of Human Alcohol Dependence. <i>Alcoholism: Clinical and Experimental Research</i> , 2014, 38, 144-151.	1.4	15

#	ARTICLE	IF	CITATIONS
37	Transcriptional Dys-regulation in Anxiety and Major Depression: 5-HT1A Gene Promoter Architecture as a Therapeutic Opportunity. <i>Current Pharmaceutical Design</i> , 2014, 20, 3738-3750.	0.9	38
38	Role of protein kinase C in agonist-induced desensitization of 5-HT1A receptor coupling to calcium channels in F11 cells. <i>European Journal of Pharmacology</i> , 2013, 706, 84-91.	1.7	2
39	Effects of COMT genotype on sensory gating and its modulation by nicotine: Differences in low and high P50 suppressors. <i>Neuroscience</i> , 2013, 241, 147-156.	1.1	23
40	The neurobiology of depression—revisiting the serotonin hypothesis. II. Genetic, epigenetic and clinical studies. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2013, 368, 20120535.	1.8	79
41	DEAF1 Is a Pellino1-interacting Protein Required for Interferon Production by Sendai Virus and Double-stranded RNA*. <i>Journal of Biological Chemistry</i> , 2013, 288, 24569-24580.	1.6	28
42	Drugs for kids: Good or bad?. <i>Journal of Psychiatry and Neuroscience</i> , 2012, 37, 293-295.	1.4	1
43	Mechanistic Role for a Novel Glucocorticoid-KLF11 (TIEG2) Protein Pathway in Stress-induced Monoamine Oxidase A Expression. <i>Journal of Biological Chemistry</i> , 2012, 287, 24195-24206.	1.6	80
44	The neurobiology of depression—revisiting the serotonin hypothesis. I. Cellular and molecular mechanisms. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2012, 367, 2378-2381.	1.8	155
45	Transcriptional regulation of the 5-HT _{1A} receptor: implications for mental illness. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2012, 367, 2402-2415.	1.8	102
46	Increased Serotonin-1A (5-HT1A) Autoreceptor Expression and Reduced Raphe Serotonin Levels in Deformed Epidermal Autoregulatory Factor-1 (Deaf-1) Gene Knock-out Mice. <i>Journal of Biological Chemistry</i> , 2012, 287, 6615-6627.	1.6	67
47	17 β -Estradiol-Induced Regulation of the Novel 5-HT1A-Related Transcription Factors NUDR and Freud-1 in SH SY5Y Cells. <i>Cellular and Molecular Neurobiology</i> , 2012, 32, 517-521.	1.7	13
48	Ser ³ Thr residues at ± 3 loop of G \pm s are important in morphine-induced adenylyl cyclase sensitization but not mitogen-activated protein kinase phosphorylation. <i>FEBS Journal</i> , 2012, 279, 650-660.	2.2	7
49	Brain derived neurotrophic factor, cardiopulmonary fitness and cognition in patients with coronary artery disease. <i>Brain, Behavior, and Immunity</i> , 2011, 25, 1264-1271.	2.0	39
50	The moderating role of the dopamine transporter 1 gene on P50 sensory gating and its modulation by nicotine. <i>Neuroscience</i> , 2011, 180, 148-156.	1.1	25
51	Freud-2/CC2D1B mediates dual repression of the serotonin-1A receptor gene. <i>European Journal of Neuroscience</i> , 2011, 33, 214-223.	1.2	13
52	Region-specific regulation of 5-HT1A receptor expression by Pet ¹ -dependent mechanisms <i>in vivo</i> . <i>Journal of Neurochemistry</i> , 2011, 116, 1066-1076.	2.1	26
53	A functional alternative splicing mutation in human tryptophan hydroxylase-2. <i>Molecular Psychiatry</i> , 2011, 16, 1169-1176.	4.1	21
54	Transcriptional dysregulation of 5-HT1A autoreceptors in mental illness. <i>Molecular Brain</i> , 2011, 4, 21.	1.3	112

#	ARTICLE	IF	CITATIONS
55	Neurotensin Triggers Dopamine D2 Receptor Desensitization through a Protein Kinase C and β^2 -Arrestin1-dependent Mechanism. <i>Journal of Biological Chemistry</i> , 2011, 286, 9174-9184.	1.6	50
56	The Reduction of R1, a Novel Repressor Protein for Monoamine Oxidase A, in Major Depressive Disorder. <i>Neuropsychopharmacology</i> , 2011, 36, 2139-2148.	2.8	73
57	What is a functional genetic polymorphism? Defining classes of functionality. <i>Journal of Psychiatry and Neuroscience</i> , 2011, 36, 363-365.	1.4	36
58	TNFAIP8: A new effector for Galpha(i) coupling to reduce cell death and induce cell transformation. <i>Journal of Cellular Physiology</i> , 2010, 225, 865-874.	2.0	46
59	Modifying 5-HT1A receptor gene expression as a new target for antidepressant therapy. <i>Frontiers in Neuroscience</i> , 2010, 4, 35.	1.4	66
60	Decreased expression of Freud-1/CC2D1A, a transcriptional repressor of the 5-HT1A receptor, in the prefrontal cortex of subjects with major depression. <i>International Journal of Neuropsychopharmacology</i> , 2010, 13, 1089-1101.	1.0	32
61	Epigenetics in mental illness: Hope or hype?. <i>Journal of Psychiatry and Neuroscience</i> , 2010, 35, 366-368.	1.4	13
62	Effects of nicotine on the amplitude and gating of the auditory P50 and its influence by dopamine D2 receptor gene polymorphism. <i>Neuroscience</i> , 2010, 166, 145-156.	1.1	35
63	Gender-specific decrease in NUDR and 5-HT1A receptor proteins in the prefrontal cortex of subjects with major depressive disorder. <i>International Journal of Neuropsychopharmacology</i> , 2009, 12, 155.	1.0	71
64	F.86. Deaf1 Isoforms Control Changes in Peripheral Tissue Antigen Gene Expression in the Non-obese Diabetic Mouse Pancreatic Lymph Node during Type I Diabetes Pathogenesis. <i>Clinical Immunology</i> , 2009, 131, S117.	1.4	0
65	Deaf1 isoforms control the expression of genes encoding peripheral tissue antigens in the pancreatic lymph nodes during type 1 diabetes. <i>Nature Immunology</i> , 2009, 10, 1026-1033.	7.0	134
66	Differential regulation of the serotonin 1 A transcriptional modulators five prime repressor element under dual repression-1 and nuclear-deformed epidermal autoregulatory factor by chronic stress. <i>Neuroscience</i> , 2009, 163, 1119-1127.	1.1	26
67	Human Freud-2/CC2D1B: A Novel Repressor of Postsynaptic Serotonin-1A Receptor Expression. <i>Biological Psychiatry</i> , 2009, 66, 214-222.	0.7	36
68	A Nurr1 point mutant, implicated in Parkinson's disease, uncouples ERK1/2-dependent regulation of tyrosine hydroxylase transcription. <i>Neurobiology of Disease</i> , 2008, 29, 117-122.	2.1	43
69	HES1 regulates 5-HT1A receptor gene transcription at a functional polymorphism: Essential role in developmental expression. <i>Molecular and Cellular Neurosciences</i> , 2008, 38, 349-358.	1.0	29
70	Transcriptional regulation at a HTR1A polymorphism associated with mental illness. <i>Neuropharmacology</i> , 2008, 55, 977-985.	2.0	158
71	Roles of G protein and β^2 -arrestin in dopamine D 2 receptor-mediated ERK activation. <i>Biochemical and Biophysical Research Communications</i> , 2008, 377, 705-709.	1.0	23
72	GAP1(IP4BP)/RASA3 Mediates $G_{\beta\gamma}$ -induced Inhibition of Mitogen-activated Protein Kinase. <i>Journal of Biological Chemistry</i> , 2008, 283, 35908-35917.	1.6	12

#	ARTICLE	IF	CITATIONS
73	Differential Repression by Freud-1/CC2D1A at a Polymorphic Site in the Dopamine-D2 Receptor Gene*. Journal of Biological Chemistry, 2007, 282, 20897-20905.	1.6	33
74	Role of Cdk5-Mediated Phosphorylation of Prx2 in MPTP Toxicity and Parkinson's Disease. Neuron, 2007, 55, 37-52.	3.8	225
75	The Freud-1/CC2D1A family: Transcriptional regulators implicated in mental retardation. Journal of Neuroscience Research, 2007, 85, 2833-2838.	1.3	24
76	Computerized measurement of facial expression of emotions in schizophrenia. Journal of Neuroscience Methods, 2007, 163, 350-361.	1.3	39
77	Characterization of rat rostral raphe primary cultures: Multiplex quantification of serotonergic markers. Journal of Neuroscience Methods, 2007, 164, 59-67.	1.3	12
78	The mental retardation gene CC2D1A/Freudâ€¹1 encodes a long isoform that binds conserved DNA elements to repress gene transcription. European Journal of Neuroscience, 2007, 26, 965-974.	1.2	22
79	Differential signaling of dopamine-D2S and -D2L receptors to inhibit ERK1/2 phosphorylation. Journal of Neurochemistry, 2007, 102, 1796-1804.	2.1	18
80	Cell-type specific induction of tryptophan hydroxylase-2 transcription by calcium mobilization. Journal of Neurochemistry, 2007, 103, 2047-2057.	2.1	19
81	Differential desensitization of dopamine D2 receptor isoforms by protein kinase C: The importance of receptor phosphorylation and pseudosubstrate sites. European Journal of Pharmacology, 2007, 577, 44-53.	1.7	25
82	Identification of Novel Transcriptional Regulators in the Nervous System. Frontiers in Neuroscience, 2007, , 81-103.	0.0	0
83	RGS17/RGSZ2 and the RZ/A family of regulators of G-protein signaling. Seminars in Cell and Developmental Biology, 2006, 17, 390-399.	2.3	38
84	Molecular Determinants in the Second Intracellular Loop of the 5-Hydroxytryptamine-1A Receptor for G-Protein Coupling. Molecular Pharmacology, 2006, 69, 1518-1526.	1.0	28
85	Cell-Specific Repressor or Enhancer Activities of Deaf-1 at a Serotonin 1A Receptor Gene Polymorphism. Journal of Neuroscience, 2006, 26, 1864-1871.	1.7	124
86	Specific residues of the 5â€¹HT1A receptor second and third intracellular domain Câ€¹terminal determine GÎ²3 or GÎ±i coupling specificity, respectively. FASEB Journal, 2006, 20, A918.	0.2	0
87	Coupling of 5-HT1A autoreceptors to inhibition of mitogen-activated protein kinase activation via GÎ²3 subunit signaling. European Journal of Neuroscience, 2005, 21, 721-732.	1.2	47
88	Differential Roles of Nuclear and Cytoplasmic Cyclin-Dependent Kinase 5 in Apoptotic and Excitotoxic Neuronal Death. Journal of Neuroscience, 2005, 25, 8954-8966.	1.7	122
89	5-HT1A Receptors, Gene Repression, and Depression: Guilt by Association. Neuroscientist, 2004, 10, 575-593.	2.6	223
90	The Proapoptotic Gene SIVA Is a Direct Transcriptional Target for the Tumor Suppressors p53 and E2F1. Journal of Biological Chemistry, 2004, 279, 28706-28714.	1.6	73

#	ARTICLE	IF	CITATIONS
91	RGS17/RGSZ2, a Novel Regulator of Gi/o, Gz, and Gq Signaling. <i>Journal of Biological Chemistry</i> , 2004, 279, 26314-26322.	1.6	78
92	Cell type-dependent recruitment of trichostatin A-sensitive repression of the human 5-HT1A receptor gene. <i>Journal of Neurochemistry</i> , 2004, 88, 857-868.	2.1	45
93	Forskolin-resistant Y1 adrenal cell mutants are deficient in adenylyl cyclase type 4. <i>Molecular and Cellular Endocrinology</i> , 2004, 214, 155-165.	1.6	10
94	Expression of adenylyl cyclase-4 (AC-4) in Y1 and forskolin-resistant adrenal cells. <i>Molecular and Cellular Endocrinology</i> , 2004, 215, 101-108.	1.6	7
95	5-HT1A-mediated promotion of mitogen-activated T and B cell survival and proliferation is associated with increased translocation of NF- κ B to the nucleus. <i>Brain, Behavior, and Immunity</i> , 2004, 18, 24-34.	2.0	65
96	Association of the C(-1019)G 5-HT1A functional promoter polymorphism with antidepressant response. <i>International Journal of Neuropsychopharmacology</i> , 2004, 7, 501-506.	1.0	175
97	Diacylglycerol and ceramide formation induced by dopamine D2S receptors via G β γ -subunits in Balb/c-3T3 cells. <i>American Journal of Physiology - Cell Physiology</i> , 2003, 284, C640-C648.	2.1	9
98	Impaired Repression at a 5-Hydroxytryptamine 1A Receptor Gene Polymorphism Associated with Major Depression and Suicide. <i>Journal of Neuroscience</i> , 2003, 23, 8788-8799.	1.7	662
99	Freud-1: A Neuronal Calcium-Regulated Repressor of the 5-HT1A Receptor Gene. <i>Journal of Neuroscience</i> , 2003, 23, 7415-7425.	1.7	94
100	Editorial: Dopamine-D2-Mediated Inhibition of TRH-Induced PLC Activation in Pituitary Cells-Direct or Indirect?. <i>Endocrinology</i> , 2002, 143, 744-746.	1.4	3
101	G Protein Preferences for Dopamine D2 Inhibition of Prolactin Secretion and DNA Synthesis in GH4 Pituitary Cells. <i>Molecular Endocrinology</i> , 2002, 16, 1903-1911.	3.7	20
102	Dopamine-D2S Receptor Inhibition of Calcium Influx, Adenylyl Cyclase, and Mitogen-Activated Protein Kinase in Pituitary Cells: Distinct G β γ and G β γ Requirements. <i>Molecular Endocrinology</i> , 2002, 16, 2393-2404.	3.7	38
103	Growth Hormone-induced Diacylglycerol and Ceramide Formation via G β γ and G β γ in GH4 Pituitary Cells. <i>Journal of Biological Chemistry</i> , 2002, 277, 48427-48433.	1.6	8
104	G protein specificity. <i>Cellular Signalling</i> , 2002, 14, 407-418.	1.7	156
105	Identification of an Endogenous 5-Hydroxytryptamine2A Receptor in NIH-3T3 Cells: Agonist-Induced Down-Regulation Involves Decreases in Receptor RNA and Number. <i>Journal of Neurochemistry</i> , 2002, 68, 1998-2011.	2.1	22
106	TATA-Driven Transcriptional Initiation and Regulation of the Rat Serotonin 5-HT1A Receptor Gene. <i>Journal of Neurochemistry</i> , 2002, 72, 2238-2247.	2.1	37
107	A critical protein kinase C phosphorylation site on the 5-HT1A receptor controlling coupling to N-type calcium channels. <i>Journal of Physiology</i> , 2002, 538, 41-51.	1.3	27
108	Receptor signaling and structure: insights from serotonin-1 receptors. <i>Trends in Endocrinology and Metabolism</i> , 2001, 12, 453-460.	3.1	67

#	ARTICLE	IF	CITATIONS
109	Heterodimerization of Mineralocorticoid and Glucocorticoid Receptors at a Novel Negative Response Element of the 5-HT1A Receptor Gene. <i>Journal of Biological Chemistry</i> , 2001, 276, 14299-14307.	1.6	151
110	Transcriptional Mechanisms for Induction of 5-HT1A Receptor mRNA and Protein in Activated B and T Lymphocytes. <i>Journal of Biological Chemistry</i> , 2001, 276, 4382-4388.	1.6	62
111	APAF1 is a key transcriptional target for p53 in the regulation of neuronal cell death. <i>Journal of Cell Biology</i> , 2001, 155, 207-216.	2.3	184
112	Distinct Roles for G α i 2 and G α i 3 in Signaling to DNA Synthesis and G α i 3 in Cellular Transformation by Dopamine D2S Receptor Activation in BALB/c 3T3 Cells. <i>Molecular and Cellular Biology</i> , 2000, 20, 1497-1506.	1.1	42
113	Novel Dual Repressor Elements for Neuronal Cell-specific Transcription of the Rat 5-HT1A Receptor Gene. <i>Journal of Biological Chemistry</i> , 2000, 275, 8161-8168.	1.6	62
114	Receptor Selectivity of the Cloned Opossum G Protein-Coupled Receptor Kinase 2 (GRK2) in Intact Opossum Kidney Cells: Role in Desensitization of Endogenous β 2C-Adrenergic but Not Serotonin 1B Receptors. <i>Molecular Endocrinology</i> , 1999, 13, 138-147.	3.7	19
115	Distinct Roles for G α i 2, G α i 3, and G α i 3 in Modulation of Forskolin- or Gs-mediated cAMP Accumulation and Calcium Mobilization by Dopamine D2S Receptors. <i>Journal of Biological Chemistry</i> , 1999, 274, 9238-9245.	1.6	76
116	Stimulation of cAMP Synthesis by Gi-coupled Receptors upon Ablation of Distinct G α i Protein Expression. <i>Journal of Biological Chemistry</i> , 1999, 274, 16444-16450.	1.6	60
117	Constitutive Gi2-dependent Activation of Adenylyl Cyclase Type II by the 5-HT1A Receptor. <i>Journal of Biological Chemistry</i> , 1999, 274, 35469-35474.	1.6	58
118	Receptor Selectivity of the Cloned Opossum G Protein-Coupled Receptor Kinase 2 (GRK2) in Intact Opossum Kidney Cells: Role in Desensitization of Endogenous β 2C-Adrenergic but Not Serotonin 1B Receptors. <i>Molecular Endocrinology</i> , 1999, 13, 138-147.	3.7	11
119	A Putative alpha-helical Gbetagamma-coupling Domain in the Second Intracellular Loop of the 5-HT1A Receptor. <i>Annals of the New York Academy of Sciences</i> , 1998, 861, 146-161.	1.8	22
120	Endogenous serotonin-2A and -2C receptors in Balb/c-3T3 cells revealed in serotonin-free medium. <i>Biochemical Pharmacology</i> , 1998, 56, 1347-1357.	2.0	36
121	Selective Antagonism of Receptor Signaling Using Antisense RNA to Deplete G-Protein Subunits. , 1998, 84, 107-122.		0
122	Homology Cloning of cDNA or Genomic DNA. <i>Current Protocols in Neuroscience</i> , 1997, 00, 4.1.1-4.1.6.	2.6	0
123	A Conserved Threonine Residue in the Second Intracellular Loop of the 5-Hydroxytryptamine 1A Receptor Directs Signaling Specificity. <i>Molecular Pharmacology</i> , 1997, 52, 164-171.	1.0	47
124	A Novel cdc2-Related Protein Kinase Expressed in the Nervous System. <i>Journal of Neurochemistry</i> , 1997, 69, 348-364.	2.1	30
125	Mechanisms of Dopaminergic Regulation of Prolactin Secretion. , 1997, , 359-381.		6
126	Correspondence. <i>Neuropsychopharmacology</i> , 1996, 15, 213-214.	2.8	8

#	ARTICLE	IF	CITATIONS
127	The 5-HT1A receptor: Signaling, desensitization, and gene transcription. <i>Neuropsychopharmacology</i> , 1996, 14, 19-25.	2.8	108
128	Antisense knockouts: molecular scalpels for the dissection of signal transduction. <i>Trends in Pharmacological Sciences</i> , 1994, 15, 250-254.	4.0	48
129	Heterologous Expression of G Protein-Linked Receptors in Pituitary and Fibroblast Cell Lines. <i>Vitamins and Hormones</i> , 1994, 48, 59-109.	0.7	23
130	Deletions of the Synenkephalin Domain Which Do Not Alter Cell-Specific Proteolytic Processing or Secretory Targeting of Human Proenkephalin. <i>Journal of Neurochemistry</i> , 1993, 60, 1325-1334.	2.1	4
131	Cholera toxin-sensitive 3',5'-cyclic adenosine monophosphate and calcium signals of the human dopamine-D1 receptor: selective potentiation by protein kinase A. <i>Molecular Endocrinology</i> , 1992, 6, 1815-1824.	3.7	42
132	Molecular biology of the 5-HT1A receptor: Low-stringency cloning and eukaryotic expression. <i>Journal of Chemical Neuroanatomy</i> , 1992, 5, 283-288.	1.0	8
133	Differential Sensitivity of the Short and Long Human Dopamine D ₂ Receptor Subtypes to Protein Kinase C. <i>Journal of Neurochemistry</i> , 1992, 59, 2311-2317.	2.1	68
134	Cloning and expression of a rat D2 dopamine receptor cDNA. <i>Nature</i> , 1988, 336, 783-787.	13.7	1,121
135	The next frontier in the molecular biology of the opioid system. <i>Molecular Neurobiology</i> , 1987, 1, 373-391.	1.9	15