Gerardo Gold Gold-Bouchot

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3262857/publications.pdf

Version: 2024-02-01

60 papers

1,285 citations

304602 22 h-index 33 g-index

60 all docs

60 does citations

60 times ranked

1446 citing authors

#	Article	IF	Citations
1	Determination of esterase activity and characterization of cholinesterases in the reef fish Haemulon plumieri. Ecotoxicology and Environmental Safety, 2008, 71, 787-797.	2.9	65
2	Organochlorine pesticides in soils and air of southern Mexico: Chemical profiles and potential for soil emissions. Atmospheric Environment, 2008, 42, 7737-7745.	1.9	61
3	Characterization of cholinesterase activity from different tissues of Nile tilapia (Oreochromis) Tj ETQq1 1 0.7843	14 rgBT /C	Overlock 10 T
4	Large-scale environmental influences on the benthic macroinfauna of the southern Gulf of Mexico. Estuarine, Coastal and Shelf Science, 2003, 58, 825-841.	0.9	50
5	Histopathological effects of petroleum hydrocarbons and heavy metals on the American oyster (Crassostrea virginica) from Tabasco, Mexico. Marine Pollution Bulletin, 1995, 31, 439-445.	2.3	47
6	Lead contamination in the Mexican Caribbean recorded by the coral Montastraea annularis (Ellis and) Tj ETQq0 0	0 rgBT /O	verlock 10 Tf
7	Cell wall composition affects Cd2+ accumulation and intracellular thiol peptides in marine red algae. Aquatic Toxicology, 2007, 81, 65-72.	1.9	46
8	Organochlorine Pesticides and Polychlorinated Biphenyls Levels in Human Milk from Chelem, Yucatán, MÁ©xico. Bulletin of Environmental Contamination and Toxicology, 2008, 80, 255-259.	1.3	45
9	Passive Air Sampling of Organochlorine Pesticides in Mexico. Environmental Science & Emp; Technology, 2009, 43, 704-710.	4.6	45
10	Polynuclear Aromatic Hydrocarbons in American Oysters Crassostrea virginica from the Terminos Lagoon, Campeche, Mexico. Marine Pollution Bulletin, 1999, 38, 637-645.	2.3	42
11	Anaerobic–Aerobic Biodegradation of DDT (Dichlorodiphenyl Trichloroethane) in Soils. Bulletin of Environmental Contamination and Toxicology, 1999, 63, 219-225.	1.3	36
12	Effect of Pyrene on Hepatic Cytochrome P450 1A (CYP1A) Expression in Nile Tilapia (Oreochromis) Tj ETQq0 0 0) rgBJ /Ove	erlock 10 Tf 50
13	CICHLIDOGYRUS SCLEROSUS (MONOGENEA: ANCYROCEPHALINAE) AND ITS HOST, THE NILE TILAPIA (OREOCHROMIS NILOTICUS), AS BIOINDICATORS OF CHEMICAL POLLUTION. Journal of Parasitology, 2007, 93, 1097-1106.	0.3	35
14	Environmental monitoring using acetylcholinesterase inhibition in vitro. A case study in two Mexican lagoons. Marine Environmental Research, 2000, 50, 357-360.	1.1	33
15	Potential interactions between metazoan parasites of the Mayan catfish Ariopsis assimilis and chemical pollution in Chetumal Bay, Mexico. Journal of Helminthology, 2003, 77, 173-184.	0.4	33
16	The checkered puffer (Spheroides testudineus) and its helminths as bioindicators of chemical pollution in Yucatan coastal lagoons. Science of the Total Environment, 2009, 407, 2315-2324.	3.9	33
17	Persistent organic pollutants and histological lesions in Mayan catfish Ariopsis assimilis from the Bay of Chetumal, Mexico. Marine Pollution Bulletin, 2004, 48, 263-269.	2.3	30

o,pâ \in 2-DDT induction of vitellogenesis and its inhibition by tamoxifen in Nile tilapia (Oreochromis) Tj ETQq0 0 0 rgBT.1/Overlock 10 Tf 50 tilapia (Oreochromis)

18

#	Article	IF	CITATIONS
19	Endocrine disruption mechanism of o,p′-DDT in mature male tilapia (Oreochromis niloticus). Toxicology and Applied Pharmacology, 2007, 221, 158-167.	1.3	26
20	Biomarkers and pollutants in the Nile Tilapia, Oreochromis niloticus, in four lakes from San Miguel, Chiapas, Mexico. International Journal of Environment and Pollution, 2006, 26, 129.	0.2	24
21	Pollutants and biomarker responses in two reef fish species (Haemulon aurolineatum and Ocyurus) Tj ETQq $1\ 1$	0.784314 2.3	rgBT/Overloc
22	The pink shrimpFarfantepenaeus duorarum, its symbionts and helminths as bioindicators of chemical pollution in Campeche Sound, Mexico. Journal of Helminthology, 2006, 80, 159-174.	0.4	23
23	Hydrocarbon Concentrations in Oysters (Crassostrea virginica) and Recent Sediments from Three Coastal Lagoons in Tabasco, Mexico. Bulletin of Environmental Contamination and Toxicology, 1997, 59, 430-437.	1.3	21
24	Towards a coastal condition assessment and monitoring of the Gulf of Mexico Large Marine Ecosystem (GoM LME): Terminos Lagoon pilot site. Environmental Development, 2013, 7, 72-79.	1.8	21
25	Thiol peptides induction in the seagrass Thalassia testudinum (Banks ex König) in response to cadmium exposure. Aquatic Toxicology, 2008, 86, 12-19.	1.9	20
26	Benthic infauna variability in relation to environmental factors and organic pollutants in tropical coastal lagoons from the northern Yucatan Peninsula. Marine Pollution Bulletin, 2012, 64, 2725-2733.	2.3	20
27	Biological responses of shoal flounder (Syacium gunteri) to toxic environmental pollutants from the southern Gulf of Mexico. Environmental Pollution, 2020, 258, 113669.	3.7	20
28	Chlorinated pesticides in the Rio Palizada, Campeche, Mexico. Marine Pollution Bulletin, 1993, 26, 648-650.	2.3	19
29	Organochlorine pesticide residue concentrations in biota and sediments from R� Palizada, Mexico. Bulletin of Environmental Contamination and Toxicology, 1995, 54, 554-61.	1.3	19
30	Hydrocarbon and Organochlorine Residue Concentrations in Sediments from Bay of Chetumal, Mexico. Bulletin of Environmental Contamination and Toxicology, 1998, 61, 80-87.	1.3	17
31	Toxicity of sediments from BahıÌa de Chetumal, México, as assessed by hepatic EROD induction and histology in nile tilapia Oreochromis niloticus. Marine Environmental Research, 2000, 50, 385-391.	1.1	17
32	Biological effects of environmental pollutants in American Oyster, Crassostrea virginica: a field study in Laguna de Terminos, Mexico. International Journal of Environment and Health, 2007, 1, 171.	0.3	17
33	The characterization of cytosolic glutathione transferase from four species of sea turtles: Loggerhead (Caretta caretta), green (Chelonia mydas), olive ridley (Lepidochelys olivacea), and hawksbill (Eretmochelys imbricata). Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology, 2009, 150, 279-284.	1.3	17
34	Effects of oil spill related chemical pollution on helminth parasites in Mexican flounder Cyclopsetta chittendeni from the Campeche Sound, Gulf of Mexico. Ecotoxicology and Environmental Safety, 2015, 119, 162-169.	2.9	16
35	Vitellogenin Induction and Increased Plasma $17\hat{l}^2$ -Estradiol Concentrations in Male Nile Tilapia, Oreochromis niloticus, Exposed to Organochlorine Pollutants and Polycyclic Aromatics Hydrocarbons. Bulletin of Environmental Contamination and Toxicology, 2008, 81, 543-547.	1.3	14
36	A mesocosm experiment to determine half-lives of individual hydrocarbons in simulated oil spill scenarios with and without the dispersant, Corexit. Marine Pollution Bulletin, 2020, 151, 110804.	2.3	13

#	Article	IF	CITATIONS
37	Hydrocarbon concentrations in the American oyster, Crassostrea virginica, in Laguna de Terminos, Campeche, Mexico. Bulletin of Environmental Contamination and Toxicology, 1995, 54, 222-7.	1.3	12
38	Preliminary studies of biochemical changes (ethoxycoumarin O-deethylase activities and vitellogenin) Tj ETQq0 O Gulf of Mexico. Ecotoxicology and Environmental Safety, 2005, 61, 98-104.	0 rgBT /C 2.9	overlock 10 Tf 12
39	Biodegradation of hexadecane using sediments from rivers and lagoons of the Southern Gulf of Mexico. Marine Pollution Bulletin, 2018, 128, 202-207.	2.3	12
40	Trace metals in the American oyster, Crassostrea virginica, and sediments from the coastal lagoons Mecoacan, Carmen and Machona, Tabasco, Mexico. Chemosphere, 1997, 34, 2437-2450.	4.2	11
41	Heavy Metals and Hydrocarbons in Sediments from Three Lakes from San Miguel, Chiapas, Mexico. Bulletin of Environmental Contamination and Toxicology, 2004, 73, 762-9.	1.3	10
42	Transitional carbonate-terrigenous shelf sub-environments inferred from textural characteristics of surficial sediments in the Southern Gulf of Mexico. Continental Shelf Research, 2005, 25, 1836-1852.	0.9	10
43	Ecotoxicological effects of POPs on ariidae Ariopsis felis (Linnaeus, 1766) from three coastal ecosystems in the Southern Gulf of Mexico and Yucatan Peninsula. Journal of Environmental Science and Health - Part A Toxic/Hazardous Substances and Environmental Engineering, 2007, 42, 1513-1520.	0.9	10
44	Temporal Variation of Persistent Organic Pollutant (POP) Residue Concentrations in Sediments from the Bay of Chetumal, Mexico. Bulletin of Environmental Contamination and Toxicology, 2007, 79, 141-146.	1.3	10
45	Environmental and anthropogenic factors affecting the probability of occurrence of Oncomegas wageneri (Cestoda: Trypanorhyncha) in the southern Gulf of Mexico. Parasites and Vectors, 2015, 8, 609.	1.0	10
46	Effect of the use of Bacillus spp. on the characteristics of dissolved fluorescent organic matter and the phytochemical quality of Stevia rebaudiana grown in a recirculating aquaponic system. Environmental Science and Pollution Research, 2021, 28, 36326-36343.	2.7	9
47	The metazoan parasite communities of the shoal flounder (Syacium gunteri) as bioindicators of chemical contamination in the southern Gulf of Mexico. Parasites and Vectors, 2014, 7, 541.	1.0	8
48	Inter-laboratory calibration of estimated oil equivalent (EOE) concentrations of a water accommodated fraction (WAF) of oil and a chemically enhanced WAF (CEWAF). Heliyon, 2019, 5, e01174.	1.4	7
49	An aromatase inhibitor and tamoxifen decrease plasma levels of o,p′-DDT and its metabolites in Nile tilapia (Oreochromis niloticus). Marine Environmental Research, 2004, 58, 337-342.	1.1	6
50	Polybrominated diphenyl ethers (PBDE) and hexabromocyclododecane (HBCD) in liver of checkered puffer (Sphoeroides testudineus) from Ria Lagartos, Yucatan, Mexico. Marine Pollution Bulletin, 2019, 146, 488-492.	2.3	6
51	Spatial and temporal variability of sea breezes and synoptic influences over the surface wind field of the Yucat \tilde{A}_i n Peninsula. , 2020, 33, 123-142.		6
52	Trace Metals in Sediments from Bahia de Chetumal, Mexico. Bulletin of Environmental Contamination and Toxicology, 2003, 70, 1228-1234.	1.3	5
53	Brominated Flame Retardants in Sediments of Four Coastal Lagoons of Yucatan, Mexico. Bulletin of Environmental Contamination and Toxicology, 2018, 101, 160-165.	1.3	5
54	Chromophoric dissolved organic matter (CDOM) in a subtropical estuary (Galveston Bay, USA) and the impact of Hurricane Harvey. Environmental Science and Pollution Research, 2021, 28, 53045-53057.	2.7	5

#	Article	IF	CITATIONS
55	PERSISTENT ORGANIC POLLUTANTS IN SERUM AND BREAST MILK OF FERTILE-AGED WOMEN. Revista Internacional De Contaminacion Ambiental, 2019, 35, 281-293.	0.1	4
56	Hydrocarbon concentrations in sediments and clams (Rangia cuneata) in Laguna de Pom, Mexico. Bulletin of Environmental Contamination and Toxicology, 1994, 52, 39-45.	1.3	3
57	Oil Pollution in the Southern Gulf of Mexico: Field and Laboratory Studies. Bulletin of Environmental Contamination and Toxicology, 2022, 108, 1-2.	1.3	2
58	Tamoxifen Affects the Toxicokinetics of o,p′-DDT in Male Nile Tilapia (Oreochromis niloticus). Bulletin of Environmental Contamination and Toxicology, 2010, 85, 545-549.	1.3	1
59	Microbial Activity in Marine Sediments Exposed to Hexadecane: A Laboratory Study. Clean - Soil, Air, Water, 2017, 45, 1700531.	0.7	1
60	Characteristics of Chromophoric Dissolved Organic Matter (CDOM) Produced by Heterotrophic Bacteria Isolated from Aquaculture Systems. Journal of Marine Science and Engineering, 2022, 10, 672.	1.2	1