Luis H Canani

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3263856/publications.pdf

Version: 2024-02-01

101384 5,983 126 36 citations h-index papers

g-index 142 142 142 8219 docs citations times ranked citing authors all docs

82410

72

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Diabetic Nephropathy: Diagnosis, Prevention, and Treatment. Diabetes Care, 2005, 28, 164-176. | 4.3 | 1,347 |
| 2 | Genome-Wide Association Scan for Diabetic Nephropathy Susceptibility Genes in Type 1 Diabetes. Diabetes, 2009, 58, 1403-1410. | 0.3 | 259 |
| 3 | Standards of Medical Care in Diabetes-2008: Response to Hirsch, Inzucchi, and Kirkman. Diabetes Care, 2008, 31, e44-e44. | 4.3 | 234 |
| 4 | Oral Semaglutide Versus Empagliflozin in Patients With Type 2 Diabetes Uncontrolled on Metformin: The PIONEER 2 Trial. Diabetes Care, 2019, 42, 2272-2281. | 4.3 | 225 |
| 5 | Diabetic Retinopathy Predicts All-Cause Mortality and Cardiovascular Events in Both Type 1 and 2 Diabetes. Diabetes Care, 2011, 34, 1238-1244. | 4.3 | 194 |
| 6 | The Type 2 Deiodinase A/G (Thr92Ala) Polymorphism Is Associated with Decreased Enzyme Velocity and Increased Insulin Resistance in Patients with Type 2 Diabetes Mellitus. Journal of Clinical Endocrinology and Metabolism, 2005, 90, 3472-3478. | 1.8 | 191 |
| 7 | Coronary artery calcium score prediction of all cause mortality and cardiovascular events in people with type 2 diabetes: systematic review and meta-analysis. BMJ, The, 2013, 346, f1654-f1654. | 3.0 | 140 |
| 8 | Diabetic nephropathy. Diabetology and Metabolic Syndrome, 2009, 1, 10. | 1.2 | 101 |
| 9 | Meta-Analysis Reveals the Association of Common Variants in the Uncoupling Protein (UCP) 1–3 Genes with Body Mass Index Variability. PLoS ONE, 2014, 9, e96411. | 1.1 | 99 |
| 10 | Masked Hypertension, Urinary Albumin Excretion Rate, and Echocardiographic Parameters in Putatively Normotensive Type 2 Diabetic Patients. Diabetes Care, 2007, 30, 1255-1260. | 4.3 | 96 |
| 11 | The role of the uncoupling protein 1 (UCP1) on the development of obesity and type 2 diabetes mellitus. Arquivos Brasileiros De Endocrinologia E Metabologia, 2012, 56, 215-225. | 1.3 | 92 |
| 12 | Nitric oxide system and diabetic nephropathy. Diabetology and Metabolic Syndrome, 2014, 6, 17. | 1.2 | 79 |
| 13 | The role of uncoupling protein 2 (UCP2) on the development of type 2 diabetes mellitus and its chronic complications. Arquivos Brasileiros De Endocrinologia E Metabologia, 2011, 55, 239-248. | 1.3 | 78 |
| 14 | The Human Peroxisome Proliferator-Activated Receptor Â2 (PPARÂ2) Pro12Ala Polymorphism Is Associated With Decreased Risk of Diabetic Nephropathy in Patients With Type 2 Diabetes. Diabetes, 2003, 52, 3010-3013. | 0.3 | 76 |
| 15 | Nitric oxide levels in patients with diabetes mellitus: A systematic review and meta-analysis. Nitric Oxide - Biology and Chemistry, 2016, 61, 1-9. | 1.2 | 71 |
| 16 | Clinical and Laboratory Profile of Patients With Type 2 Diabetes With Low Glomerular Filtration Rate and Normoalbuminuria. Diabetes Care, 2007, 30, 1998-2000. | 4.3 | 68 |
| 17 | Evidence for different susceptibility genes for proteinuria and ESRD in type 2 diabetes. Advances in Chronic Kidney Disease, 2005, 12, 155-169. | 0.6 | 66 |
| 18 | Prevalence of adults with type 1 diabetes who meet the goals of care in daily clinical practice: A nationwide multicenter study in Brazil. Diabetes Research and Clinical Practice, 2012, 97, 63-70. | 1.1 | 63 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Associations between UCP1 -3826A/G, UCP2 -866G/A, Ala55Val and Ins/Del, and UCP3 -55C/T Polymorphisms and Susceptibility to Type 2 Diabetes Mellitus: Case-Control Study and Meta-Analysis. PLoS ONE, 2013, 8, e54259. | 1.1 | 58 |
| 20 | The â^374A allele of the receptor for advanced glycation end products gene is associated with a decreased risk of ischemic heart disease in African-Brazilians with type 2 diabetes. Molecular Genetics and Metabolism, 2005, 85, 149-156. | 0.5 | 57 |
| 21 | Minor Effect of GLUT1 Polymorphisms on Susceptibility to Diabetic Nephropathy in Type 1 Diabetes. Diabetes, 2002, 51, 2264-2269. | 0.3 | 55 |
| 22 | Endothelin-1 levels and albuminuria in patients with type 2 diabetes mellitus. Diabetes Research and Clinical Practice, 2008, 80, 299-304. | 1.1 | 53 |
| 23 | Polymorphism in Ecto-Nucleotide Pyrophosphatase/Phosphodiesterase 1 Gene (ENPP1/PC-1) and Early Development of Advanced Diabetic Nephropathy in Type 1 Diabetes. Diabetes, 2002, 51, 1188-1193. | 0.3 | 51 |
| 24 | High-Density Single Nucleotide Polymorphism Genome-Wide Linkage Scan for Susceptibility Genes for Diabetic Nephropathy in Type 1 Diabetes. Diabetes, 2008, 57, 2519-2526. | 0.3 | 51 |
| 25 | Polymorphisms of the <i>UCP2</i> gene are associated with proliferative diabetic retinopathy in patients with diabetes mellitus. Clinical Endocrinology, 2010, 72, 612-619. | 1.2 | 51 |
| 26 | MicroRNA expression profile in plasma from type 1 diabetic patients: Case-control study and bioinformatic analysis. Diabetes Research and Clinical Practice, 2018, 141, 35-46. | 1.1 | 49 |
| 27 | Polymorphisms in genes encoding miR-155 and miR-146a are associated with protection to type 1 diabetes mellitus. Acta Diabetologica, 2017, 54, 433-441. | 1.2 | 47 |
| 28 | Plantar thermography is useful in the early diagnosis of diabetic neuropathy. Clinics, 2012, 67, 1419-1425. | 0.6 | 46 |
| 29 | The Fatty Acid-Binding Protein-2 A54T Polymorphism Is Associated With Renal Disease in Patients With Type 2 Diabetes. Diabetes, 2005, 54, 3326-3330. | 0.3 | 45 |
| 30 | Polymorphisms in the TLR3 gene are associated with risk for type 1 diabetes mellitus. European Journal of Endocrinology, 2014, 170, 519-527. | 1.9 | 44 |
| 31 | Circulating miRNAs in diabetic kidney disease: case–control study and in silico analyses. Acta Diabetologica, 2019, 56, 55-65. | 1.2 | 41 |
| 32 | The Catalase –262C/T Promoter Polymorphism and Diabetic Complications in Caucasians with Type 2 Diabetes. Disease Markers, 2006, 22, 355-359. | 0.6 | 40 |
| 33 | Association of the UCP polymorphisms with susceptibility to obesity: case–control study and meta-analysis. Molecular Biology Reports, 2014, 41, 5053-5067. | 1.0 | 40 |
| 34 | The role of progranulin in diabetes and kidney disease. Diabetology and Metabolic Syndrome, 2015, 7, 117. | 1.2 | 39 |
| 35 | Determinants of body weight regulation in humans. Archives of Endocrinology and Metabolism, 2016, 60, 152-162. | 0.3 | 39 |
| 36 | Functional Vascular Endothelial Growth Factor -634G>C SNP Is Associated With Proliferative Diabetic Retinopathy: A case-control study in a Brazilian population of European ancestry. Diabetes Care, 2007, 30, 275-279. | 4.3 | 38 |

| # | Article | IF | Citations |
|----|--|-----|-----------|
| 37 | Choroidal Thickness in Patients with Diabetes and Microalbuminuria. Ophthalmology, 2014, 121, 2071-2073. | 2.5 | 38 |
| 38 | Impact of White-Coat Hypertension on Microvascular Complications in Type 2 Diabetes. Diabetes Care, 2008, 31, 2233-2237. | 4.3 | 37 |
| 39 | Relationship between adherence to diet, glycemic control and cardiovascular risk factors in patients with type 1 diabetes: a nationwide survey in Brazil. Nutrition Journal, 2014, 13, 19. | 1.5 | 37 |
| 40 | The <i>UCP1 </i> according to the second state of the second seco | | 35 |
| 41 | Plasma levels of miRâ€29b and miRâ€200b in type 2 diabetic retinopathy. Journal of Cellular and Molecular Medicine, 2019, 23, 1280-1287. | 1.6 | 34 |
| 42 | Identification of a Common Risk Haplotype for Diabetic Nephropathy at the Protein Kinase C- \hat{l}^21 (PRKCB1) Gene Locus. Journal of the American Society of Nephrology: JASN, 2003, 14, 2015-2024. | 3.0 | 33 |
| 43 | Effect of dietary lipids on circulating adiponectin: a systematic review with meta-analysis of randomised controlled trials. British Journal of Nutrition, 2014, 112, 1235-1250. | 1.2 | 33 |
| 44 | Relationship of endothelial nitric oxide synthase (<i>eNOS</i>) gene polymorphisms with diabetic retinopathy in Caucasians with type 2 diabetes. Ophthalmic Genetics, 2012, 33, 23-27. | 0.5 | 31 |
| 45 | Irisin-encoding gene (FNDC5) variant is associated with changes in blood pressure and lipid profile in type 2 diabetic women but not in men. Metabolism: Clinical and Experimental, 2015, 64, 952-957. | 1.5 | 31 |
| 46 | Association of eNOS gene polymorphisms with renal disease in Caucasians with type 2 diabetes. Diabetes Research and Clinical Practice, 2011, 91, 353-362. | 1.1 | 30 |
| 47 | Physical Therapy Reduces Bone Resorption and Increases Bone Formation in Preterm Infants. American Journal of Perinatology, 2012, 29, 573-578. | 0.6 | 29 |
| 48 | Regional differences in clinical care among patients with type 1 diabetes in Brazil: Brazilian Type 1 Diabetes Study Group. Diabetology and Metabolic Syndrome, 2012, 4, 44. | 1.2 | 29 |
| 49 | Messenger RNA levels of podocyte-associated proteins in subjects with different degrees of glucose tolerance with or without nephropathy. BMC Nephrology, 2013, 14, 214. | 0.8 | 29 |
| 50 | The TCF7L2 rs7903146 (C/T) polymorphism is associated with risk to type 2 diabetes mellitus in Southern-Brazil. Arquivos Brasileiros De Endocrinologia E Metabologia, 2014, 58, 918-925. | 1.3 | 29 |
| 51 | MiR-30e-5p and MiR-15a-5p Expressions in Plasma and Urine of Type 1 Diabetic Patients With Diabetic Kidney Disease. Frontiers in Genetics, 2019, 10, 563. | 1.1 | 29 |
| 52 | Masked hypertension, nocturnal blood pressure and retinopathy in normotensive patients with type 1 diabetes. Diabetes Research and Clinical Practice, 2010, 87, 240-245. | 1.1 | 28 |
| 53 | Association of genetic variants in the promoter region of genes encoding p22phox (CYBA) and glutamate cysteine ligase catalytic subunit (GCLC) and renal disease in patients with type 1 diabetes mellitus. BMC Medical Genetics, 2011, 12, 129. | 2.1 | 28 |
| 54 | Polymorphisms of the UCP2 Gene Are Associated with Glomerular Filtration Rate in Type 2 Diabetic Patients and with Decreased UCP2 Gene Expression in Human Kidney. PLoS ONE, 2015, 10, e0132938. | 1.1 | 27 |

| # | Article | IF | Citations |
|----|--|------------|-------------------------|
| 55 | The â^106CC genotype of the aldose reductase gene is associated with an increased risk of proliferative diabetic retinopathy in Caucasian-Brazilians with type 2 diabetes. Molecular Genetics and Metabolism, 2006, 88, 280-284. | 0.5 | 26 |
| 56 | Major components of metabolic syndrome and adiponectin levels: a cross-sectional study. Diabetology and Metabolic Syndrome, 2014, 6, 26. | 1.2 | 26 |
| 57 | Cataract and type 1 diabetes mellitus. Diabetes Research and Clinical Practice, 2008, 82, 324-328. | 1.1 | 25 |
| 58 | Endothelin-1 and Endothelin A Receptor Immunoreactivity Is Increased in Patients with Diabetic Nephropathy. Renal Failure, 2012, 34, 308-315. | 0.8 | 25 |
| 59 | Urinary Albumin Excretion Rate Is Associated With Increased Ambulatory Blood Pressure in Normoalbuminuric Type 2 Diabetic Patients. Diabetes Care, 2005, 28, 1724-1729. | 4.3 | 23 |
| 60 | Prevalence of 15 mitochondrial DNA mutations among type 2 diabetic patients with or without clinical characteristics of maternally inherited diabetes and deafness. Arquivos Brasileiros De Endocrinologia E Metabologia, 2008, 52, 1228-1235. | 1.3 | 23 |
| 61 | Prevalence and characteristics of diabetic polyneuropathy in Passo Fundo, South of Brazil. Arquivos Brasileiros De Endocrinologia E Metabologia, 2007, 51, 987-992. | 1.3 | 22 |
| 62 | Genetics of diabetic nephropathy. Arquivos Brasileiros De Endocrinologia E Metabologia, 2010, 54, 253-261. | 1.3 | 22 |
| 63 | Association of ADIPOQ variants, total and high molecular weight adiponectin levels with coronary artery disease in diabetic and non-diabetic Brazilian subjects. Journal of Diabetes and Its Complications, 2012, 26, 94-98. | 1.2 | 22 |
| 64 | The -308G>A Polymorphism of the TNF Gene Is Associated With Proliferative Diabetic Retinopathy in Caucasian Brazilians With Type 2 Diabetes. Investigative Ophthalmology and Visual Science, 2015, 56, 1184-1190. | 3.3 | 22 |
| 65 | Familial history of type 2 diabetes in patients from Southern Brazil and its influence on the clinical characteristics of this disease. Arquivos Brasileiros De Endocrinologia E Metabologia, 2006, 50, 862-868. | 1.3 | 22 |
| 66 | Toll-like receptor 3 (TLR3) and the development of type 1 diabetes mellitus. Archives of Endocrinology and Metabolism, 2015, 59, 4-12. | 0.3 | 21 |
| 67 | COL18A1 is highly expressed during human adipocyte differentiation and the SNP c.1136C > T in its "frizzled" motif is associated with obesity in diabetes type 2 patients. Anais Da Academia Brasileira De Ciencias, 2008, 80, 167-177. | 0.3 | 21 |
| 68 | The ACE Insertion/Deletion Polymorphism Is Not Associated With the Metabolic Syndrome (WHO) Tj ETQq0 0 (| O rgBT/Ove | rlo <u>၄</u> k 10 Tf 50 |
| 69 | Cardiovascular autonomic neuropathy in type 2 diabetes mellitus patients with peripheral artery disease. Diabetology and Metabolic Syndrome, 2013, 5, 54. | 1.2 | 20 |
| 70 | The A Allele of the rs1990760 Polymorphism in the IFIH1 Gene Is Associated with Protection for Arterial Hypertension in Type 1 Diabetic Patients and with Expression of This Gene in Human Mononuclear Cells. PLoS ONE, 2013, 8, e83451. | 1.1 | 20 |
| 71 | The Presence of At Least Three Alleles of the <i>ADRB3 </i> Trp64Arg (C/T) and <i>UCP1 </i> â^3826A/G Polymorphisms Is Associated with Protection to Overweight/Obesity and with Higher High-Density Lipoprotein Cholesterol Levels in Caucasian-Brazilian Patients with Type 2 Diabetes. Metabolic Syndrome and Related Disorders, 2014, 12, 16-24. | 0.5 | 19 |
| 72 | Diabetes and cardiovascular disease: from evidence to clinical practice – position statement 2014 of Brazilian Diabetes Society. Diabetology and Metabolic Syndrome, 2014, 6, 58. | 1.2 | 19 |

| # | Article | IF | Citations |
|----|--|-----|-----------|
| 73 | Exendin-4 protects rat islets against loss of viability and function induced by brain death. Molecular and Cellular Endocrinology, 2015, 412, 239-250. | 1.6 | 19 |
| 74 | The role of ecto-nucleotide pyrophosphatase/phosphodiesterase 1 in diabetic nephropathy. Arquivos Brasileiros De Endocrinologia E Metabologia, 2011, 55, 677-685. | 1.3 | 18 |
| 75 | The C Allele of â^'634G/C Polymorphism in the <i>VEGFA</i> Gene Is Associated with Increased <i>VEGFA</i> Gene Expression in Human Retinal Tissue., 2012, 53, 6411. | | 17 |
| 76 | Endothelin-1 gene polymorphisms and diabetic kidney disease in patients with type 2 diabetes mellitus. Diabetology and Metabolic Syndrome, 2015, 7, 103. | 1.2 | 17 |
| 77 | Microvascular Complications of Posttransplant Diabetes Mellitus in Kidney Transplant Recipients: A Longitudinal Study. Journal of Clinical Endocrinology and Metabolism, 2019, 104, 557-567. | 1.8 | 16 |
| 78 | Type 2 Deiodinase Thr92Ala Polymorphism Is Not Associated With Arterial Hypertension in Type 2 Diabetes Mellitus Patients. Hypertension, 2007, 49, e47; author reply e48. | 1.3 | 15 |
| 79 | Contrasting effects of preexisting hyperglycemia and higher body size on hospital mortality in critically ill patients: a prospective cohort study. BMC Endocrine Disorders, 2014, 14, 50. | 0.9 | 15 |
| 80 | Association between the ENPP1 K121Q Polymorphism and Risk of Diabetic Kidney Disease: A Systematic Review and Meta-Analysis. PLoS ONE, 2015, 10, e0118416. | 1.1 | 15 |
| 81 | Serum and Urinary Progranulin in Diabetic Kidney Disease. PLoS ONE, 2016, 11, e0165177. | 1.1 | 15 |
| 82 | A Method for Developing High-Density SNP Maps and Its Application at the Type 1 Angiotensin II Receptor (AGTR1) Locus. Genomics, 2002, 79, 326-332. | 1.3 | 14 |
| 83 | HNF1α mutations are present in half of clinically defined MODY patients in South-Brazilian individuals. Arquivos Brasileiros De Endocrinologia E Metabologia, 2008, 52, 1326-1331. | 1.3 | 14 |
| 84 | The presence of the â^'866A/55Val/Ins haplotype in the uncoupling protein 2 (UCP2) gene is associated with decreased UCP2 gene expression in human retina. Experimental Eye Research, 2012, 94, 49-55. | 1.2 | 14 |
| 85 | Catalase activity, allelic variations in the catalase gene and risk of kidney complications in patients with type 1 diabetes. Diabetologia, 2013, 56, 2733-2742. | 2.9 | 14 |
| 86 | Advances in GLP-1 treatment: focus on oral semaglutide. Diabetology and Metabolic Syndrome, 2021, 13, 99. | 1.2 | 11 |
| 87 | Late afternoon blood pressure increase is associated with diabetic retinopathy in normotensive type 2 diabetes mellitus patients. Diabetes Research and Clinical Practice, 2009, 84, e12-e14. | 1.1 | 10 |
| 88 | Absence of diabetic retinopathy in a patient who has had diabetes mellitus for 69 years, and inadequate glycemic control: case presentation. Diabetology and Metabolic Syndrome, 2009, 1, 13. | 1.2 | 10 |
| 89 | Which patients with diabetes should undergo ambulatory blood pressure monitoring?. Journal of Hypertension, 2011, 29, 236-241. | 0.3 | 10 |
| 90 | The rs2292239 polymorphism in ERBB3 gene is associated with risk for type 1 diabetes mellitus in a Brazilian population. Gene, 2018, 644, 122-128. | 1.0 | 10 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | Interaction of HSD11B1 and H6PD polymorphisms in subjects with type 2 diabetes are protective factors against obesity: a cross-sectional study. Diabetology and Metabolic Syndrome, 2019, 11, 78. | 1.2 | 10 |
| 92 | Risk factors for micro and macrovascular disease in black and white patients with type 2 Diabetes mellitus. Revista Da Associação Médica Brasileira, 2009, 55, 308-314. | 0.3 | 9 |
| 93 | TheAla54ThrPolymorphism of theFABP2Gene Influences the Postprandial Fatty Acids in Patients with Type 2 Diabetes. Journal of Clinical Endocrinology and Metabolism, 2010, 95, 3909-3917. | 1.8 | 9 |
| 94 | Afternoon Blood Pressure Increase: A Blood Pressure Pattern Associated With Microvascular Complications in Type 2 Diabetes Mellitus. American Journal of Hypertension, 2011, 24, 64-69. | 1.0 | 9 |
| 95 | FRMD3 gene: its role in diabetic kidney disease. A narrative review. Diabetology and Metabolic Syndrome, 2015, 7, 118. | 1.2 | 9 |
| 96 | Interleukin-10 â^'1082A > G (rs1800896) polymorphism is associated with diabetic retinopathy in type 2 diabetes. Diabetes Research and Clinical Practice, 2018, 138, 187-192. | 1.1 | 9 |
| 97 | Association between progranulin serum levels and dietary intake. PLoS ONE, 2018, 13, e0202149. | 1.1 | 9 |
| 98 | Determinants of intensive insulin therapeutic regimens in patients with type 1 diabetes: data from a nationwide multicenter survey in Brazil. Diabetology and Metabolic Syndrome, 2014, 6, 67. | 1.2 | 8 |
| 99 | Health-related quality of life in patients with type 1 diabetes mellitus in the different geographical regions of Brazil: data from the Brazilian Type 1 Diabetes Study Group. Diabetology and Metabolic Syndrome, 2015, 7, 87. | 1.2 | 8 |
| 100 | Smoking habit is associated with diabetic macular edema in Type 1 diabetes mellitus patients. Journal of Diabetes and Its Complications, 2008, 22, 430. | 1.2 | 7 |
| 101 | Mutation H63D in the HFE gene confers risk for the development of type 2 diabetes mellitus but not for chronic complications. Journal of Diabetes and Its Complications, 2011, 25, 25-30. | 1.2 | 7 |
| 102 | Association study of sorbitol dehydrogenase â^888G> C polymorphism with type 2 diabetic retinopathy in Caucasian-Brazilians. Experimental Eye Research, 2013, 115, 140-143. | 1.2 | 7 |
| 103 | Polimorfismo K121Q do gene ENPP1 e cardiopatia isquêmica em pacientes com diabete melito. Arquivos Brasileiros De Cardiologia, 2010, 94, 168-173. | 0.3 | 7 |
| 104 | Myocardial Dysfunction in Maternally Inherited Diabetes and Deafness. Diabetes Care, 2003, 26, 1323-1324. | 4.3 | 6 |
| 105 | Linkage disequilibrium with HLA-DRB1-DQB1 haplotypes explains the association of TNF-308G>A variant with type 1 diabetes in a Brazilian cohort. Gene, 2015, 568, 50-54. | 1.0 | 6 |
| 106 | Changes in choroidal thickness and volume are related to urinary albumin excretion in type 2 diabetic patients without retinopathy. Clinical Ophthalmology, 2018, Volume 12, 1405-1411. | 0.9 | 6 |
| 107 | Plasma progranulin levels in obese patients before and after Roux-en-Y gastric bariatric surgery: a longitudinal study. Surgery for Obesity and Related Diseases, 2020, 16, 1655-1660. | 1.0 | 6 |
| 108 | Does bacteriuria interfere with albuminuria measurements of patients with diabetes?. Nephrology Dialysis Transplantation, 2008, 24, 1193-1196. | 0.4 | 5 |

7

| # | Article | lF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Association of –1082A>G Polymorphism in the Interleukin-10 Gene with Estimated Glomerular Filtration Rate in Type 2 Diabetes. Kidney and Blood Pressure Research, 2017, 42, 1164-1174. | 0.9 | 5 |
| 110 | Macular choroidal thickness in pregnant women with type 1, type 2 and gestational diabetes mellitus measured by spectral-domain optical coherence tomography. Clinical Ophthalmology, 2018, Volume 12, 1259-1265. | 0.9 | 5 |
| 111 | Could serum zonulin be an intestinal permeability marker in diabetes kidney disease?. PLoS ONE, 2021, 16, e0253501. | 1.1 | 5 |
| 112 | Microalbuminuria Is Associated With Early Retinal Neurodegeneration in Patients With Type 2 Diabetes. Ophthalmic Surgery Lasers and Imaging Retina, 2018, 49, e36-e43. | 0.4 | 5 |
| 113 | Prevalence, Awareness, and Treatment of Hypertension in Patients with Type 1 Diabetes: A Nationwide Multicenter Study in Brazil. International Journal of Hypertension, 2013, 2013, 1-8. | 0.5 | 4 |
| 114 | rs1888747 polymorphism in the FRMD3 gene, gene and protein expression: role in diabetic kidney disease. Diabetology and Metabolic Syndrome, 2016, 8, 3. | 1.2 | 4 |
| 115 | The A allele of the UCP2 -866G/A polymorphism changes UCP2 promoter activity in HUVECs treated with high glucose. Molecular Biology Reports, 2019, 46, 4735-4741. | 1.0 | 4 |
| 116 | Progranulin serum levels in human kidney transplant recipients: A longitudinal study. PLoS ONE, 2018, 13, e0192959. | 1.1 | 4 |
| 117 | Body Fat Estimation in Kidney Transplant Recipients: Skinfolds Thickness Compared With Dual-Energy X-Ray Absorptiometry., 2019, 29, 556-562. | | 3 |
| 118 | Aspirin therapy is still underutilized among patients with type 2 diabetes. Arquivos Brasileiros De Endocrinologia E Metabologia, 2006, 50, 1014-1019. | 1.3 | 3 |
| 119 | The A allele of the rs759853 single nucleotide polymorphism in the AKR1B1 gene confers risk for diabetic kidney disease in patients with type 2 diabetes from a Brazilian population. Archives of Endocrinology and Metabolism, 2022, , . | 0.3 | 3 |
| 120 | The rs2304256 Polymorphism in TYK2 Gene Is Associated with Protection for Type 1 Diabetes Mellitus. Diabetes and Metabolism Journal, 2021, 45, 899-908. | 1.8 | 2 |
| 121 | Visceral obesity is associated with higher urinary albumin excretion levels in normoalbuminuric type 2 diabetic patients. Arquivos Brasileiros De Endocrinologia E Metabologia, 2006, 50, 466-471. | 1.3 | 1 |
| 122 | K121Q polymorphism in the Ectonucleotide Pyrophosphatase/Phosphodiesterase 1 gene is associated with acute kidney rejection. PLoS ONE, 2019, 14, e0219062. | 1.1 | 1 |
| 123 | The rs2442598 polymorphism in the ANGPT-2 gene is associated with risk for diabetic retinopathy in patients with type 1 diabetes mellitus in a Brazilian population. Archives of Endocrinology and Metabolism, 2021, 65, . | 0.3 | 1 |
| 124 | Author reply. Ophthalmology, 2015, 122, e43. | 2.5 | 0 |
| 125 | PTPN2 gene polymorphisms are associated with type 1 diabetes mellitus in Brazilian subjects?. Arquivos Brasileiros De Endocrinologia E Metabologia, 2014, 58, 980-981. | 1.3 | 0 |
| 126 | Different Course of the Microvascular Complications of Diabetes Mellitus in Kidney Transplant Recipients with Posttransplant Diabetes—A Longitudinal Study. Diabetes, 2018, 67, 1551-P. | 0.3 | 0 |