

# Guillermina García-Rivera

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3272299/publications.pdf>

Version: 2024-02-01

12  
papers

226  
citations

1163117

8  
h-index

1199594

12  
g-index

13  
all docs

13  
docs citations

13  
times ranked

239  
citing authors

#	ARTICLE	IF	CITATIONS
1	Protein Sumoylation Is Crucial for Phagocytosis in <i>Entamoeba histolytica</i> Trophozoites. <i>International Journal of Molecular Sciences</i> , 2021, 22, 5709.	4.1	3
2	EhVps23: A Component of ESCRT-I That Participates in Vesicular Trafficking and Phagocytosis of <i>Entamoeba histolytica</i> . <i>Frontiers in Cellular and Infection Microbiology</i> , 2021, 11, 770759.	3.9	4
3	A noncanonical GATA transcription factor of <i>Entamoeba histolytica</i> modulates genes involved in phagocytosis. <i>Molecular Microbiology</i> , 2020, 114, 1019-1037.	2.5	2
4	EhRabB mobilises the EhCPADH complex through the actin cytoskeleton during phagocytosis of <i>Entamoeba histolytica</i> . <i>Cellular Microbiology</i> , 2019, 21, e13071.	2.1	10
5	Identification of the gene encoding the TATA box-binding protein-associated factor 1 (TAF1) and its putative role in the heat shock response in the protozoan parasite <i>Entamoeba histolytica</i> . <i>Parasitology Research</i> , 2019, 118, 517-538.	1.6	6
6	The Conserved ESCRT-III Machinery Participates in the Phagocytosis of <i>Entamoeba histolytica</i> . <i>Frontiers in Cellular and Infection Microbiology</i> , 2018, 8, 53.	3.9	40
7	The <i>Entamoeba histolytica</i> TBP and TRF1 transcription factors are GAAC-box binding proteins, which display differential gene expression under different stress stimuli and during the interaction with mammalian cells. <i>Parasites and Vectors</i> , 2018, 11, 153.	2.5	8
8	EhNPC1 and EhNPC2 Proteins Participate in Trafficking of Exogenous Cholesterol in <i>Entamoeba histolytica</i> Trophozoites: Relevance for Phagocytosis. <i>PLoS Pathogens</i> , 2016, 12, e1006089.	4.7	24
9	EhVps32 Is a Vacuole-Associated Protein Involved in Pinocytosis and Phagocytosis of <i>Entamoeba histolytica</i> . <i>PLoS Pathogens</i> , 2015, 11, e1005079.	4.7	32
10	EhADH112 Is a Bro1 Domain-Containing Protein Involved in the <i>Entamoeba histolytica</i> Multivesicular Bodies Pathway. <i>Journal of Biomedicine and Biotechnology</i> , 2012, 2012, 1-15.	3.0	26
11	mRNA Decay Proteins Are Targeted to poly(A) <sup>+</sup> RNA and dsRNA-Containing Cytoplasmic Foci That Resemble P-Bodies in <i>Entamoeba histolytica</i> . <i>PLoS ONE</i> , 2012, 7, e45966.	2.5	17
12	EhCP112 is an <i>Entamoeba histolytica</i> secreted cysteine protease that may be involved in the parasite-virulence. <i>Cellular Microbiology</i> , 2005, 7, 221-232.	2.1	53