

Patricia Lopez

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3278018/patricia-lopez-publications-by-year.pdf>

Version: 2024-04-17

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

88

papers

2,782

citations

29

h-index

50

g-index

95

ext. papers

3,399

ext. citations

4.3

avg, IF

4.97

L-index

#	Paper	IF	Citations
88	Profiling of Serum Oxylipins During the Earliest Stages of Rheumatoid Arthritis. <i>Arthritis and Rheumatology</i> , 2021 , 73, 401-413	9.5	4
87	Low-density granulocytes and monocytes as biomarkers of cardiovascular risk in systemic lupus erythematosus. <i>Rheumatology</i> , 2020 , 59, 1752-1764	3.9	10
86	GlycA Levels during the Earliest Stages of Rheumatoid Arthritis: Potential Use as a Biomarker of Subclinical Cardiovascular Disease. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	5
85	Relationship Between T-Cell Exosomes and Cellular Subsets in SLE According to Type I IFN-Signaling. <i>Frontiers in Medicine</i> , 2020 , 7, 604098	4.9	4
84	IgM anti-phosphorylcholine antibodies associate with senescent and IL-17+ T cells in SLE patients with a pro-inflammatory lipid profile. <i>Rheumatology</i> , 2020 , 59, 407-417	3.9	17
83	Vitamin D Receptor Polymorphism and DHCR7 Contribute to the Abnormal Interplay Between Vitamin D and Lipid Profile in Rheumatoid Arthritis. <i>Scientific Reports</i> , 2019 , 9, 2546	4.9	8
82	Earlier puberty in boys with type 1 diabetes mellitus compared to a simultaneously recruited group of control adolescents. <i>Pediatric Diabetes</i> , 2019 , 20, 197-201	3.6	4
81	Ovarian Function in Adolescents Conceived Using Assisted Reproductive Technologies. <i>Journal of Pediatric and Adolescent Gynecology</i> , 2019 , 32, 117-121	2	0
80	IRF4 and IRGs Delineate Clinically Relevant Gene Expression Signatures in Systemic Lupus Erythematosus and Rheumatoid Arthritis. <i>Frontiers in Immunology</i> , 2018 , 9, 3085	8.4	10
79	Anti-High-Density Lipoprotein Antibodies and Antioxidant Dysfunction in Immune-Driven Diseases. <i>Frontiers in Medicine</i> , 2018 , 5, 114	4.9	6
78	Endothelial Progenitor Cells as Mediators of the Crosstalk between Vascular Repair and Immunity: Lessons from Systemic Autoimmune Diseases. <i>Current Medicinal Chemistry</i> , 2018 , 25, 4478-4496	4.3	7
77	EPC Dysfunction and Immune Networks: Translating Opportunities for Clinical Setting in Personalized Medicine. <i>Current Medicinal Chemistry</i> , 2018 , 25, 4497-4506	4.3	7
76	Profiling of B-Cell Factors and Their Decoy Receptors in Rheumatoid Arthritis: Association With Clinical Features and Treatment Outcomes. <i>Frontiers in Immunology</i> , 2018 , 9, 2351	8.4	9
75	Could Fecal Phenylacetic and Phenylpropionic Acids Be Used as Indicators of Health Status?. <i>Journal of Agricultural and Food Chemistry</i> , 2018 , 66, 10438-10446	5.7	12
74	The role of gut microbiota in lupus: what we know in 2018?. <i>Expert Review of Clinical Immunology</i> , 2018 , 14, 787-792	5.1	7
73	Circulating microparticle subpopulations in systemic lupus erythematosus are affected by disease activity. <i>International Journal of Cardiology</i> , 2017 , 236, 138-144	3.2	19
72	Hyperandrogenism in adolescent girls: relationship with the somatotrophic axis. <i>Journal of Pediatric Endocrinology and Metabolism</i> , 2017 , 30, 561-568	1.6	3

71	High triglycerides and low high-density lipoprotein cholesterol lipid profile in rheumatoid arthritis: A potential link among inflammation, oxidative status, and dysfunctional high-density lipoprotein. <i>Journal of Clinical Lipidology</i> , 2017 , 11, 1043-1054.e2	4.9	26
70	New Diagnostic Criteria of Polycystic Ovarian Morphology for Adolescents: Impact on Prevalence and Hormonal Profile. <i>Hormone Research in Paediatrics</i> , 2017 , 88, 401-407	3.3	15
69	Heterogeneity of the Type I Interferon Signature in Rheumatoid Arthritis: A Potential Limitation for Its Use As a Clinical Biomarker. <i>Frontiers in Immunology</i> , 2017 , 8, 2007	8.4	22
68	Serum Levels of Anti-PON1 and Anti-HDL Antibodies as Potential Biomarkers of Premature Atherosclerosis in Systemic Lupus Erythematosus. <i>Thrombosis and Haemostasis</i> , 2017 , 117, 2194-2206	7	18
67	Intestinal Dysbiosis Is Associated with Altered Short-Chain Fatty Acids and Serum-Free Fatty Acids in Systemic Lupus Erythematosus. <i>Frontiers in Immunology</i> , 2017 , 8, 23	8.4	53
66	Microbiota and oxidant-antioxidant balance in systemic lupus erythematosus. <i>Nutricion Hospitalaria</i> , 2017 , 34, 934-941	1	6
65	Senescent profile of angiogenic T cells from systemic lupus erythematosus patients. <i>Journal of Leukocyte Biology</i> , 2016 , 99, 405-12	6.5	30
64	Phenolic compounds from red wine and coffee are associated with specific intestinal microorganisms in allergic subjects. <i>Food and Function</i> , 2016 , 7, 104-9	6.1	23
63	A pathogenic IFN γ /BlyS and IL-17 axis in Systemic Lupus Erythematosus patients. <i>Scientific Reports</i> , 2016 , 6, 20651	4.9	55
62	Th17 responses and natural IgM antibodies are related to gut microbiota composition in systemic lupus erythematosus patients. <i>Scientific Reports</i> , 2016 , 6, 24072	4.9	123
61	THU0079 HDL, IGG anti-HDL Antibodies and PON1 Axis in Rheumatoid Arthritis Patients: A New Target for CV Disease. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 208.1-208	2.4	
60	FRI0005 A Distinct Profile of Serum Free Fatty Acids Is Associated with Clinical Features and Skewed Th1 Response in Rheumatoid Arthritis Patients. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 427.2-427	2.4	
59	Intestinal dysbiosis in systemic lupus erythematosus: cause or consequence?. <i>Current Opinion in Rheumatology</i> , 2016 , 28, 515-22	5.3	32
58	Paraoxonase 1 Activity Is Modulated by the rs662 Polymorphism and IgG Anti-High-Density Lipoprotein Antibodies in Patients With Rheumatoid Arthritis: Potential Implications for Cardiovascular Disease. <i>Arthritis and Rheumatology</i> , 2016 , 68, 1367-76	9.5	21
57	Comment on: "A new cytofluorimetric approach to evaluate the circulating microparticles in subjects with antiphospholipid antibodies" by Niccolai et al. <i>Thrombosis Research</i> , 2016 , 139, 127	8.2	
56	Allergic Patients with Long-Term Asthma Display Low Levels of Bifidobacterium adolescentis. <i>PLoS ONE</i> , 2016 , 11, e0147809	3.7	62
55	Non-Esterified Fatty Acids Profiling in Rheumatoid Arthritis: Associations with Clinical Features and Th1 Response. <i>PLoS ONE</i> , 2016 , 11, e0159573	3.7	26
54	Pathology-confirmed cerebral arterial invasion and recurrent multiple brain metastasis from cardiac myxoma without evidence of disease after surgery and radiotherapy 2016 , 35, 84-8		3

53	Antibodies to paraoxonase 1 are associated with oxidant status and endothelial activation in rheumatoid arthritis. <i>Clinical Science</i> , 2016 , 130, 1889-99	6.5	12
52	Antibodies to high-density lipoproteins are associated with inflammation and cardiovascular disease in rheumatoid arthritis patients. <i>Translational Research</i> , 2015 , 166, 529-39	11	13
51	Ranking the impact of human health disorders on gut metabolism: systemic lupus erythematosus and obesity as study cases. <i>Scientific Reports</i> , 2015 , 5, 8310	4.9	56
50	Type I IFNs as biomarkers in rheumatoid arthritis: towards disease profiling and personalized medicine. <i>Clinical Science</i> , 2015 , 128, 449-64	6.5	23
49	Altered profile of circulating microparticles in rheumatoid arthritis patients. <i>Clinical Science</i> , 2015 , 128, 437-48	6.5	24
48	Red cell distribution width is associated with endothelial progenitor cell depletion and vascular-related mediators in rheumatoid arthritis. <i>Atherosclerosis</i> , 2015 , 240, 131-6	3.1	25
47	Red wine consumption is associated with fecal microbiota and malondialdehyde in a human population. <i>Journal of the American College of Nutrition</i> , 2015 , 34, 135-41	3.5	24
46	Good response to tumour necrosis factor alpha blockade results in an angiogenic T cell recovery in rheumatoid arthritis patients. <i>Rheumatology</i> , 2015 , 54, 1129-31	3.9	7
45	A single mutation in the gene responsible for the mucoid phenotype of <i>Bifidobacterium animalis</i> subsp. <i>lactis</i> confers surface and functional characteristics. <i>Applied and Environmental Microbiology</i> , 2015 , 81, 7960-8	4.8	33
44	Hirsutism and oligomenorrhea are appropriate screening criteria for polycystic ovary syndrome in adolescents. <i>Gynecological Endocrinology</i> , 2015 , 31, 625-9	2.4	22
43	TNF α polymorphism as marker of immunosenescence for rheumatoid arthritis patients. <i>Experimental Gerontology</i> , 2015 , 61, 123-9	4.5	8
42	Red cell distribution width is associated with cardiovascular risk and disease parameters in rheumatoid arthritis. <i>Rheumatology</i> , 2015 , 54, 641-6	3.9	24
41	SAT0017 Serum Free Fatty Acid Profile in Rheumatoid Arthritis Patients. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 655.2-655	2.4	
40	SAT0116 Antibodies to High Density Lipoprotein are Associated with Altered Blood Lipid Profile, Inflammatory Cytokines and Cardiovascular Disease in Rheumatoid Arthritis Patients. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 692.2-692	2.4	
39	Association of polyphenols from oranges and apples with specific intestinal microorganisms in systemic lupus erythematosus patients. <i>Nutrients</i> , 2015 , 7, 1301-17	6.7	47
38	The effects of <i>Bifidobacterium breve</i> on immune mediators and proteome of HT29 cells monolayers. <i>BioMed Research International</i> , 2015 , 2015, 479140	3	19
37	Angiogenic T cells are decreased in rheumatoid arthritis patients. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 921-7	2.4	23
36	Exopolysaccharide-producing <i>Bifidobacterium animalis</i> subsp. <i>lactis</i> strains and their polymers elicit different responses on immune cells from blood and gut associated lymphoid tissue. <i>Anaerobe</i> , 2014 , 26, 24-30	2.8	47

35	Antimalarial drugs inhibit IFN γ -enhanced TNF α and STAT4 expression in monocytes: implication for systemic lupus erythematosus. <i>Cytokine</i> , 2014 , 67, 13-20	4	8
34	Anti-ribosomal P antibodies are associated with elevated circulating IFN γ and IL-10 levels in systemic lupus erythematosus patients. <i>Lupus</i> , 2014 , 23, 1477-85	2.6	6
33	Interferon- γ -induced B-lymphocyte stimulator expression and mobilization in healthy and systemic lupus erythematosus monocytes. <i>Rheumatology</i> , 2014 , 53, 2249-58	3.9	32
32	A1.75 Angiogenic T cells and derived microparticles disturbances in rheumatoid arthritis patients. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, A33.1-A33	2.4	3
31	FRI0348 Microparticles in Rheumatoid Arthritis Patients: A Principal Component Analysis Approach. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 513.2-513	2.4	
30	Intestinal dysbiosis associated with systemic lupus erythematosus. <i>MBio</i> , 2014 , 5, e01548-14	7.8	309
29	Immune modulating capability of two exopolysaccharide-producing Bifidobacterium strains in a Wistar rat model. <i>BioMed Research International</i> , 2014 , 2014, 106290	3	23
28	Association of levels of antibodies from patients with inflammatory bowel disease with extracellular proteins of food and probiotic bacteria. <i>BioMed Research International</i> , 2014 , 2014, 351204 ³		15
27	IFN γ serum levels are associated with endothelial progenitor cells imbalance and disease features in rheumatoid arthritis patients. <i>PLoS ONE</i> , 2014 , 9, e86069	3.7	26
26	Relationship between FOXP3 positive populations and cytokine production in systemic lupus erythematosus. <i>Cytokine</i> , 2013 , 61, 90-6	4	11
25	Microbial targets for the development of functional foods accordingly with nutritional and immune parameters altered in the elderly. <i>Journal of the American College of Nutrition</i> , 2013 , 32, 399-406	3.5	52
24	Influence of Atg5 mutation in SLE depends on functional IL-10 genotype. <i>PLoS ONE</i> , 2013 , 8, e78756	3.7	34
23	Fatty acids intake and immune parameters in the elderly. <i>Nutricion Hospitalaria</i> , 2013 , 28, 474-8	1	6
22	Immune Modulation Capability of Exopolysaccharides Synthesised by Lactic Acid Bacteria and Bifidobacteria. <i>Probiotics and Antimicrobial Proteins</i> , 2012 , 4, 227-37	5.5	122
21	Treg-inducing membrane vesicles from Bifidobacterium bifidum LMG13195 as potential adjuvants in immunotherapy. <i>Vaccine</i> , 2012 , 30, 825-9	4.1	47
20	Characterisation of the exopolysaccharide (EPS)-producing Lactobacillus paraplantarum BGCG11 and its non-EPS producing derivative strains as potential probiotics. <i>International Journal of Food Microbiology</i> , 2012 , 158, 155-62	5.8	99
19	Exopolysaccharide-producing Bifidobacterium strains elicit different in vitro responses upon interaction with human cells. <i>Food Research International</i> , 2012 , 46, 99-107	7	86
18	Effects of glucocorticoid treatment on CD25 ⁺ FOXP3 ⁺ population and cytokine-producing cells in rheumatoid arthritis. <i>Rheumatology</i> , 2012 , 51, 1198-207	3.9	26

17	Interaction of Bifidobacterium bifidum LMG13195 with HT29 cells influences regulatory-T-cell-associated chemokine receptor expression. <i>Applied and Environmental Microbiology</i> , 2012 , 78, 2850-7	4.8	46
16	Circulating endothelial cells and their progenitors in systemic lupus erythematosus and early rheumatoid arthritis patients. <i>Rheumatology</i> , 2012 , 51, 1775-84	3.9	33
15	Dexamethasone upregulates FOXP3 expression without increasing regulatory activity. <i>Immunobiology</i> , 2011 , 216, 386-92	3.4	42
14	A flagellin-producing Lactococcus strain: interactions with mucin and enteropathogens. <i>FEMS Microbiology Letters</i> , 2011 , 318, 101-7	2.9	19
13	Immune response to Bifidobacterium bifidum strains support Treg/Th17 plasticity. <i>PLoS ONE</i> , 2011 , 6, e24776	3.7	94
12	Glucocorticoids enhance Th17/Th1 imbalance and signal transducer and activator of transcription 3 expression in systemic lupus erythematosus patients. <i>Rheumatology</i> , 2011 , 50, 1794-801	3.9	27
11	IL-10 and TNFalpha genotypes in SLE. <i>Journal of Biomedicine and Biotechnology</i> , 2010 , 2010, 838390		38
10	Distinct Bifidobacterium strains drive different immune responses in vitro. <i>International Journal of Food Microbiology</i> , 2010 , 138, 157-65	5.8	122
9	Conserved anti-proliferative effect and poor inhibition of TNFalpha secretion by regulatory CD4+CD25+ T cells in patients with systemic lupus erythematosus. <i>Clinical Immunology</i> , 2009 , 132, 385-92		9
8	Analysis of Fatty Acid Ethyl Esters in Hair by Headspace Solid-Phase Microextraction (HS-SPME) and Gas Chromatography-Mass Spectrometry (GC-MS). <i>Analytical Letters</i> , 2009 , 42, 2962-2977	2.2	9
7	IFNalpha treatment generates antigen-presenting cells insensitive to atorvastatin inhibition of MHC-II expression. <i>Clinical Immunology</i> , 2008 , 129, 350-9	9	5
6	Influence of functional interleukin 10/tumor necrosis factor-alpha polymorphisms on interferon-alpha, IL-10, and regulatory T cell population in patients with systemic lupus erythematosus receiving antimalarial treatment. <i>Journal of Rheumatology</i> , 2008 , 35, 1559-66	4.1	23
5	Systemic lupus erythematosus in Asturias, Spain: clinical and serologic features. <i>Medicine (United States)</i> , 2006 , 85, 157-168	1.8	33
4	Cytokine polymorphisms influence treatment outcomes in SLE patients treated with antimalarial drugs. <i>Arthritis Research and Therapy</i> , 2006 , 8, R42	5.7	29
3	Enrichment of CD4+ CD25high T cell population in patients with systemic lupus erythematosus treated with glucocorticoids. <i>Annals of the Rheumatic Diseases</i> , 2006 , 65, 1512-7	2.4	115
2	Differential effect of IL10 and TNF{alpha} genotypes on determining susceptibility to discoid and systemic lupus erythematosus. <i>Annals of the Rheumatic Diseases</i> , 2005 , 64, 1605-10	2.4	60
1	Epidemiology of systemic lupus erythematosus in a northern Spanish population: gender and age influence on immunological features. <i>Lupus</i> , 2003 , 12, 860-5	2.6	114