

# Yoshinori Asakawa

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3280072/publications.pdf>

Version: 2024-02-01

345  
papers

9,745  
citations

46918

47  
h-index

102304

66  
g-index

359  
all docs

359  
docs citations

359  
times ranked

4029  
citing authors

#	ARTICLE	IF	CITATIONS
1	Chemosystematics of the Hepaticae. <i>Phytochemistry</i> , 2004, 65, 623-669.	1.4	206
2	Phytochemical and biological studies of bryophytes. <i>Phytochemistry</i> , 2013, 91, 52-80.	1.4	199
3	Biologically active compounds from bryophytes. <i>Pure and Applied Chemistry</i> , 2007, 79, 557-580.	0.9	175
4	Recent advances in phytochemistry of bryophytes-acetogenins, terpenoids and bis(bibenzyl)s from selected Japanese, Taiwanese, New Zealand, Argentinean and European liverworts. <i>Phytochemistry</i> , 2001, 56, 297-312.	1.4	159
5	Chemical Constituents of Bryophytes: Structures and Biological Activity. <i>Journal of Natural Products</i> , 2018, 81, 641-660.	1.5	141
6	Novel neurotrophic isocuparane-type sesquiterpene dimers, mastigophorenes A, B, C and D, isolated from the liverwort <i>Mastigophora diclados</i> . <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1991, , 2737.	0.9	103
7	Cyclic bis(bibenzyls) and related compounds from the liverworts <i>Marchantia polymorpha</i> and <i>Marchantia palmata</i> . <i>Phytochemistry</i> , 1987, 26, 1811-1816.	1.4	91
8	New Bibenzyl Cannabinoid from the New Zealand Liverwort <i>Radula marginata</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 2002, 50, 1390-1392.	0.6	90
9	Prenyl bibenzyls from the liverworts <i>Radula perrottetii</i> and <i>Radula complanata</i> . <i>Phytochemistry</i> , 1991, 30, 235-251.	1.4	89
10	Chemical Constituents of the Ascomycete <i>Daldinia concentrica</i> . <i>Journal of Natural Products</i> , 2002, 65, 1869-1874.	1.5	88
11	Chemical structures of macrocyclic bis(bibenzyls) isolated from liverworts (Hepaticae). <i>Spectroscopy</i> , 2000, 14, 149-175.	0.8	86
12	Introduction. <i>Progress in the Chemistry of Organic Natural Products</i> , 2013, 95, 1-16.	0.8	77
13	Riccardin C, a novel cyclic bibenzyl derivative from <i>Reboulia hemisphaerica</i> . <i>Phytochemistry</i> , 1982, 21, 2143-2144.	1.4	74
14	Inedible mushrooms: a good source of biologically active substances. <i>Chemical Record</i> , 2006, 6, 79-99.	2.9	74
15	Anti-Influenza Activity of Marchantins, Macrocyclic Bisbibenzyls Contained in Liverworts. <i>PLoS ONE</i> , 2011, 6, e19825.	1.1	73
16	Bryophytes: Bio- and Chemical Diversity, Bioactivity and Chemosystematics. <i>Heterocycles</i> , 2009, 77, 99.	0.4	70
17	Phenolic constituents of the liverwort: Four novel cyclic bisbibenzyl dimers from <i>Blasia pusilla</i> L. <i>Tetrahedron</i> , 1996, 52, 14487-14500.	1.0	69
18	Cohaerins A and B, azaphilones from the fungus <i>Hypoxyton cohaerens</i> , and comparison of HPLC-based metabolite profiles in <i>Hypoxyton</i> sect. <i>Annulata</i> . <i>Phytochemistry</i> , 2005, 66, 797-809.	1.4	67

#	ARTICLE	IF	CITATIONS
19	Riccardin C: A natural product that functions as a liver X receptor (LXR) <sup>1</sup> agonist and an LXR <sup>2</sup> antagonist. <i>FEBS Letters</i> , 2005, 579, 5299-5304.	1.3	66
20	A potent cytotoxic warburganal and related drimane-type sesquiterpenoids from <i>Polygonum hydropiper</i> . <i>Phytochemistry</i> , 1980, 21, 2895-2898.	1.4	65
21	Neogrifolin derivatives possessing anti-oxidative activity from the mushroom <i>Albatrellus ovinus</i> . <i>Phytochemistry</i> , 2002, 59, 731-737.	1.4	65
22	Riccardin A and riccardin B, two novel cyclic bis(bibenzylyls) possessing cytotoxicity from the liverwort <i>Riccardia multifida</i> (L.) S. Gray. <i>Journal of Organic Chemistry</i> , 1983, 48, 2164-2167.	1.7	64
23	Biosynthesis of cyclic bis(bibenzylyls) in <i>Marchantia polymorpha</i> . <i>Phytochemistry</i> , 1999, 50, 589-598.	1.4	64
24	Biologically Active Substances of Japanese Inedible Mushrooms. <i>Heterocycles</i> , 1998, 47, 1067.	0.4	62
25	Grifolin derivatives from <i>Albatrellus caeruleoporus</i> , new inhibitors of nitric oxide production in RAW 264.7 cells. <i>Bioorganic and Medicinal Chemistry</i> , 2006, 14, 164-168.	1.4	62
26	Perrottetins E, F, and G from (liverwort)–isolation, structure determination, and synthesis of perrottetin e. <i>Tetrahedron Letters</i> , 1985, 26, 6097-6100.	0.7	58
27	New Chlorinated Cyclic Bis(bibenzylyls) from the Liverworts <i>Herbertus sakuraii</i> and <i>Mastigophora diclados</i> . <i>Tetrahedron</i> , 2000, 56, 3153-3159.	1.0	57
28	In vitro antitrypanosomal activity of plant terpenes against <i>Trypanosoma brucei</i> . <i>Phytochemistry</i> , 2011, 72, 2024-2030.	1.4	57
29	Ent-sesquiterpenoids and cyclic bis(bibenzylyls) from the german liverwort <i>Marchantia Polymorpha</i> . <i>Phytochemistry</i> , 1990, 29, 1577-1584.	1.4	55
30	(+)-Cavicularin: A novel optically active cyclic bibenzyl-dihydrophenanthrene derivative from the liverwort <i>Cavicularia densa</i> Steph. <i>Tetrahedron Letters</i> , 1996, 37, 4745-4748.	0.7	54
31	Changes in secondary metabolism during stomatal ontogeny of <i>Hypoxylon fragiforme</i> . <i>Mycological Research</i> , 2006, 110, 811-820.	2.5	54
32	Sesquiterpenes from <i>Porella species</i> . <i>Phytochemistry</i> , 1978, 17, 457-460.	1.4	53
33	Distribution of novel cyclic bisbibenzyls in <i>Marchantia</i> and <i>Riccardia</i> species. <i>Phytochemistry</i> , 1983, 22, 1413-1415.	1.4	53
34	Structures of a Novel Binaphthyl and Three Novel Benzophenone Denvatives with Plant-Growth Inhibitory Activity from the Fungus <i>Daldinia concentrica</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 1528-1530.	0.6	53
35	(+)-Osmundalactone, $\beta$ -lactones and spiromentins from the fungus <i>Paxillus atrotomentosus</i> . <i>Phytochemistry</i> , 1995, 40, 1251-1257.	1.4	53
36	Terpenoids of the liverwort <i>Frullanoides densifolia</i> and <i>Trocholejeunea sandvicensis</i> . <i>Phytochemistry</i> , 1993, 32, 335-348.	1.4	52

#	ARTICLE	IF	CITATIONS
37	Bryophytes as a source of bioactive volatile terpenoids – A review. Food and Chemical Toxicology, 2019, 132, 110649.	1.8	52
38	Seven new bibenzyls and a dihydrochalcone from <i>Radula variabilis</i> . Phytochemistry, 1978, 17, 2005-2010.	1.4	51
39	Total assignment of <sup>1</sup> H and <sup>13</sup> C NMR spectra of marchantins isolated from liverworts and its application to structure determination of two new macrocyclic bis(bibenzyls) from and. Tetrahedron Letters, 1985, 26, 4735-4738.	0.7	51
40	Cryptoporic acids A-G, drimane-type sesquiterpenoid ethers of isocitric acid from the fungus <i>Cryptoporus volvatus</i> . Phytochemistry, 1992, 31, 579-592.	1.4	51
41	Terpenoids and aromatic compounds from six liverworts. Phytochemistry, 1996, 41, 207-211.	1.4	50
42	Pinguisane and dimeric pinguisane-type sesquiterpenoids from the Japanese liverwort <i>Porella acutifolia</i> subsp. <i>tosana</i> . Phytochemistry, 2000, 53, 593-604.	1.4	50
43	Cyclic azaphilones daldinins E and F from the ascomycete fungus <i>Hypoxylon fuscum</i> (Xylariaceae). Phytochemistry, 2004, 65, 469-473.	1.4	50
44	Novel cytotoxic kaurane-type diterpenoids from the New Zealand Liverwort <i>Jungermannia</i> species. Tetrahedron, 2005, 61, 4531-4544.	1.0	50
45	Inhibition of Nitric Oxide Production in RAW 264.7 Cells by Azaphilones from Xylariaceous Fungi. Biological and Pharmaceutical Bulletin, 2006, 29, 34-37.	0.6	50
46	Chemical Constituents of Bryophytes. Progress in the Chemistry of Organic Natural Products, 2013, , .	0.8	50
47	Sesquiterpenes from Japanese liverworts. Phytochemistry, 1981, 20, 2359-2366.	1.4	49
48	A comparative study on three chemo-types of the liverwort <i>Conocephalum conicum</i> using volatile constituents. Phytochemistry, 1997, 44, 1265-1270.	1.4	48
49	Cytochromes P-450 catalyze the formation of marchantins A and C in <i>Marchantia polymorpha</i> . Phytochemistry, 1999, 52, 1195-1202.	1.4	48
50	Antioxidant Activity of Curtisians I - L from the Inedible Mushroom <i>Paxillus curtisii</i> . Planta Medica, 2003, 69, 1063-1066.	0.7	48
51	Prenyl bibenzyls from the liverwort <i>Radula kojana</i> . Phytochemistry, 1991, 30, 219-234.	1.4	47
52	Sesquiterpenoids from the liverworts <i>Bazzania trilobata</i> and <i>Porella canariensis</i> . Phytochemistry, 1996, 42, 1361-1366.	1.4	47
53	Molecular chemotaxonomy of <i>Daldinia</i> and other Xylariaceae. Mycological Research, 2001, 105, 1191-1205.	2.5	47
54	Cytotoxic and Apoptosis-Inducing <i>ent</i> -Kaurane-Type Diterpenoids from the Japanese Liverwort <i>Jungermannia truncata</i> N <sup>small</sup> EES <sup>small</sup> . Chemical and Pharmaceutical Bulletin, 2002, 50, 808-813.	0.6	47

#	ARTICLE	IF	CITATIONS
55	New Azaphilones from the Inedible Mushroom <i>Hypoxylon rubiginosum</i> . <i>Journal of Natural Products</i> , 2004, 67, 1152-1155.	1.5	47
56	Novel bibenzyl derivatives and ent-cuparene-type sesquiterpenoids from <i>Radula</i> species. <i>Phytochemistry</i> , 1982, 21, 2481-2490.	1.4	46
57	Fusicoccane-, dolabellane- and rearranged labdane-type diterpenoids from the liverwort <i>Pleurozia gigantea</i> . <i>Phytochemistry</i> , 1990, 29, 2597-2603.	1.4	46
58	Structures of Daldinins A-C, Three Novel Azaphilone Derivatives from Ascomycetous Fungus <i>Daldinia concentrica</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 2397-2399.	0.6	46
59	Herbertane-type sesquiterpenoids from the liverwort <i>Herbertus sakuraii</i> . <i>Phytochemistry</i> , 2000, 55, 247-253.	1.4	46
60	Bis(bibenzyls) from Liverworts Inhibit Lipopolysaccharide-Induced Inducible NOS in RAW 264.7 Cells: A Study of Structure-Activity Relationships and Molecular Mechanism. <i>Journal of Natural Products</i> , 2005, 68, 1779-1781.	1.5	46
61	Isoprenyl Phenyl Ethers from Liverworts of the Genus <i>Trichocolea</i> : Cytotoxic Activity, Structural Corrections, and Synthesis. <i>Journal of Natural Products</i> , 1996, 59, 729-733.	1.5	45
62	Isoplagiochins C and D, New Type of Macrocyclic Bis(bibenzyls), Having Two Biphenyl Linkages from the Liverwort <i>Plagiochila fruticosa</i> . <i>Chemistry Letters</i> , 1996, 25, 741-742.	0.7	45
63	Cytochalasins from a <i>Daldinia</i> sp. of fungus. <i>Phytochemistry</i> , 1996, 41, 821-828.	1.4	45
64	Iridoid glucosides from roots of Vietnamese <i>Paederia scandens</i> . <i>Phytochemistry</i> , 2002, 60, 505-514.	1.4	45
65	Sassafrins A-D, new antimicrobial azaphilones from the fungus <i>Creosphaeria sassafras</i> . <i>Tetrahedron</i> , 2005, 61, 1743-1748.	1.0	45
66	Terpenoids and bibenzyls of 25 liverwort <i>Frullania</i> species. <i>Phytochemistry</i> , 1981, 20, 2187-2194.	1.4	42
67	Application of Chromatographic and Spectroscopic Methods towards the Quality Assessment of Ginger ( <i>Zingiber officinale</i> ) Rhizomes from Ecological Plantations. <i>International Journal of Molecular Sciences</i> , 2017, 18, 452.	1.8	42
68	New ent-Kaurene-Type Diterpenoids Possessing Cytotoxicity from the New Zealand Liverwort <i>Jungermannia</i> Species. <i>Chemical and Pharmaceutical Bulletin</i> , 2003, 51, 1189-1192.	0.6	41
69	Sesquiterpenes of six <i>Porella</i> species (hepaticae). <i>Phytochemistry</i> , 1976, 15, 1929-1931.	1.4	40
70	Isotachin A and isotachin B, two sulphur-containing acrylates from the liverwort <i>isotachis japonica</i> . <i>Phytochemistry</i> , 1985, 24, 1505-1508.	1.4	40
71	Cytotoxic, radical scavenging and antimicrobial activities of sesquiterpenoids from the Tahitian liverwort <i>Mastigophora diclados</i> (Brid.) Nees (Mastigophoraceae). <i>Journal of Natural Medicines</i> , 2010, 64, 417-422.	1.1	40
72	Homomono- and sesquiterpenoids from the liverwort <i>Lophocolea heterophylla</i> . <i>Phytochemistry</i> , 1990, 29, 2334-2337.	1.4	39

#	ARTICLE	IF	CITATIONS
73	Zierane sesquiterpene lactone, cembrane and fusicoccane diterpenoids, from the Tahitian liverwort <i>Chandonanthus hirtellus</i> . <i>Phytochemistry</i> , 2010, 71, 1387-1394.	1.4	39
74	Marchantin A Trimethyl Ether: Its Molecular Structure and Tubocurarine-like Skeletal Muscle Relaxation Activity.. <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 52-56.	0.6	38
75	Novel skeletal diterpenoids from the Japanese liverwort <i>Pallavicinia subciliata</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 1998, 46, 178-180.	0.6	38
76	Volatile components of selected species of the liverwort genera <i>Frullania</i> and <i>Schusterella</i> ( <i>Frullaniaceae</i> ) from New Zealand, Australia and South America: a chemosystematic approach. <i>Phytochemistry</i> , 2003, 62, 439-452.	1.4	38
77	Apoptosis-Inducing Properties of ent-Kaurene-Type Diterpenoids from the Liverwort <i>Jungermannia truncata</i> . <i>Planta Medica</i> , 2003, 69, 377-379.	0.7	38
78	Kaurene Diterpene Induces Apoptosis in Human Leukemia Cells Partly through a Caspase-8-Dependent Pathway. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2004, 311, 115-122.	1.3	38
79	Antimicrobial Azaphilones from the Fungus <i>Hypoxyylon multiforme</i> . <i>Planta Medica</i> , 2005, 71, 1058-1062.	0.7	38
80	Plagiochins A, B, C, and D, new type of macrocyclic bis(bibenzyls) having a biphenyl linkage between the ortho positions to the benzyl methylenes, from the liverwort subsp.. <i>Tetrahedron Letters</i> , 1987, 28, 6295-6298.	0.7	37
81	Sesquiterpene and other constituents of the liverwort <i>Dumortiera hirsuta</i> . <i>Phytochemistry</i> , 1997, 44, 293-298.	1.4	37
82	Chemical Constituents of Malagasy Liverworts, Part II: Mastigophoric Acid Methyl Ester of Biogenetic Interest from <i>Mastigophora dicladus</i> ( <i>Lepicoleaceae</i> Subf. <i>Mastigophoroideae</i> ). <i>Chemical and Pharmaceutical Bulletin</i> , 2004, 52, 1382-1384.	0.6	37
83	Antimitotic activity of two macrocyclic bis(bibenzyls), isoplagiochins A and B from the Liverwort <i>Plagiochila fruticosa</i> . <i>Bioorganic and Medicinal Chemistry Letters</i> , 2009, 19, 493-496.	1.0	37
84	Bibenzyl derivatives from the Australian liverwort <i>Frullania falciloba</i> . <i>Phytochemistry</i> , 1987, 26, 1023-1025.	1.4	36
85	Five 10-phenyl-[11]-cytochalasans from a <i>Daldinia</i> fungal species. <i>Phytochemistry</i> , 1995, 40, 135-140.	1.4	36
86	Terpenoids from the Japanese liverworts <i>Jackiella javanica</i> and <i>Jungermannia infusca</i> . <i>Phytochemistry</i> , 1997, 46, 1203-1208.	1.4	36
87	Sesqui- and diterpenoids from two Japanese and three European liverworts. <i>Phytochemistry</i> , 2001, 56, 347-352.	1.4	36
88	Thelephantins A, B and C: three benzoyl p-terphenyl derivatives from the inedible mushroom <i>Thelephora aurantiotincta</i> . <i>Phytochemistry</i> , 2003, 62, 109-113.	1.4	36
89	Thelephantins D-H: five p-terphenyl derivatives from the inedible mushroom <i>Thelephora aurantiotincta</i> . <i>Phytochemistry</i> , 2003, 63, 919-924.	1.4	36
90	Chemical Constituents of Malagasy Liverworts, Part V: Prenyl Bibenzyls and Clerodane Diterpenoids with Nitric Oxide Inhibitory Activity from <i>Radula appressa</i> and <i>Thysananthus spatulistipus</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2006, 54, 1046-1049.	0.6	36

#	ARTICLE	IF	CITATIONS
91	Synthesis of riccardin C and its seven analogues. Part 1: The role of their phenolic hydroxy groups as LXRI± agonists. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2009, 19, 738-741.	1.0	36
92	Fatty acids and cyclic bis(bibenzyls) from the New Zealand liverwort <i>Monoclea forsteri</i> . <i>Phytochemistry</i> , 1988, 27, 2603-2608.	1.4	35
93	Terpenoids and aromatic compounds from selected Ecuadorian liverworts*. <i>Phytochemistry</i> , 1991, 30, 215-217.	1.4	35
94	Cytotoxic 2,3-Secoaromadendrane-Type Sesquiterpenoids from the Liverwort <i>Plagiochila ovalifolia</i> . <i>Planta Medica</i> , 1998, 64, 462-464.	0.7	35
95	Eremophilanolides and other constituents from the Argentine liverwort <i>Frullania brasiliensis</i> . <i>Phytochemistry</i> , 2002, 59, 205-213.	1.4	35
96	Anent-Kaurene Diterpene Enhances Apoptosis Induced by Tumor Necrosis Factor in Human Leukemia Cells. <i>Planta Medica</i> , 2004, 70, 723-727.	0.7	35
97	Efficient synthesis of isoplagiochin D, a macrocyclic bis(bibenzyls), by utilizing an intramolecular Suzuki-Miyaura reaction. <i>Tetrahedron Letters</i> , 2004, 45, 6941-6945.	0.7	35
98	Africane-Type Sesquiterpenoids from the Argentine Liverwort <i>Porella swartziana</i> and Their Antibacterial Activity. <i>Journal of Natural Products</i> , 2004, 67, 31-36.	1.5	35
99	ent-Verticillane-type diterpenoids from the Japanese liverwort <i>Jackiella javanica</i> . <i>Phytochemistry</i> , 2005, 66, 1662-1670.	1.4	35
100	New bibenzyls from <i>Radula complanata</i> . <i>Phytochemistry</i> , 1978, 17, 2115-2117.	1.4	34
101	A Highly efficient preparation of lunularic acid and some biological activities of stilbene and dihydrostilbene derivatives. <i>Phytochemistry</i> , 1988, 27, 109-113.	1.4	34
102	Highlights in phytochemistry of hepaticae-biologically active terpenoids and aromatic compounds. <i>Pure and Applied Chemistry</i> , 1994, 66, 2193-2196.	0.9	34
103	A Novel Class of Plant Type III Polyketide Synthase Involved in Orsellinic Acid Biosynthesis from <i>Rhododendron dauricum</i> . <i>Frontiers in Plant Science</i> , 2016, 7, 1452.	1.7	34
104	Sesquiterpenoids from <i>Chiloscyphus</i> , <i>Clasmatocolea</i> and <i>Frullania</i> species. <i>Phytochemistry</i> , 1983, 22, 961-964.	1.4	33
105	Sesquiterpenoids from the liverwort <i>Porella acutifolia</i> subsp. <i>Tosana</i> . <i>Phytochemistry</i> , 1991, 30, 567-573.	1.4	33
106	Terpenoids from six lophoziaceae liverworts. <i>Phytochemistry</i> , 1993, 34, 181-190.	1.4	33
107	Epi-neoverrucosane- and ent-clerodane-type diterpenoids and ent-2,3-secoaromadendrane- and calamenene-type sesquiterpenoids from the liverwort <i>heteroscyphus planus</i> . <i>Phytochemistry</i> , 1995, 38, 119-127.	1.4	33
108	Sesqui- and Diterpenoids from the Japanese Liverwort <i>Jungermannia infusca</i> . <i>Journal of Natural Products</i> , 2001, 64, 1309-1317.	1.5	33

#	ARTICLE	IF	CITATIONS
109	The Novel Compounds That Activate Farnesoid X Receptor: the Diversity of Their Effects on Gene Expression. <i>Journal of Pharmacological Sciences</i> , 2008, 107, 285-294.	1.1	33
110	Distribution of Bibenzyls, Prenyl Bibenzyls, Bis-bibenzyls, and Terpenoids in the Liverwort Genus <i>Radula</i> . <i>Journal of Natural Products</i> , 2020, 83, 756-769.	1.5	33
111	Concentriols B, C and D, three squalene-type triterpenoids from the ascomycete <i>Daldinia concentrica</i> . <i>Phytochemistry</i> , 2002, 61, 345-353.	1.4	32
112	Composition of the essential oil of the liverwort <i>Radula perrottetii</i> of Japanese origin. <i>Phytochemistry</i> , 2005, 66, 941-949.	1.4	32
113	Terrestrins A-G: p-Terphenyl derivatives from the inedible mushroom <i>Thelephora terrestris</i> . <i>Phytochemistry</i> , 2005, 66, 1052-1059.	1.4	32
114	Distribution of Cyclic and Acyclic Bis-bibenzyls in the Marchantiophyta (Liverworts), Ferns and Higher Plants and Their Biological Activities, Biosynthesis, and Total Synthesis. <i>Heterocycles</i> , 2012, 86, 891.	0.4	32
115	The superoxide release inhibitors, cryptoporin acids C, D, and E; dimeric drimane sesquiterpenoid ethers of isocitric acid from the fungus <i>Cryptoporus volvatus</i> . <i>Journal of the Chemical Society Chemical Communications</i> , 1989, , 258.	2.0	31
116	Cadinane-type sesquiterpenoids from the liverwort <i>Scapania undulata</i> . <i>Phytochemistry</i> , 1994, 37, 1323-1325.	1.4	31
117	Chenopodene, marchantin P and riccardin G from the liverwort <i>Marchantia chenopoda</i> . <i>Phytochemistry</i> , 1994, 36, 73-76.	1.4	31
118	Chapter Five: Distribution of Terpenoids and Aromatic Compounds in Selected Southern Hemispheric Liverworts. <i>Fieldiana Botany</i> , 2008, 47, 37.	0.5	31
119	Structures of four novel macrocyclic bis(bibenzyl) dimers, pusilatins A-D from the liverwort <i>Blasia pusilla</i> . <i>Tetrahedron Letters</i> , 1994, 35, 909-910.	0.7	30
120	Sesqui- and diterpenoids from the Japanese liverwort <i>Jungermannia hattoriana</i> . <i>Phytochemistry</i> , 1997, 45, 353-363.	1.4	30
121	Sesquiterpenoids from the three Japanese liverworts <i>Lejeunea aquatica</i> , <i>L. flava</i> and <i>L. japonica</i> . <i>Phytochemistry</i> , 1997, 46, 145-150.	1.4	30
122	Butenolides from <i>Marchantia paleacea</i> subspecies <i>diptera</i> . <i>Phytochemistry</i> , 1997, 46, 293-296.	1.4	30
123	Sesquiterpenoids, hopanoids and bis(bibenzyls) from the Argentine liverwort <i>Plagiochasma rupestre</i> . <i>Phytochemistry</i> , 1999, 52, 1323-1329.	1.4	30
124	Isolation, Synthesis and Biological Activity of Grifolic Acid Derivatives from the Inedible Mushroom <i>Albatrellus dispansus</i> . <i>Heterocycles</i> , 2005, 65, 2431.	0.4	30
125	Cryptoporin acids A and B, novel bitter drimane sesquiterpenoid ethers of isocitric acid, from the fungus. <i>Tetrahedron Letters</i> , 1987, 28, 6303-6304.	0.7	29
126	First syntheses of 1,13- and 1,15-dihydroxyherbertenes, and herbertenolide by applying intramolecular Heck reaction for the construction of adjacent quaternary centers. <i>Tetrahedron</i> , 2001, 57, 9299-9307.	1.0	29



#	ARTICLE	IF	CITATIONS
127	New Humulane-Type Sesquiterpenes from the Liverworts <i>Tylimanthus tenellus</i> and <i>Marchantia emarginata</i> subsp. <i>tosana</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2004, 52, 481-484.	0.6	29
128	Cinnamacrins A-C, Cinnafagrins D, and Cytostatic Metabolites with $\beta$ -Glucosidase Inhibitory Activity from <i>Cinnamosma macrocarpa</i> . <i>Journal of Natural Products</i> , 2007, 70, 277-282.	1.5	29
129	Marchantiophyta (Liverworts): Rich Sources of Macrocyclic Bis(bibenzyls). <i>Heterocycles</i> , 2008, 76, 99.	0.4	29
130	Cyclopropanochroman derivatives from the liverwort <i>Radula javanica</i> . <i>Phytochemistry</i> , 1991, 30, 325-328.	1.4	28
131	Terpenoids from some German and Russian liverworts. <i>Phytochemistry</i> , 1993, 33, 1445-1448.	1.4	28
132	A 10-phenyl-[11]-cytochalasan from a species of <i>Daldinia</i> . <i>Phytochemistry</i> , 1996, 42, 173-176.	1.4	28
133	Scapaundulins A and B, two novel dimeric labdane diterpenoids, and related compounds from the Japanese liverwort <i>Scapania undulata</i> (L.) Dum.. <i>Tetrahedron Letters</i> , 1997, 38, 1975-1978.	0.7	28
134	Volatile components of the liverworts <i>Archilejeunea olivacea</i> , <i>cheilolejeunea imbricata</i> and <i>Leptolejeunea elliptica</i> . <i>Phytochemistry</i> , 1997, 44, 1261-1264.	1.4	28
135	Ptychantols A-C, macrocyclic bis(bibenzyls), possessing a trans-stilbene structure from the liverwort <i>Ptychanthus striatus</i> . <i>Phytochemistry</i> , 1999, 52, 501-509.	1.4	28
136	Occurrence of Polygodial and 1-(2,4,6-Trimethoxyphenyl)-but-2-en-1-one from Some Ferns and Liverworts: Role of Pungent Components in Bryophytes and Pteridophytes Evolution.. <i>Chemical and Pharmaceutical Bulletin</i> , 2001, 49, 1380-1381.	0.6	28
137	Terpenoids and Aromatic Compounds from the New Zealand Liverworts <i>Plagiochila</i> , <i>Schistochila</i> , and <i>Heteroscyphus</i> Species. <i>Chemical and Pharmaceutical Bulletin</i> , 2004, 52, 556-560.	0.6	28
138	Insect Antifeedant Sesquiterpene Acetals from the Liverwort <i>Lepidolaena clavigera</i> . 2. Structures, Artifacts, and Activity. <i>Journal of Natural Products</i> , 2008, 71, 258-261.	1.5	28
139	Sesqui- and diterpenoids from <i>Plagiochila</i> species. <i>Phytochemistry</i> , 1994, 36, 1425-1430.	1.4	27
140	The absolute structures of new 1.BETA.-hydroxysacculatane-type diterpenoids with piscidal activity from the liverwort <i>Pellia endiviifolia</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 1995, 43, 2030-2032.	0.6	27
141	Terpenoid constituents of the liverwort <i>Heteroscyphus coalitus</i> . <i>Phytochemistry</i> , 1996, 41, 575-580.	1.4	27
142	Identification of cryptic species within liverwort <i>Conocephalum conicum</i> based on the volatile components. <i>Phytochemistry</i> , 2013, 95, 234-241.	1.4	27
143	Dietary intake of metals by the young adult population of Eastern Poland: Results from a market basket study. <i>Journal of Trace Elements in Medicine and Biology</i> , 2016, 35, 36-42.	1.5	27
144	Bibenzyl derivatives from <i>Frullania</i> Species. <i>Phytochemistry</i> , 1987, 26, 1117-1122.	1.4	26

#	ARTICLE	IF	CITATIONS
145	Labdane type diterpenoids from the liverwort <i>Frullania hamachiloba</i> . <i>Phytochemistry</i> , 1988, 27, 1789-1793.	1.4	26
146	Curtisians M-Q: Five Novel p-Terphenyl Derivatives from the Mushroom <i>Paxillus curtisii</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2003, 51, 1064-1067.	0.6	26
147	A Comparison of Apoptosis and Necrosis Induced by ent-Kaurene-Type Diterpenoids in HL-60 Cells. <i>Planta Medica</i> , 2004, 70, 401-406.	0.7	26
148	In vitro antitrypanosomal activity of bis(bibenzyl)s and bibenzyls from liverworts against <i>Trypanosoma brucei</i> . <i>Journal of Natural Medicines</i> , 2012, 66, 377-382.	1.1	26
149	Occurrence of a potent piscicidal diterpenoidal in the liverwort <i>Riccardia lobata</i> var. <i>Yakushimensis</i> . <i>Phytochemistry</i> , 1985, 24, 261-262.	1.4	25
150	Isotachin c and balantiolide two aromatic compounds from the new zealand liverwort <i>balantiopsis rosea</i> . <i>Phytochemistry</i> , 1986, 25, 2543-2546.	1.4	25
151	Novel ent-vibane- and dolabellane-type diterpenoids from the liverwort <i>Odontoschisma denudatum</i> . <i>Tetrahedron Letters</i> , 1998, 39, 579-582.	0.7	25
152	<i>Phytochemistry of Bryophytes.</i> , 1999, , 319-342.		25
153	Occurrence of a Bis-bibenzyl Derivative in the Japanese Fern <i>Hymenophyllum barbatum</i> : First Isolation and Identification of Perrottetin H from the Pteridophytes.. <i>Chemical and Pharmaceutical Bulletin</i> , 1999, 47, 297-298.	0.6	25
154	Identification and Characterization of Daurichromenic Acid Synthase Active in Anti-HIV Biosynthesis. <i>Plant Physiology</i> , 2017, 174, 2213-2230.	2.3	25
155	Bibenzyls from <i>Radula tokiensis</i> and <i>R. Japonica</i> . <i>Phytochemistry</i> , 1981, 20, 858-859.	1.4	24
156	Cuparane- and isocuparane-type sesquiterpenoids in liverworts of the genus <i>Herbertus</i> . <i>Phytochemistry</i> , 1982, 21, 2471-2473.	1.4	24
157	Distribution of cyclic bis(bibenzyls) in the South African liverwort <i>Marchantia polymorpha</i> . <i>Phytochemistry</i> , 1988, 27, 161-163.	1.4	24
158	Mastigophorenes: novel dimeric isocuparane-type sesquiterpenoids from the liverwort <i>Mastigophora diclados</i> . <i>Journal of the Chemical Society Chemical Communications</i> , 1988, , 1341.	2.0	24
159	Two novel macrocyclic bis(Bibenzyls), Isoplagiochins A and B from the liverwort <i>Plagiochila fruticosa</i> . <i>Tetrahedron Letters</i> , 1994, 35, 911-912.	0.7	24
160	Dumortenols, Novel Skeletal Sesquiterpenoids from the Argentinian Liverwort <i>Dumortiera hirsuta</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 1997, 45, 2119-2121.	0.6	24
161	Eudesmane-type sesquiterpene lactones from the Japanese liverwort <i>Frullania densiloba</i> . <i>Phytochemistry</i> , 1997, 45, 555-558.	1.4	24
162	A 7-nordumortenone and other dumortane derivatives from the Argentine liverwort <i>Dumortiera hirsuta</i> . <i>Phytochemistry</i> , 1999, 51, 281-287.	1.4	24

#	ARTICLE	IF	CITATIONS
163	ent-Isopimarane-type diterpenoids from the New Zealand liverwort <i>Trichocolea mollissima</i> . <i>Phytochemistry</i> , 2003, 64, 1319-1325.	1.4	24
164	New Sesquiterpenoids from the New Zealand Liverwort <i>Chiloscyphus subporosus</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2004, 52, 949-952.	0.6	24
165	Synthetic Transformation of Ptychantin into Forskolin and 1,9-Dideoxyforskolin. <i>Journal of Organic Chemistry</i> , 2006, 71, 4619-4624.	1.7	24
166	Heterocyclic Compounds Found in Bryophytes. <i>Heterocycles</i> , 1997, 46, 795.	0.4	24
167	Studies on the liverwort sesquiterpene alcohol tamariscol. Synthesis and absolute configuration. <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1991, , 435.	0.9	23
168	Neurotrophic secoaromadendrane-type sesquiterpenes from the liverwort <i>Plagiochila fruticosa</i> . <i>Phytochemistry</i> , 1991, 30, 4061-4065.	1.4	23
169	Two novel Diels-Alder reaction-type dimeric pinguisane sesquiterpenoids and related compounds from the liverwort <i>Porella acutifolia</i> subsp. <i>tosana</i> . <i>Tetrahedron Letters</i> , 1998, 39, 2977-2980.	0.7	23
170	Triterpenoid constituents of the moss <i>Floribundaria aurea</i> subsp. <i>Nipponica</i> . <i>Phytochemistry</i> , 1998, 48, 297-299.	1.4	23
171	Curtisians Eâ€”H: four p-terphenyl derivatives from the inedible mushroom <i>Paxillus curtisii</i> . <i>Phytochemistry</i> , 2003, 64, 649-654.	1.4	23
172	Thelephantins Iâ€”N p-terphenyl derivatives from the inedible mushroom <i>Hydnellum caeruleum</i> . <i>Phytochemistry</i> , 2004, 65, 1179-1184.	1.4	23
173	Bioactive compounds from liverworts: Inhibition of lipopolysaccharide-induced inducible NOS mRNA in RAW 264.7 cells by Herbertenoids and Cuparenoids. <i>Phytomedicine</i> , 2007, 14, 486-491.	2.3	23
174	An Aromatic Farnesyltransferase Functions in Biosynthesis of the Anti-HIV Meroterpenoid Daurichromenic Acid. <i>Plant Physiology</i> , 2018, 178, 535-551.	2.3	23
175	Sesquiterpene Lactones and Acetogenin Lactones from the Hepaticae and Chemosystematics of the Liverworts <i>Frullania</i> , <i>Plagiochila</i> and <i>Porella</i> . <i>Heterocycles</i> , 2001, 54, 1057.	0.4	23
176	Terpenoids and bibenzyls from some New Zealand liverworts. <i>Phytochemistry</i> , 1982, 21, 2663-2667.	1.4	22
177	Terpenoids from the french liverwort <i>targionia hypophylla</i> . <i>Phytochemistry</i> , 1986, 25, 2555-2556.	1.4	22
178	Sesqui- and diterpenoids from the panamanian liverwort <i>Bryopteris filicina</i> . <i>Phytochemistry</i> , 1994, 37, 433-439.	1.4	22
179	New acorane- and cuparane-type sesqui- and new labdane- and seco-labdane-type diterpenoids from the Japanese liverwort <i>Jungermannia infusca</i> (Mitt.) Steph.. <i>Tetrahedron</i> , 1999, 55, 9117-9132.	1.0	22
180	Dimeric azaphilones from the xylariaceous ascomycete <i>Hypoxylon rutilum</i> . <i>Tetrahedron</i> , 2005, 61, 8451-8455.	1.0	22

#	ARTICLE	IF	CITATIONS
181	Diterpenoids and Aromatic Compounds from the Three New Zealand Liverworts <i>Jamesoniella kirkii</i> , <i>Balantiopsis rosea</i> , and <i>Radula</i> Species. <i>Chemical and Pharmaceutical Bulletin</i> , 2006, 54, 902-906.	0.6	22
182	Fingerprinting of Secondary Metabolites of Liverworts: Chemosystematic Approach. <i>Journal of AOAC INTERNATIONAL</i> , 2014, 97, 1234-1243.	0.7	22
183	Cytotoxic Activity of Riccardin and Perrottetin Derivatives from the Liverwort <i>Lunularia cruciata</i> . <i>Journal of Natural Products</i> , 2019, 82, 694-701.	1.5	22
184	Inhibitory Effect of Cryptoporin Acid E, a Product from Fungus <i>Cryptoporus volvatus</i> , on Colon Carcinogenesis Induced with N-Methyl-N-Nitrosourea in Rats and with 1,2-Dimethylhydrazine in Mice. <i>Japanese Journal of Cancer Research</i> , 1992, 83, 830-834.	1.7	21
185	Structures of five new highly oxygenated labdane-type diterpenoids, ptychantins A-E, closely related to forskolin from the liverwort <i>Ptychanthus striatus</i> . <i>Tetrahedron Letters</i> , 1994, 35, 5457-5460.	0.7	21
186	Total Synthesis and Absolute Configuration of Riccardiphenols A and B, Isolated from the Liverwort <i>Riccardia crassa</i> . <i>Journal of Organic Chemistry</i> , 1996, 61, 5362-5370.	1.7	21
187	Induction of Apoptosis by Newent-Kaurene-Type Diterpenoids Isolated from the New Zealand Liverwort <i>Jungermannia</i> Species. <i>Planta Medica</i> , 2005, 71, 1005-1009.	0.7	21
188	Chemical constituents of the Vietnamese inedible mushroom <i>Xylaria intracolorata</i> . <i>Natural Product Research</i> , 2006, 20, 317-321.	1.0	21
189	Secondary Metabolites of <i>Cinnamosma madagascariensis</i> and Their $\beta$ -Glucosidase Inhibitory Properties. <i>Journal of Natural Products</i> , 2008, 71, 123-126.	1.5	21
190	MAO-A Inhibitory Potential of Terpene Constituents from Ginger Rhizomes: A Bioactivity Guided Fractionation. <i>Molecules</i> , 2018, 23, 1301.	1.7	21
191	Bis-bibenzyls, Bibenzyls, and Terpenoids in 33 Genera of the Marchantiophyta (Liverworts): Structures, Synthesis, and Bioactivity. <i>Journal of Natural Products</i> , 2022, 85, 729-762.	1.5	21
192	Drimane-type sesquiterpenoids from <i>Cryptoporus volvatus</i> infected by <i>Paecilomyces varioti</i> . <i>Phytochemistry</i> , 1993, 33, 1055-1059.	1.4	20
193	Phytol esters and phaeophytins from the hornwort <i>Megaceros flagellaris</i> . <i>Phytochemistry</i> , 1996, 41, 1373-1376.	1.4	20
194	Chemical Conversion of Labdane-Type Diterpenoid Isolated from the Liverwort <i>Porella perrottetiana</i> into (-)-Ambrox. <i>Heterocycles</i> , 1998, 49, 315.	0.4	20
195	Biotransformation of Monoterpenoids. , 2010, , 669-801.		20
196	Vasorelaxant effects of macrocyclic bis(bibenzyls) from liverworts. <i>Bioorganic and Medicinal Chemistry</i> , 2011, 19, 4051-4056.	1.4	20
197	Total synthesis of riccardin C and ( $\pm$ )-cavicularin via Pd-catalyzed Ar-Ar cross couplings. <i>Tetrahedron</i> , 2013, 69, 6959-6968.	1.0	20
198	Clerodane- and halimane-type diterpenoids from the liverwort <i>Jungermannia hyalina</i> . <i>Phytochemistry</i> , 1995, 40, 209-212.	1.4	19

#	ARTICLE	IF	CITATIONS
199	Ent-kaurane-type diterpenoids from the liverwort <i>Jungermannia exsertifolia</i> ssp. <i>cordifolia</i> . <i>Phytochemistry</i> , 1996, 41, 1129-1141.	1.4	19
200	Isolation, Structure Elucidation, and Chemical Derivatization of a New Cyclic Bisbibenzyl Dimer, Pusilatin E, from the Liverwort <i>Riccardia multifida</i> subsp. <i>decrescens</i> . <i>Journal of Natural Products</i> , 1997, 60, 145-147.	1.5	19
201	Diterpenoids from the Japanese liverwort <i>Jungermannia INFUSCA</i> <sup>fn1</sup> <sup>fn1</sup> This paper is dedicated, with best wishes, to Professor G. H. Neil Towers on the occasion of his 75th birthday, in honour of professor G. H. Neil Towers 75TH birthday. <i>Phytochemistry</i> , 1998, 49, 601-608.	1.4	19
202	Occurrence of ent-Sesquiterpene in the Japanese Moss- <i>Plagiomnium acutum</i> : First Isolation and Identification of the ent-Sesqui- and Dolabellane-type Diterpenoids from the Musci.. <i>Chemical and Pharmaceutical Bulletin</i> , 1998, 46, 1488-1489.	0.6	19
203	Highly oxygenated sesquiterpenes from the liverwort <i>Trocholejeunea sandvicensis</i> . <i>Phytochemistry</i> , 2000, 53, 271-276.	1.4	19
204	Chemical Constituents of Malagasy Liverworts, Part III: Sesquiterpenoids from <i>Bazzania decrescens</i> and <i>Bazzania madagassa</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2005, 53, 515-518.	0.6	19
205	Lanostane triterpenoids from the inedible mushroom <i>Fomitopsis spraguei</i> . <i>Phytochemistry</i> , 2005, 66, 1656-1661.	1.4	19
206	Bitter kaurane-type diterpene glucosides from the liverwort <i>Jungermannia infusca</i> . <i>Phytochemistry</i> , 1990, 29, 1619-1623.	1.4	18
207	Stereostructure of Plagiochiline A and Conversion of Plagiochiline A and Stearoylvelutinal into Hot-Tasting Compounds by Human Saliva.. <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 1542-1544.	0.6	18
208	Terpenoids from the liverworts <i>Symphyogyna brasiliensis</i> and unidentified <i>Frullania</i> species. <i>Phytochemistry</i> , 1995, 39, 99-103.	1.4	18
209	Novel sesquiterpenoids from the Colombian liverwort <i>Porella swartziana</i> . <i>Tetrahedron</i> , 1996, 52, 6339-6354.	1.0	18
210	Eudesmane-type sesquiterpenoids from the liverwort <i>Lepidozia vitrea</i> . <i>Phytochemistry</i> , 1996, 41, 833-836.	1.4	18
211	Terpenoid constituents of the New Zealand liverwort <i>Jamesoniella tasmanica</i> . <i>Phytochemistry</i> , 1996, 43, 1057-1064.	1.4	18
212	Sesquiterpene constituents from the liverwort <i>Bazzania japonica</i> . <i>Phytochemistry</i> , 2003, 63, 581-587.	1.4	18
213	Chemotaxonomic value of essential oil components in liverwort species. A review. <i>Flavour and Fragrance Journal</i> , 2015, 30, 189-196.	1.2	18
214	Geographical distribution of tamariscol, a mossy odorous sesquiterpene alcohol, in the liverwort <i>Frullania tamarisci</i> and related species. <i>Phytochemistry</i> , 1991, 30, 2295-2300.	1.4	17
215	Differential effects of cryptoporin acids D and E, inhibitors of superoxide anion radical release, on tumor promotion of okadaic acid in mouse skin. <i>Carcinogenesis</i> , 1991, 12, 1129-1131.	1.3	17
216	Volatile constituent of the liverwort <i>Chiloscyphus pallidus</i> (Mitt.) Engel & Schuster. <i>Flavour and Fragrance Journal</i> , 1994, 9, 237-240.	1.2	17

#	ARTICLE	IF	CITATIONS
217	Eudesmane-Type Sesquiterpenoids from Japanese Liverwort <i>Frullania tamarisci</i> subsp. <i>obscura</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 1998, 46, 542-544.	0.6	17
218	Chemical constituents of Malagasy liverworts: Cyclomyltalanoids from <i>Bazzania madagassa</i> . <i>Phytochemistry</i> , 2006, 67, 2616-2622.	1.4	17
219	Volatile Components from Selected Mexican, Ecuadorian, Greek, German and Japanese Liverworts. <i>Natural Product Communications</i> , 2008, 3, 1934578X0800300.	0.2	17
220	Bryophytes: Liverworts, Mosses, and Hornworts: Extraction and Isolation Procedures. <i>Methods in Molecular Biology</i> , 2013, 1055, 1-20.	0.4	17
221	Trifarienols A and B, Isolated from the liverwort <i>Cheilolejeunea trifaria</i> . Sesquiterpenes having a new carbon skeleton, trifarane. <i>Tetrahedron Letters</i> , 1994, 35, 4787-4788.	0.7	16
222	Sesquiterpenes and a phenolic compound from the liverwort <i>Omphalanthus filiformis</i> . <i>Phytochemistry</i> , 1995, 38, 651-653.	1.4	16
223	Sesqui- and di-terpenoids from the liverwort <i>Jungermannia vulcanicola</i> . <i>Phytochemistry</i> , 1996, 42, 93-96.	1.4	16
224	New Sesqui- and Diterpenoids from the Japanese Liverwort <i>Jungermannia infusca</i> (MITT.) STEPH.. <i>Chemical and Pharmaceutical Bulletin</i> , 1998, 46, 1184-1185.	0.6	16
225	Sesqui- and diterpenoids from <i>Ptilidium ciliare</i> and <i>Barbilophozia</i> species (liverworts). <i>Phytochemistry</i> , 1999, 51, 563-566.	1.4	16
226	A revision of the positive sign of the optical rotation and its maximum value of $\hat{\alpha}_D^{25}$ -eudesmol. <i>Phytochemistry</i> , 1999, 52, 689-694.	1.4	16
227	Tyromycic Acids B <sup>+</sup> E, New Lanostane Triterpenoids from the Mushroom <i>Tyromyces fissilis</i> . <i>Journal of Natural Products</i> , 2004, 67, 148-151.	1.5	16
228	<i>Penicillium sclerotiorum</i> Catalyzes the Conversion of Herbertenediol into Its Dimers: Mastigophorenes A and B. <i>Chemical and Pharmaceutical Bulletin</i> , 2005, 53, 256-257.	0.6	16
229	Chemical constituents of the Vietnamese liverwort <i>Porella densifolia</i> . <i>F<sub>2</sub>-totera p<sub>2</sub></i> , 2010, 81, 659-661.	1.1	16
230	Studies on the Genus <i>Thysananthus</i> (Marchantiophyta, Lejeuneaceae) 3. Terpenoid Chemistry and Chemotaxonomy of Selected Species of <i>Thysananthus</i> and <i>Dendrolejeunea fruticosa</i> . <i>Cryptogamie, Bryologie</i> , 2011, 32, 199-209.	0.1	16
231	Phytochemical investigations and bioactivity evaluation of liverworts as a function of anti-inflammatory and antinociceptive properties in animal models. <i>Pharmaceutical Biology</i> , 2013, 51, 1008-1013.	1.3	16
232	Sesquiterpene derivatives and a norsesquiterpenoid from the liverworts <i>Riccardia crassa</i> and <i>Porella caespitans</i> var. <i>Setigera</i> . <i>Phytochemistry</i> , 1992, 32, 137-140.	1.4	15
233	Rearranged ent-eudesmane- and ent-eremophilane-type sesquiterpenoids from the liverwort <i>Frullania dilatata</i> . <i>Phytochemistry</i> , 1994, 37, 1317-1321.	1.4	15
234	Sesquiterpenoids from the liverwort <i>dicranolejeunea yoshinaga</i> (HATT.) MIZUT.. <i>Chemical and Pharmaceutical Bulletin</i> , 1995, 43, 714-716.	0.6	15

#	ARTICLE	IF	CITATIONS
235	Structure of conocephalenol, a sesquiterpenoid alcohol from the European liverwort <i>Conocephalum conicum</i> : determination of the absolute configuration by total synthesis. <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1995, , 593.	0.9	15
236	Isolation, structure, and synthesis of chenopodanol and the absolute configuration of chenopodene and chenopodanol. <i>Canadian Journal of Chemistry</i> , 1997, 75, 634-640.	0.6	15
237	First synthesis of 1,9-dideoxyforskolin from ptychantin A. <i>Tetrahedron Letters</i> , 2003, 44, 2305-2306.	0.7	15
238	Tyromycic Acids F and G: Two New Triterpenoids from the Mushroom <i>Tyromyces fissilis</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2003, 51, 1441-1443.	0.6	15
239	Constituents of <i>Isotachis aubertii</i> (Isotachidaceae) collected in Madagascar. <i>Biochemical Systematics and Ecology</i> , 2004, 32, 1073-1078.	0.6	15
240	Inhibitory activity of nitric oxide production in RAW 264.7 cells of daldinols A-C from the fungus <i>Daldinia childiae</i> and other metabolites isolated from inedible mushrooms. <i>Journal of Natural Medicines</i> , 2006, 60, 303-307.	1.1	15
241	Insecticidal Constituents from the Argentine Liverwort <i>Plagiochila bursata</i> . <i>Chemistry and Biodiversity</i> , 2010, 7, 1855-1861.	1.0	15
242	Terpenoids and Bibenzyls from Three Argentine Liverworts. <i>Molecules</i> , 2011, 16, 10471-10478.	1.7	15
243	Terpenoids and Aromatic Compounds from Bryophytes and their Central Nervous System Activity. <i>Current Organic Chemistry</i> , 2020, 24, 113-128.	0.9	15
244	Bis(bibenzyl) ethers from <i>Pellia endiviifolia</i> . <i>Phytochemistry</i> , 1991, 30, 1523-1530.	1.4	14
245	Trifarinenols A-E, trifarane-type sesquiterpenoids from the Malaysian liverwort <i>Cheilelejeunea trifaria</i> . <i>Phytochemistry</i> , 1995, 40, 171-176.	1.4	14
246	Sesquiterpenoids from a Cell Suspension Culture of the Liverwort <i>Porella vernicosa</i> Lindb. , 1996, 11, 53-56.		14
247	A phenethyl glycoside from <i>Conocephalum conicum</i> . <i>Phytochemistry</i> , 1996, 43, 1087-1088.	1.4	14
248	A Novel Skeletal Diterpenoid from the German Liverwort <i>Barbilophozia hatcheri</i> (Evans) Loeske.. <i>Chemical and Pharmaceutical Bulletin</i> , 1999, 47, 138-139.	0.6	14
249	New Chroman Derivatives from the Japanese Liverworts <i>Metacalypogeia cordifolia</i> and <i>Cephalozia otaruensis</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 2001, 49, 924-926.	0.6	14
250	Occurrence of a High Concentration of Spider Pheromones in the Ascomycete Fungus <i>Hypoxyylon truncatum</i> . <i>Journal of Natural Products</i> , 2003, 66, 1613-1614.	1.5	14
251	Cytotoxic bibenzyls, and germacrane- and pinguisane-type sesquiterpenoids from Indonesian, Tahitian and Japanese liverworts. <i>Natural Product Communications</i> , 2011, 6, 303-9.	0.2	14
252	Chemosystematics of <i>Porella</i> (Marchantiophyta, Porellaceae). <i>Natural Product Communications</i> , 2011, 6, 315-21.	0.2	14



#	ARTICLE	IF	CITATIONS
253	Fusicorrugatol from the Venezuelan liverwort <i>Plagiochila corrugata</i> . Chemical and Pharmaceutical Bulletin, 1994, 42, 2650-2652.	0.6	13
254	Africane- and monocyclofarnesane-type sesquiterpenoids from the liverwort <i>Porella subobtusa</i> . Phytochemistry, 1996, 43, 1285-1291.	1.4	13
255	Two novel skeletal diterpenoids, neodenudatenones A and B, from the liverwort <i>Odontoschisma denudatum</i> . Tetrahedron Letters, 1998, 39, 3791-3794.	0.7	13
256	Synthesis of cryptoporin acid A methyl ester. Tetrahedron Letters, 2000, 41, 3099-3102.	0.7	13
257	Activation of p38 Mitogen-Activated Protein Kinase during $11\beta$ -Hydroxy-16-kauren-15-one-Induced Apoptosis in Human Leukemia HL-60 Cells. Planta Medica, 2005, 71, 275-277.	0.7	13
258	Chemical Constituents of Malagasy Liverworts. 6. A Myltaylane Caffeeate with Nitric Oxide Inhibitory Activity from <i>Bazzania nitida</i> . Journal of Natural Products, 2007, 70, 856-858.	1.5	13
259	Volatile components from selected Tahitian liverworts. Natural Product Communications, 2009, 4, 1387-92.	0.2	13
260	Distribution of drimane sesquiterpenoids and tocopherols in liverworts, ferns and higher plants: Polygonaceae, Canellaceae and Winteraceae species. Natural Product Communications, 2012, 7, 685-92.	0.2	13
261	Bisbibenzyl and sesquiterpenoids from the liver wort <i>Jubula japonica</i> . Phytochemistry, 1993, 34, 1135-1137.	1.4	12
262	Sesquiterpene esters from the liverwort <i>Plagiochila porelloides</i> . Phytochemistry, 1994, 37, 1091-1093.	1.4	12
263	Synthesis and the absolute configuration of the sesquiterpene aldehyde tridensal from the Taiwanese liverwort <i>Bazzania tridens</i> . Journal of the Chemical Society Perkin Transactions 1, 1995, , 1513.	0.9	12
264	Nudenoic acid: A novel tricyclic sesquiterpenoid from the Taiwanese liverwort <i>Mylia nuda</i> . Tetrahedron Letters, 1996, 37, 9307-9308.	0.7	12
265	Bazzanane Sesquiterpenoids from the New Zealand Liverwort <i>Frullania falciloba</i> . Chemical and Pharmaceutical Bulletin, 2006, 54, 1347-1349.	0.6	12
266	New ent-Verticillane Diterpenoids from the Japanese Liverwort <i>Jackiella javanica</i> . Chemical and Pharmaceutical Bulletin, 2008, 56, 1184-1188.	0.6	12
267	Evaluation of anti-melanoma and tyrosinase inhibitory properties of marchantin A, a natural macrocyclic bisbibenzyl isolated from <i>Marchantia</i> species. Phytochemistry Letters, 2019, 31, 192-195.	0.6	12
268	Setiformenol, isolated from the liverwort <i>tetralophozia setiformis</i> , the first example of cembrane-type diterpene from bryophytes. Tetrahedron Letters, 1993, 34, 643-644.	0.7	11
269	Naphthalene and isocoumarin derivatives from the liverwort <i>Wettsteinia schusterana</i> . Phytochemistry, 1994, 37, 233-235.	1.4	11
270	A New Monocyclic Diterpene from the Liverwort <i>Jungermannia infusca</i> (MITT.) STEPH.. Chemical and Pharmaceutical Bulletin, 1996, 44, 1628-1630.	0.6	11



#	ARTICLE	IF	CITATIONS
271	Ent-kaurane-type diterpenoids from the liverwort <i>Jungermannia rotundata</i> . <i>Phytochemistry</i> , 1997, 44, 653-657.	1.4	11
272	Isolation and Structure of Striatenic Acid from Liverwort <i>Cheilolejeunea serpentina</i> and the Absolute Configuration by Synthesis. <i>Tetrahedron</i> , 2000, 56, 1655-1659.	1.0	11
273	Hydnellins A and B, Nitrogen-Containing Terphenyls from the Mushrooms <i>Hydnellum suaveolens</i> and <i>Hydnellum geogierum</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2006, 54, 912-914.	0.6	11
274	Volatile Components from Selected Tahitian Liverworts. <i>Natural Product Communications</i> , 2009, 4, 1934578X0900401.	0.2	11
275	Volatile Components of the Stressed Liverwort <i>Conocephalum Conicum</i> . <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.2	11
276	Chemotypes and Biomarkers of Seven Species of New Caledonian Liverworts from the Bazzanioideae Subfamily. <i>Molecules</i> , 2018, 23, 1353.	1.7	11
277	Phytochemicals from bryophytes: Structures and biological activity. <i>Journal of the Serbian Chemical Society</i> , 2021, 86, 1139-1175.	0.4	11
278	Sesquiterpenoids from the liverwort <i>Marsupella aquatica</i> . <i>Phytochemistry</i> , 1994, 37, 777-779.	1.4	10
279	Ent-longipinane-type sesquiterpenoid from the liverwort <i>Marsupella emarginata</i> . <i>Phytochemistry</i> , 1994, 37, 1767-1768.	1.4	10
280	Clerodane-type Diterpenoids from the Japanese Liverwort <i>Jungermannia infusca</i> (Mitt.) Steph.. <i>Chemical and Pharmaceutical Bulletin</i> , 2000, 48, 1818-1821.	0.6	10
281	Sesquiterpene hydrocarbons with trifarane backbone in the liverwort <i>Trocholejeunea sandvicensis</i> . <i>Phytochemistry</i> , 2001, 57, 499-506.	1.4	10
282	Expedient Synthetic Transformation of Ptychantins into Forskolin. <i>Synlett</i> , 2008, 2008, 929-931.	1.0	10
283	Chemical Constituents of Selected Japanese and New Zealand Liverworts. <i>Natural Product Communications</i> , 2008, 3, 1934578X0800300.	0.2	10
284	Liverworts-Potential Source of Medicinal Compounds. , 2012, 01, .		10
285	Identification of sesquiterpene lactones in the Bryophyta (mosses) <i>Takakia</i> : <i>Takakia</i> species are closely related chemically to the Marchantiophyta (liverworts). <i>Natural Product Communications</i> , 2015, 10, 5-8.	0.2	10
286	Volatile Components of the Stressed Liverwort <i>Conocephalum conicum</i> . <i>Natural Product Communications</i> , 2016, 11, 103-4.	0.2	10
287	New Highly Oxygenated 6,7-Seco-kaurane- and Bis-kaurane-Type Diterpenoids from The Liverwort <i>Jungermannia exsertifolia</i> STEPH. ssp. <i>cordifolia</i> (DUM.) VANA.. <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 2656-2658.	0.6	9
288	Structures of Four Novel Highly Oxygenated Labdane-Type Diterpenoids, Ptychantins F-I, from the Liverwort <i>Ptychanthus striatus</i> . <i>Chemistry Letters</i> , 1995, 24, 481-482.	0.7	9

#	ARTICLE	IF	CITATIONS
289	seco-Cuparane-type sesquiterpenoid from the Japanese liverwort <i>Jungermannia infusca</i> . <i>Phytochemistry</i> , 2001, 56, 807-810.	1.4	9
290	A Novel Dimeric Lactone Bis-osmundalactone from the Japanese Inedible Mushroom <i>Paxillus atromentosus</i> var. <i>bambusinus</i> . <i>Heterocycles</i> , 2002, 56, 581.	0.4	9
291	Sesqui- and diterpenoids from three New Zealand liverworts: <i>Bazzania novae-zelandiae</i> , <i>Gackstroemia</i> sp. and <i>Dendromastigophora</i> sp.. <i>Natural Product Research</i> , 2010, 24, 68-75.	1.0	9
292	Diversity of Secondary Metabolites in the Liverwort <i>Syzygiella rubricaulis</i> (<sc>Nees</sc>) <sc>Stephani</sc> (Jamesoniellaceae, Marchantiophyta) from Neotropical High Mountains. <i>Chemistry and Biodiversity</i> , 2018, 15, e1800239.	1.0	9
293	Daurichromenic acid-producing oxidocyclase in the young leaves of <i>Rhododendron dauricum</i> . <i>Natural Product Communications</i> , 2014, 9, 1329-32.	0.2	9
294	Malagasy Liverworts, Source of New and Biologically Active Compounds. <i>Natural Product Communications</i> , 2007, 2, 1934578X0700200.	0.2	8
295	Pungent Aromatic Compound from New Zealand Liverwort <i>Hymenophyton flabellatum</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2009, 57, 1015-1018.	0.6	8
296	Cytotoxic Bibenzyls, and Germacrane- and Pinguisane-type Sesquiterpenoids from Indonesian, Tahitian and Japanese Liverworts. <i>Natural Product Communications</i> , 2011, 6, 1934578X1100600.	0.2	8
297	<strong>Chemosystematics of selected liverworts collected in Borneo</strong>. <i>Bryophyte Diversity and Evolution</i> , 2015, 31, 33.	1.0	8
298	Bryophytes: chemical diversity, synthesis and biotechnology. A review.. <i>Flavour and Fragrance Journal</i> , 2011, 26, n/a-n/a.	1.2	7
299	Pungent and Bitter, Cytotoxic and Antiviral Terpenoids from Some Bryophytes and Inedible Fungi. <i>Natural Product Communications</i> , 2014, 9, 1934578X1400900.	0.2	7
300	Phytochemistry of three selected liverworts: <i>Conocepalum conicum</i> , <i>Plagiochila barteri</i> and <i>P. terebrans</i> . <i>Arkivoc</i> , 2007, 2007, 22-29.	0.3	7
301	Recent Advances of Biologically Active Substances from the Marchantiophyta. <i>Natural Product Communications</i> , 2008, 3, 1934578X0800300.	0.2	6
302	Biotransformation of Sesquiterpenoids. , 2010, , 803-892.		6
303	Chemosystematics of <i>Porella</i> (Marchantiophyta, Porellaceae). <i>Natural Product Communications</i> , 2011, 6, 1934578X1100600.	0.2	6
304	Constituents of the Argentine Liverwort <i>Plagiochila diversifolia</i> and Their Insecticidal Activities. <i>Chemistry and Biodiversity</i> , 2017, 14, e1700229.	1.0	6
305	Transcriptome Analysis of Marchantin Biosynthesis from the Liverwort <i>Marchantia polymorpha</i> . <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200.	0.2	6
306	Characteristic Scent from the Tahitian Liverwort, &lt;i>Cyathodium foetidissim&lt;/i>. <i>Journal of Oleo Science</i> , 2018, 67, 1265-1269.	0.6	6

#	ARTICLE	IF	CITATIONS
307	Syntheses of isocitric acid derivatives and biological evaluation. <i>Tetrahedron Letters</i> , 2000, 41, 3095-3098.	0.7	5
308	Pregnane-Type Steroids from the Inedible Mushroom <i>Thelephora terrestris</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2005, 53, 309-312.	0.6	5
309	Volatile Components of Selected Liverworts, and Cytotoxic, Radical Scavenging and Antimicrobial Activities of Their Crude Extracts. <i>Natural Product Communications</i> , 2010, 5, 1934578X1000500.	0.2	5
310	Distribution of Drimane Sesquiterpenoids and Tocopherols in Liverworts, Ferns and Higher Plants: Polygonaceae, Canellaceae and Winteraceae Species. <i>Natural Product Communications</i> , 2012, 7, 1934578X1200700.	0.2	5
311	Chemical variability of the Tahitian <i>Marchantia hexaptera</i> Reich.. <i>Phytochemistry Letters</i> , 2014, 10, xcix-ciii.	0.6	5
312	Biotransformation of Perrottetin F by <i>Aspergillus niger</i> : New Bioactive Secondary Metabolites. <i>Records of Natural Products</i> , 2021, 15, 281-292.	1.3	5
313	Volatile Components Emitted from the Liverwort <i>Marchantia paleacea</i> subsp. <i>diptera</i> . <i>Natural Product Communications</i> , 2016, 11, 263-4.	0.2	5
314	Chemical Relationships between Liverworts of the Family Lejeuneaceae (Porellales,) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 462 Td (Jung</i>	0.2	4
315	Volatile Components Emitted from the Liverwort <i>Marchantia Paleacea</i> Subsp. <i>Diptera</i> . <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.2	4
316	Management of Diabetic Bacterial Foot Infections with Organic Extracts of Liverwort <i>Marchantia debilis</i> from Cameroon. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.2	4
317	Chemotaxonomy and cytotoxicity of the liverwort <i>Porella Viridissima</i> . <i>Natural Product Research</i> , 2021, 35, 2099-2102.	1.0	4
318	Biotransformation of Sesquiterpenoids from Liverworts by Fungi and Mammals. <i>Natural Product Communications</i> , 2010, 5, 1934578X1000500.	0.2	3
319	Chemical Constituents of Marchantiophyta. <i>Progress in the Chemistry of Organic Natural Products</i> , 2013, , 25-561.	0.8	3
320	Influence of Thermal Processing and <i>in vitro</i> Digestion on the Antioxidant Potential of Ginger and Ginger Containing Products. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.2	3
321	Microbial Transformation of Some Natural and Synthetic Aromatic Compounds by Fungi: <i>Aspergillus</i> and <i>Neurospora</i> Strains. <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200.	0.2	3
322	Chemo- and biocatalytic esterification of marchantin A and cytotoxic activity of ester derivatives. <i>FÄ-toterapÄ-ÄÇ</i> , 2020, 142, 104520.	1.1	3
323	Chemical Diversity of Liverworts From <i>Frullania</i> Genus. <i>Natural Product Communications</i> , 2021, 16, 1934578X2199538.	0.2	3
324	Hunting for bis-bibenzylys in <i>Primula veris</i> subsp. <i>macrocalyx</i> (Bunge) <i>LÄ¼di</i> : Organ-specific accumulation and cytotoxic activity. <i>Phytochemistry Letters</i> , 2021, 44, 90-97.	0.6	3

#	ARTICLE	IF	CITATIONS
325	Modification of valencene by bio- and chemical transformation. <i>Natural Product Communications</i> , 2013, 8, 859-62.	0.2	3
326	Chemical Relationships Between Algae, Bryophytes, and Pteridophytes. <i>Progress in the Chemistry of Organic Natural Products</i> , 2013, , 705-726.	0.8	2
327	GC/MS Fingerprinting of Solvent Extracts and Essential Oils Obtained from Liverwort Species. <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200.	0.2	2
328	Biotransformation of sesquiterpenoids from liverworts by fungi and mammals. <i>Natural Product Communications</i> , 2010, 5, 695-707.	0.2	2
329	Influence of Thermal Processing and in vitro Digestion on the Antioxidant Potential of Ginger and Ginger Containing Products. <i>Natural Product Communications</i> , 2016, 11, 1153-1156.	0.2	2
330	Global phytochemistry: research in Japan. <i>Phytochemistry</i> , 2003, 64, 909-912.	1.4	1
331	Chemosystematics of the Hepaticae. <i>ChemInform</i> , 2004, 35, no.	0.1	1
332	Guy Ourisson 1926â€“2006. <i>Phytochemistry</i> , 2007, 68, 1350-1351.	1.4	1
333	Biologically Active Compounds of the Marchantiophyta and Bryophyta. <i>Progress in the Chemistry of Organic Natural Products</i> , 2013, , 619-638.	0.8	1
334	Chemical Diversity of Bryophytes. <i>Progress in the Chemistry of Organic Natural Products</i> , 2013, , 21-24.	0.8	1
335	Chemosystematics of the Thai Liverwort <i>Cheilolejeunea</i> (Marchantiophyta, Lejeuneaceae). <i>Natural Product Communications</i> , 2013, 8, 1934578X1300800.	0.2	1
336	Chemical Evidence for the Liverwort Complex, <i>Chiloscyphus concavus</i> and <i>C. horizontalis</i> . <i>Natural Product Communications</i> , 2014, 9, 1934578X1400900.	0.2	1
337	Bis-bibenzyls from the Cameroon Liverwort <i>Marchantia debilis</i> . <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.2	1
338	Terpenoids, Flavonoids and Acetogenins from Some Malagasy Plants. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.2	1
339	Dimeric and esterified sesquiterpenes from the liverwort <i>Chiastocaulon caledonicum</i> . <i>Phytochemistry</i> , 2020, 179, 112495.	1.4	1
340	Biotransformation of Sesquiterpenoids, Ionones, Damascones, Adamantanes, and Aromatic Compounds by Green Algae, Fungi, and Mammals. , 2009, , 737-841.		1
341	p-Terphenyl Compounds Possessing Antioxidative Activity from Japanese Inedible Mushrooms. <i>International Journal of Medicinal Mushrooms</i> , 2005, 7, 412.	0.9	0
342	p-Terphenyl Compounds Possessing Antioxidative Activity from Japanese Inedible Mushrooms. <i>International Journal of Medicinal Mushrooms</i> , 2005, 7, 410-411.	0.9	0

#	ARTICLE	IF	CITATIONS
343	Chemosystematics of Marchantiophyta. Progress in the Chemistry of Organic Natural Products, 2013, , 639-704.	0.8	0
344	Sesqui- and Diterpenoids from Tahitian and Japanese Liverworts <i>Jungermannia</i> species. Natural Product Communications, 2017, 12, 1934578X1701200.	0.2	0
345	Heterocyclic Stilbene and Bibenzyl Derivatives in Liverworts: Distribution, Structures, Total Synthesis and Biological Activity. Heterocycles, 2022, 105, 179.	0.4	0