

# Chenchen Feng

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3283420/publications.pdf>

Version: 2024-02-01

49  
papers

680  
citations

471509

17  
h-index

642732

23  
g-index

50  
all docs

50  
docs citations

50  
times ranked

1083  
citing authors

#	ARTICLE	IF	CITATIONS
1	Alveolar macrophage-derived exosomal tRF-22-8BWS7K092 activates Hippo signaling pathway to induce ferroptosis in acute lung injury. <i>International Immunopharmacology</i> , 2022, 107, 108690.	3.8	16
2	Collagen synthesis and gap junctions: the highway for metastasis of ovarian cancer. <i>Laboratory Investigation</i> , 2021, 101, 540-542.	3.7	5
3	COLGALT2 is overexpressed in ovarian cancer and interacts with PLOD3. <i>Clinical and Translational Medicine</i> , 2021, 11, e370.	4.0	10
4	BCL2L2 loss renders 14q renal cancer dependent on BCL2L1 that mediates resistance to tyrosine kinase inhibitors. <i>Clinical and Translational Medicine</i> , 2021, 11, e348.	4.0	3
5	PFKFB4 is overexpressed in clear-cell renal cell carcinoma promoting pentose phosphate pathway that mediates Sunitinib resistance. <i>Journal of Experimental and Clinical Cancer Research</i> , 2021, 40, 308.	8.6	23
6	ASXL1 promotes adrenocortical carcinoma and is associated with chemoresistance to EDP regimen. <i>Aging</i> , 2021, 13, 22286-22297.	3.1	2
7	Treatment-related adverse events and response rate to immune checkpoint inhibition. <i>Journal of International Medical Research</i> , 2020, 48, 030006051988645.	1.0	0
8	Association between copy-number alteration of +20q, 14q and 18p and cross-sensitivity to tyrosine kinase inhibitors in clear-cell renal cell carcinoma. <i>Cancer Cell International</i> , 2020, 20, 482.	4.1	3
9	Comprehensive analysis of copy number variance and sensitivity to common targeted therapy in clear cell renal cell carcinoma: In silico analysis with in vitro validation. <i>Cancer Medicine</i> , 2020, 9, 6020-6029.	2.8	8
10	Overexpression of miR-15b Promotes Resistance to Sunitinib in Renal Cell Carcinoma. <i>Journal of Cancer</i> , 2019, 10, 3389-3396.	2.5	26
11	Overexpression of steroid sulfotransferase genes is associated with worsened prognosis and with immune exclusion in clear cell-renal cell carcinoma. <i>Aging</i> , 2019, 11, 9209-9219.	3.1	2
12	Nutlin-3a as a novel anticancer agent for adrenocortical carcinoma with CTNNB1 mutation. <i>Cancer Medicine</i> , 2018, 7, 1440-1449.	2.8	14
13	Knockdown of tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein zeta (YWHAZ) enhances tumorigenesis both in vivo and in vitro in bladder cancer. <i>Oncology Reports</i> , 2018, 39, 2127-2135.	2.6	11
14	Overexpression of low density lipoprotein receptor-related protein 1 (LRP1) is associated with worsened prognosis and decreased cancer immunity in clear-cell renal cell carcinoma. <i>Biochemical and Biophysical Research Communications</i> , 2018, 503, 1537-1543.	2.1	25
15	Body-mass index and metastatic melanoma outcomes. <i>Lancet Oncology</i> , The, 2018, 19, e223.	10.7	3
16	&beta;-Catenin-driven adrenocortical carcinoma is characterized with immune exclusion. <i>OncoTargets and Therapy</i> , 2018, Volume 11, 2029-2036.	2.0	24
17	Association Study of Reported Significant Loci at 5q35.3, 7p14.3, 13q14.1 and 16p12.3 with Urolithiasis in Chinese Han Ethnicity. <i>Scientific Reports</i> , 2017, 7, 45766.	3.3	10
18	Tumor Wide Horizontal Invasion Predicts Local Recurrence for Scrotal Extramammary Paget's Disease. <i>Scientific Reports</i> , 2017, 7, 44933.	3.3	7

#	ARTICLE	IF	CITATIONS
19	Epigallocatechin gallate inhibits the growth and promotes the apoptosis of bladder cancer cells. <i>Experimental and Therapeutic Medicine</i> , 2017, 14, 3513-3518.	1.8	15
20	Overexpression of HSD17B4 exerts tumor suppressive function in adrenocortical carcinoma and is not associated with hormone excess. <i>Oncotarget</i> , 2017, 8, 114736-114745.	1.8	7
21	Lipocalin 2 over-expression facilitates progress of castration-resistant prostate cancer via improving androgen receptor transcriptional activity. <i>Oncotarget</i> , 2016, 7, 64309-64317.	1.8	20
22	Prognostic value of Ki67 and p63 expressions in bladder cancer patients who underwent radical cystectomy. <i>International Urology and Nephrology</i> , 2016, 48, 495-501.	1.4	19
23	Genetic alteration in notch pathway is associated with better prognosis in renal cell carcinoma. <i>BioFactors</i> , 2016, 42, 41-48.	5.4	16
24	Coexistence of YWHAZ amplification predicts better prognosis in muscle-invasive bladder cancer with CDKN2A or TP53 loss. <i>Oncotarget</i> , 2016, 7, 34752-34758.	1.8	6
25	TFPI-2 expression is decreased in bladder cancer and is related to apoptosis. <i>Journal of B U on</i> , 2016, 21, 1518-1523.	0.4	10
26	PI3K $\hat{\imath}^2$ Inhibitor TGX221 Selectively Inhibits Renal Cell Carcinoma Cells with Both VHL and SETD2 mutations and Links Multiple Pathways. <i>Scientific Reports</i> , 2015, 5, 9465.	3.3	42
27	Loss of MLH1 confers resistance to PI3K $\hat{\imath}^2$ inhibitors in renal clear cell carcinoma with SETD2 mutation. <i>Tumor Biology</i> , 2015, 36, 3457-3464.	1.8	19
28	PD33-07 PI3K $\hat{\imath}^2$ INHIBITOR TGX221 SELECTIVELY INHIBITS RENAL CELL CARCINOMA WITH COMMON MUTATIONS AND LINKS MULTIPLE PATHWAYS. <i>Journal of Urology</i> , 2015, 193, .	0.4	0
29	MP39-03 LOSS OF MLH1 CONFERS RESISTANCE TO PI3K $\hat{\imath}^2$ INHIBITORS IN RENAL CLEAR CELL CARCINOMA WITH SETD2 MUTATION. <i>Journal of Urology</i> , 2015, 193, .	0.4	0
30	BLCA1 expression is associated with angiogenesis of bladder cancer and is correlated with common pro-angiogenic factors. <i>International Journal of Clinical and Experimental Medicine</i> , 2015, 8, 16259-65.	1.3	5
31	Urinary BLCA1 is specific for urothelial cancer detection in Chinese ethnicity. <i>International Journal of Clinical and Experimental Medicine</i> , 2015, 8, 18819-24.	1.3	0
32	RNA Sequencing Reveals Upregulation of RUNX1-RUNX1T1 Gene Signatures in Clear Cell Renal Cell Carcinoma. <i>BioMed Research International</i> , 2014, 2014, 1-8.	1.9	15
33	Ki67 and TP53 expressions predict recurrence of non-muscle-invasive bladder cancer. <i>Tumor Biology</i> , 2014, 35, 2989-2995.	1.8	36
34	Predictive Value of Clinicopathological Markers for the Metachronous Bladder Cancer and Prognosis of Upper Tract Urothelial Carcinoma. <i>Scientific Reports</i> , 2014, 4, 4015.	3.3	18
35	Ureteroscopic Holmium:YAG laser lithotripsy is effective for ureteral steinstrasse post-SWL. <i>Minimally Invasive Therapy and Allied Technologies</i> , 2013, 22, 279-282.	1.2	5
36	Paraneoplastic symptoms: Cachexia, polycythemia, and hypercalcemia are, respectively, related to vascular endothelial growth factor (VEGF) expression in renal clear cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013, 31, 1820-1825.	1.6	20

#	ARTICLE	IF	CITATIONS
37	Expression of pigment epithelium-derived factor and tumor necrosis factor- $\alpha$ is correlated in bladder tumor and is related to tumor angiogenesis. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013, 31, 241-246.	1.6	39
38	Med19 promotes bone metastasis and invasiveness of bladder urothelial carcinoma via bone morphogenetic protein 2. <i>Annals of Diagnostic Pathology</i> , 2013, 17, 259-264.	1.3	36
39	Adrenal myelolipoma: A mingle of progenitor cells?. <i>Medical Hypotheses</i> , 2013, 80, 819-822.	1.5	27
40	Endoureterotomy is not a sufficient treatment for intrinsic ureteral endometriosis. <i>Wideochirurgia I Inne Techniki Maloinwazyjne</i> , 2013, 3, 187-191.	0.7	4
41	Combination of Rapamycin, CI-1040, and 17-AAG Inhibits Metastatic Capacity of Prostate Cancer via Slug Inhibition. <i>PLoS ONE</i> , 2013, 8, e77400.	2.5	26
42	Mesenchymal stem cells in prostate cancer have higher expressions of SDF-1, CXCR4 and VEGF. <i>General Physiology and Biophysics</i> , 2013, 32, 245-250.	0.9	7
43	Relationship of TP53 and Ki67 expression in bladder cancer under WHO 2004 classification. <i>Journal of BU on</i> , 2013, 18, 420-4.	0.4	9
44	The Association Between Polymorphisms in Prooxidant or Antioxidant Enzymes (Myeloperoxidase,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 Clinical Genitourinary Cancer, 2012, 10, 251-255.	1.9	18
45	Use of NTrap during ureteroscopic Holmium:YAG laser lithotripsy of upper ureteral calculi. <i>Minimally Invasive Therapy and Allied Technologies</i> , 2012, 21, 78-82.	1.2	10
46	BLCA-4 expression is related to MMP-9, VEGF, IL-1 $\alpha$ and IL-8 in bladder cancer but not to PEDF, TNF- $\alpha$ or angiogenesis. <i>Pathologie Et Biologie</i> , 2012, 60, e36-e40.	2.2	17
47	Ureteroscopic holmium:YAG laser endopyelotomy is effective in distinctive ureteropelvic junction obstructions. <i>Wideochirurgia I Inne Techniki Maloinwazyjne</i> , 2011, 3, 144-149.	0.7	8
48	Pigment epithelium-derived factor expression is down-regulated in bladder tumors and correlates with vascular endothelial growth factor and matrix metalloproteinase-9. <i>International Urology and Nephrology</i> , 2011, 43, 383-390.	1.4	20
49	Expression of pigment epithelium-derived factor in bladder tumour is correlated with interleukin-8 yet not with interleukin-1 $\alpha$ . <i>Journal of Huazhong University of Science and Technology [Medical Sciences]</i> , 2011, 31, 21-25.	1.0	13