

Kate Hallsworth

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3286421/publications.pdf>

Version: 2024-02-01

28
papers

1,917
citations

394286

19
h-index

526166

27
g-index

29
all docs

29
docs citations

29
times ranked

2773
citing authors

#	ARTICLE	IF	CITATIONS
1	Resistance exercise reduces liver fat and its mediators in non-alcoholic fatty liver disease independent of weight loss. <i>Gut</i> , 2011, 60, 1278-1283.	6.1	382
2	Modified high-intensity interval training reduces liver fat and improves cardiac function in non-alcoholic fatty liver disease: a randomized controlled trial. <i>Clinical Science</i> , 2015, 129, 1097-1105.	1.8	165
3	Exercise Reduces Liver Lipids and Visceral Adiposity in Patients With Nonalcoholic Steatohepatitis in a Randomized Controlled Trial. <i>Clinical Gastroenterology and Hepatology</i> , 2017, 15, 96-102.e3.	2.4	163
4	Lifestyle modification in NAFLD/NASH: Facts and figures. <i>JHEP Reports</i> , 2019, 1, 468-479.	2.6	147
5	High intensity intermittent exercise improves cardiac structure and function and reduces liver fat in patients with type 2 diabetes: a randomised controlled trial. <i>Diabetologia</i> , 2016, 59, 56-66.	2.9	141
6	Cardiac structure and function are altered in adults with non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , 2013, 58, 757-762.	1.8	122
7	Non-alcoholic fatty liver disease: A patient guideline. <i>JHEP Reports</i> , 2021, 3, 100322.	2.6	109
8	Effects of Community Exercise Therapy on Metabolic, Brain, Physical, and Cognitive Function Following Stroke. <i>Neurorehabilitation and Neural Repair</i> , 2015, 29, 623-635.	1.4	102
9	Non-alcoholic fatty liver disease is associated with higher levels of objectively measured sedentary behaviour and lower levels of physical activity than matched healthy controls. <i>Frontline Gastroenterology</i> , 2015, 6, 44-51.	0.9	91
10	Health-related Quality of Life in Nonalcoholic Fatty Liver Disease Associates With Hepatic Inflammation. <i>Clinical Gastroenterology and Hepatology</i> , 2019, 17, 2085-2092.e1.	2.4	79
11	Effect of Left Ventricular Assist Device Implantation and Heart Transplantation on Habitual Physical Activity and Quality of Life. <i>American Journal of Cardiology</i> , 2014, 114, 88-93.	0.7	65
12	The degree of hepatic steatosis associates with impaired cardiac and autonomic function. <i>Journal of Hepatology</i> , 2019, 70, 1203-1213.	1.8	45
13	Barriers and Facilitators to Mediterranean Diet Adoption by Patients With Nonalcoholic Fatty Liver Disease in Northern Europe. <i>Clinical Gastroenterology and Hepatology</i> , 2019, 17, 1364-1371.e3.	2.4	42
14	Lifestyle Behavior Change in Patients With Nonalcoholic Fatty Liver Disease: A Qualitative Study of Clinical Practice. <i>Clinical Gastroenterology and Hepatology</i> , 2017, 15, 1968-1971.	2.4	37
15	Targeting Lifestyle Behavior Change in Adults with NAFLD During a 20-min Consultation: Summary of the Dietary and Exercise Literature. <i>Current Gastroenterology Reports</i> , 2016, 18, 11.	1.1	34
16	Using the theoretical domains framework to identify barriers and enabling factors to implementation of guidance for the diagnosis and management of nonalcoholic fatty liver disease: a qualitative study. <i>Translational Behavioral Medicine</i> , 2020, 10, 1016-1030.	1.2	34
17	Feasibility of a Very Low Calorie Diet to Achieve a Sustainable 10% Weight Loss in Patients With Nonalcoholic Fatty Liver Disease. <i>Clinical and Translational Gastroenterology</i> , 2020, 11, e00231.	1.3	28
18	Unsupervised high-intensity interval training improves glycaemic control but not cardiovascular autonomic function in type 2 diabetes patients: A randomised controlled trial. <i>Diabetes and Vascular Disease Research</i> , 2019, 16, 69-76.	0.9	26

#	ARTICLE	IF	CITATIONS
19	Real-world management of non-alcoholic steatohepatitis differs from clinical practice guideline recommendations and across regions. <i>JHEP Reports</i> , 2022, 4, 100411.	2.6	24
20	A study of physical activity comparing people with Charcot-Marie-Tooth disease to normal control subjects. <i>Disability and Rehabilitation</i> , 2017, 39, 1753-1758.	0.9	19
21	Implementation of a care bundle improves the management of patients with non-alcoholic fatty liver disease. <i>Frontline Gastroenterology</i> , 2021, 12, 578-585.	0.9	17
22	Digital Intervention With Lifestyle Coach Support to Target Dietary and Physical Activity Behaviors of Adults With Nonalcoholic Fatty Liver Disease: Systematic Development Process of VITALISE Using Intervention Mapping. <i>Journal of Medical Internet Research</i> , 2021, 23, e20491.	2.1	12
23	Effects of Exercise on Liver Fat and Metabolism in Alcohol Drinkers. <i>Clinical Gastroenterology and Hepatology</i> , 2017, 15, 1596-1603.e3.	2.4	9
24	Assessing the feasibility and acceptability of Changing Health for the management of prediabetes: protocol for a pilot study of a digital behavioural intervention. <i>Pilot and Feasibility Studies</i> , 2019, 5, 139.	0.5	8
25	Exercise therapy in primary biliary cirrhosis: the importance of moving while sitting on a surgical waiting list—a case study: Table 1. <i>Frontline Gastroenterology</i> , 2016, 7, 167-169.	0.9	7
26	Factors associated with engagement and adherence to a low-energy diet to promote 10% weight loss in patients with clinically significant non-alcoholic fatty liver disease. <i>BMJ Open Gastroenterology</i> , 2021, 8, e000678.	1.1	6
27	Adiposity predicts low cardiorespiratory fitness in individuals with metabolic diseases. <i>Diabetes Research and Clinical Practice</i> , 2018, 146, 300-304.	1.1	3
28	Physical Activity, Inactivity and Sleep in Patients with Significant Non-Alcoholic Fatty Liver Disease. <i>American Journal of the Medical Sciences</i> , 2022, 363, 80-83.	0.4	0