

Kate Hallsworth

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3286421/publications.pdf>

Version: 2024-02-01

28
papers

1,917
citations

393982

19
h-index

525886

27
g-index

29
all docs

29
docs citations

29
times ranked

2773
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Physical Activity, Inactivity and Sleep in Patients with Significant Non-Alcoholic Fatty Liver Disease. American Journal of the Medical Sciences, 2022, 363, 80-83. | 0.4 | 0 |
| 2 | Real-world management of non-alcoholic steatohepatitis differs from clinical practice guideline recommendations and across regions. JHEP Reports, 2022, 4, 100411. | 2.6 | 24 |
| 3 | Implementation of a care bundle improves the management of patients with non-alcoholic fatty liver disease. Frontline Gastroenterology, 2021, 12, 578-585. | 0.9 | 17 |
| 4 | Factors associated with engagement and adherence to a low-energy diet to promote 10% weight loss in patients with clinically significant non-alcoholic fatty liver disease. BMJ Open Gastroenterology, 2021, 8, e000678. | 1.1 | 6 |
| 5 | Non-alcoholic fatty liver disease: A patient guideline. JHEP Reports, 2021, 3, 100322. | 2.6 | 109 |
| 6 | Digital Intervention With Lifestyle Coach Support to Target Dietary and Physical Activity Behaviors of Adults With Nonalcoholic Fatty Liver Disease: Systematic Development Process of VITALISE Using Intervention Mapping. Journal of Medical Internet Research, 2021, 23, e20491. | 2.1 | 12 |
| 7 | Using the theoretical domains framework to identify barriers and enabling factors to implementation of guidance for the diagnosis and management of nonalcoholic fatty liver disease: a qualitative study. Translational Behavioral Medicine, 2020, 10, 1016-1030. | 1.2 | 34 |
| 8 | Feasibility of a Very Low Calorie Diet to Achieve a Sustainable 10% Weight Loss in Patients With Nonalcoholic Fatty Liver Disease. Clinical and Translational Gastroenterology, 2020, 11, e00231. | 1.3 | 28 |
| 9 | Lifestyle modification in NAFLD/NASH: Facts and figures. JHEP Reports, 2019, 1, 468-479. | 2.6 | 147 |
| 10 | Health-related Quality of Life in Nonalcoholic Fatty Liver Disease Associates With Hepatic Inflammation. Clinical Gastroenterology and Hepatology, 2019, 17, 2085-2092.e1. | 2.4 | 79 |
| 11 | The degree of hepatic steatosis associates with impaired cardiac and autonomic function. Journal of Hepatology, 2019, 70, 1203-1213. | 1.8 | 45 |
| 12 | Barriers and Facilitators to Mediterranean Diet Adoption by Patients With Nonalcoholic Fatty Liver Disease in Northern Europe. Clinical Gastroenterology and Hepatology, 2019, 17, 1364-1371.e3. | 2.4 | 42 |
| 13 | Assessing the feasibility and acceptability of Changing Health for the management of prediabetes: protocol for a pilot study of a digital behavioural intervention. Pilot and Feasibility Studies, 2019, 5, 139. | 0.5 | 8 |
| 14 | Unsupervised high-intensity interval training improves glycaemic control but not cardiovascular autonomic function in type 2 diabetes patients: A randomised controlled trial. Diabetes and Vascular Disease Research, 2019, 16, 69-76. | 0.9 | 26 |
| 15 | Adiposity predicts low cardiorespiratory fitness in individuals with metabolic diseases. Diabetes Research and Clinical Practice, 2018, 146, 300-304. | 1.1 | 3 |
| 16 | Effects of Exercise on Liver Fat and Metabolism in Alcohol Drinkers. Clinical Gastroenterology and Hepatology, 2017, 15, 1596-1603.e3. | 2.4 | 9 |
| 17 | Lifestyle Behavior Change in Patients With Nonalcoholic Fatty Liver Disease: A Qualitative Study of Clinical Practice. Clinical Gastroenterology and Hepatology, 2017, 15, 1968-1971. | 2.4 | 37 |
| 18 | A study of physical activity comparing people with Charcot-Marie-Tooth disease to normal control subjects. Disability and Rehabilitation, 2017, 39, 1753-1758. | 0.9 | 19 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Exercise Reduces Liver Lipids and Visceral Adiposity in Patients With Nonalcoholic Steatohepatitis in a Randomized Controlled Trial. <i>Clinical Gastroenterology and Hepatology</i> , 2017, 15, 96-102.e3. | 2.4 | 163 |
| 20 | Exercise therapy in primary biliary cirrhosis: the importance of moving while sitting on a surgical waiting list—a case study: Table 1. <i>Frontline Gastroenterology</i> , 2016, 7, 167-169. | 0.9 | 7 |
| 21 | Targeting Lifestyle Behavior Change in Adults with NAFLD During a 20-min Consultation: Summary of the Dietary and Exercise Literature. <i>Current Gastroenterology Reports</i> , 2016, 18, 11. | 1.1 | 34 |
| 22 | High intensity intermittent exercise improves cardiac structure and function and reduces liver fat in patients with type 2 diabetes: a randomised controlled trial. <i>Diabetologia</i> , 2016, 59, 56-66. | 2.9 | 141 |
| 23 | Modified high-intensity interval training reduces liver fat and improves cardiac function in non-alcoholic fatty liver disease: a randomized controlled trial. <i>Clinical Science</i> , 2015, 129, 1097-1105. | 1.8 | 165 |
| 24 | Non-alcoholic fatty liver disease is associated with higher levels of objectively measured sedentary behaviour and lower levels of physical activity than matched healthy controls. <i>Frontline Gastroenterology</i> , 2015, 6, 44-51. | 0.9 | 91 |
| 25 | Effects of Community Exercise Therapy on Metabolic, Brain, Physical, and Cognitive Function Following Stroke. <i>Neurorehabilitation and Neural Repair</i> , 2015, 29, 623-635. | 1.4 | 102 |
| 26 | Effect of Left Ventricular Assist Device Implantation and Heart Transplantation on Habitual Physical Activity and Quality of Life. <i>American Journal of Cardiology</i> , 2014, 114, 88-93. | 0.7 | 65 |
| 27 | Cardiac structure and function are altered in adults with non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , 2013, 58, 757-762. | 1.8 | 122 |
| 28 | Resistance exercise reduces liver fat and its mediators in non-alcoholic fatty liver disease independent of weight loss. <i>Gut</i> , 2011, 60, 1278-1283. | 6.1 | 382 |