Jung Yoon Pyo

List of Publications by Citations

Source: https://exaly.com/author-pdf/3289299/jung-yoon-pyo-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

160 26 1,114 17 h-index g-index citations papers 176 1,435 3.5 4.71 L-index avg, IF ext. citations ext. papers

#	Paper	IF	Citations
160	Measurement of the serum leptin level could assist disease activity monitoring in rheumatoid arthritis. <i>Rheumatology International</i> , 2007 , 27, 537-40	3.6	77
159	Differences in clinical manifestations and outcomes between adult and child patients with Henoch-Schilein purpura. <i>Journal of Korean Medical Science</i> , 2014 , 29, 198-203	4.7	76
158	Application of the 2016 EULAR/ACR/PRINTO Classification Criteria for Macrophage Activation Syndrome in Patients with Adult-onset Still Disease. <i>Journal of Rheumatology</i> , 2017 , 44, 996-1003	4.1	31
157	Usefulness of serum leucine-rich alpha-2 glycoprotein as a disease activity biomarker in patients with rheumatoid arthritis. <i>Journal of Korean Medical Science</i> , 2014 , 29, 1199-204	4.7	28
156	RORE-specific transcriptional interactomic inhibition suppresses autoimmunity associated with TH17 cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014 , 111, 18673-8	11.5	28
155	Malignancies in Korean patients with inflammatory myopathy. <i>Yonsei Medical Journal</i> , 2006 , 47, 519-23	3	28
154	Systemic immune-inflammation index could estimate the cross-sectional high activity and the poor outcomes in immunosuppressive drug-nalle patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Nephrology</i> , 2019 , 24, 711-717	2.2	27
153	Alleviation of rheumatoid arthritis by cell-transducible methotrexate upon transcutaneous delivery. <i>Biomaterials</i> , 2012 , 33, 1563-72	15.6	26
152	C-Reactive Protein to Serum Albumin Ratio Is an Independent Predictor of All-Cause Mortality in Patients with ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2018 , 59, 865-871	3	25
151	Evaluation of Spleen Glucose Metabolism Using F-FDG PET/CT in Patients with Febrile Autoimmune Disease. <i>Journal of Nuclear Medicine</i> , 2017 , 58, 507-513	8.9	24
150	Malignancies in Korean patients with immunoglobulin G4-related disease. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 1028-1035	2.3	24
149	In-hospital mortality in febrile lupus patients based on 2016 EULAR/ACR/PRINTO classification criteria for macrophage activation syndrome. <i>Seminars in Arthritis and Rheumatism</i> , 2017 , 47, 216-221	5.3	21
148	Neutrophil to lymphocyte ratio at diagnosis can estimate vasculitis activity and poor prognosis in patients with ANCA-associated vasculitis: a retrospective study. <i>BMC Nephrology</i> , 2018 , 19, 187	2.7	21
147	Platelet to lymphocyte ratio is associated with the current activity of ANCA-associated vasculitis at diagnosis: a retrospective monocentric study. <i>Rheumatology International</i> , 2018 , 38, 1865-1871	3.6	21
146	Inhibition of glycogen synthase kinase-3l uppresses inflammatory responses in rheumatoid arthritis fibroblast-like synoviocytes and collagen-induced arthritis. <i>Joint Bone Spine</i> , 2014 , 81, 240-6	2.9	21
145	Soluble receptor for advanced glycation end products alleviates nephritis in (NZB/NZW)F1 mice. <i>Arthritis and Rheumatism</i> , 2013 , 65, 1902-12		21
144	No overlap between IgG4-related disease and microscopic polyangiitis and granulomatosis with polyangiitis despite elevated serum IgG4 at diagnosis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019 , 38, 1147-1154	3.9	19

(2018-2017)

143	Cell-penetrating interactomic inhibition of nuclear factor-kappa B in a mouse model of postoperative cognitive dysfunction. <i>Scientific Reports</i> , 2017 , 7, 13482	4.9	17	
142	Increased serum interleukin-32 levels in patients with Behēts disease. <i>International Journal of Rheumatic Diseases</i> , 2018 , 21, 2167-2174	2.3	17	
141	E/ESratio is more sensitive than E/A ratio for detection of left ventricular diastolic dysfunction in patients with systemic sclerosis. <i>Clinical and Experimental Rheumatology</i> , 2010 , 28, S12-7	2.2	17	
140	Safety of Tocilizumab in Rheumatoid Arthritis Patients with Resolved Hepatitis B Virus Infection: Data from Real-World Experience. <i>Yonsei Medical Journal</i> , 2018 , 59, 452-456	3	16	
139	Anti-Sm is associated with the early poor outcome of lupus nephritis. <i>International Journal of Rheumatic Diseases</i> , 2016 , 19, 897-902	2.3	16	
138	Application of the 2013 ACR/EULAR classification criteria for systemic sclerosis to patients with Raynaud& phenomenon. <i>Arthritis Research and Therapy</i> , 2015 , 17, 77	5.7	15	
137	Delta neutrophil index as an early marker for differential diagnosis of adult-onset Stills disease and sepsis. <i>Yonsei Medical Journal</i> , 2014 , 55, 753-9	3	15	
136	Clinical and prognostic features of Korean patients with MPO-ANCA, PR3-ANCA and ANCA-negative vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2017 , 35 Suppl 103, 111-118	2.2	13	
135	Delta Neutrophil Index Is Associated with Vasculitis Activity and Risk of Relapse in ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2018 , 59, 397-405	3	12	
134	Birmingham vasculitis activity score of more than 9.5 at diagnosis is an independent predictor of refractory disease in granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 1593-1605	2.3	12	
133	Low serum complement 3 level is associated with severe ANCA-associated vasculitis at diagnosis. <i>Clinical and Experimental Nephrology</i> , 2019 , 23, 223-230	2.5	12	
132	The utility of the ACR/EULAR 2017 provisional classification criteria for granulomatosis with polyangiitis in Korean patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 85-87	2.2	12	
131	Birmingham vasculitis activity score at diagnosis is a significant predictor of relapse of polyarteritis nodosa. <i>Rheumatology International</i> , 2017 , 37, 685-694	3.6	11	
130	The antibody atliximab attenuates collagen-induced arthritis by neutralizing AIMP1, an inflammatory cytokine that enhances osteoclastogenesis. <i>Biomaterials</i> , 2015 , 44, 45-54	15.6	11	
129	Extravascular manifestations of Takayasu arteritis: focusing on the features shared with spondyloarthritis. <i>Arthritis Research and Therapy</i> , 2018 , 20, 142	5.7	11	
128	Secreted frizzled-related protein 5 suppresses inflammatory response in rheumatoid arthritis fibroblast-like synoviocytes through down-regulation of c-Jun N-terminal kinase. <i>Rheumatology</i> , 2014 , 53, 1704-11	3.9	11	
127	Fibrosis-4 index at diagnosis can predict all-cause mortality in patients with rheumatoid arthritis: A retrospective monocentric study. <i>Modern Rheumatology</i> , 2020 , 30, 70-77	3.3	11	
126	Birmingham vasculitis activity and chest manifestation at diagnosis can predict hospitalised infection in ANCA-associated vasculitis. <i>Clinical Rheumatology</i> , 2018 , 37, 2133-2141	3.9	10	

125	Attenuation of nephritis in lupus-prone mice by thalidomide. <i>Rheumatology</i> , 2012 , 51, 2131-40	3.9	10
124	Chest and renal involvements, Birmingham vascular activity score more than 13.5 and five factor score (1996) more than 1 at diagnosis are significant predictors of relapse of microscopic polyangiitis. <i>Clinical and Experimental Rheumatology</i> , 2017 , 35 Suppl 103, 47-54	2.2	10
123	Five factor score of more than 1 is associated with relapse during the first 2 year-follow up in patients with eosinophilic granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 1261-1268	2.3	9
122	Decreased ex vivo production of interferon-gamma is associated with severity and poor prognosis in patients with lupus. <i>Arthritis Research and Therapy</i> , 2017 , 19, 193	5.7	9
121	Clinical role of albumin to globulin ratio in microscopic polyangiitis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019 , 38, 487-494	3.9	9
120	Echocardiographic features in patients with ANCA-associated vasculitis within 3 months before and after diagnosis. <i>Clinical Rheumatology</i> , 2017 , 36, 2751-2759	3.9	8
119	Fibrosis-4 index at diagnosis is associated with all-cause mortality in patients with microscopic polyangiitis and granulomatosis with polyangiitis. <i>BMC Gastroenterology</i> , 2019 , 19, 90	3	8
118	Prognostic nutritional index is associated with disease severity and relapse in ANCA-associated vasculitis. <i>International Journal of Rheumatic Diseases</i> , 2019 , 22, 797-804	2.3	8
117	Defective autophagy activity and its association with spinal damage in patients with ankylosing spondylitis. <i>Joint Bone Spine</i> , 2017 , 84, 583-587	2.9	8
116	Antineutrophil Cytoplasmic Antibody-Associated Vasculitis in Korea: A Narrative Review. <i>Yonsei Medical Journal</i> , 2019 , 60, 10-21	3	8
115	Risk factors associated with inadequate control of disease activity in elderly patients with rheumatoid arthritis: Results from a nationwide KOrean College of Rheumatology BIOlogics (KOBIO) registry. <i>PLoS ONE</i> , 2018 , 13, e0205651	3.7	8
114	The initial predictors of death in 153 patients with ANCA-associated vasculitis in a single Korean centre. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 65-72	2.2	8
113	Serum anti-lysozyme is associated with disease activity of BehBtS disease. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 261-268	2.3	7
112	Anti-Smith antibody is associated with disease activity in patients with new-onset systemic lupus erythematosus. <i>Rheumatology International</i> , 2019 , 39, 1937-1944	3.6	7
111	2019 American College of Rheumatology/European League Against Rheumatism classification criteria for IgG4-related disease by Wallace. <i>Annals of the Rheumatic Diseases</i> , 2020 ,	2.4	7
110	Mean platelet volume can estimate the current vasculitis activity of microscopic polyangiitis. <i>Rheumatology International</i> , 2018 , 38, 1095-1101	3.6	7
109	Atializumab, a humanized anti-aminoacyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1) antibody significantly improves nephritis in (NZB/NZW) F1 mice. <i>Biomaterials</i> , 2019 , 220, 11940)§ ^{5.6}	7
108	Predictors of mortality in autoimmune disease patients with concurrent cytomegalovirus infections detected by quantitative real-time PCR. <i>PLoS ONE</i> , 2017 , 12, e0181590	3.7	7

107	Eosinophilic Granulomatosis with Polyangiitis: Experiences in Korean Patients. <i>Yonsei Medical Journal</i> , 2019 , 60, 705-712	3	7
106	Delta neutrophil index contributes to the differential diagnosis between acute gout attack and cellulitis within 24 hours after hospitalization. <i>Rheumatology</i> , 2017 , 56, 795-801	3.9	6
105	Intranuclear delivery of the transcription modulation domain of Tbet-improved lupus nephritis in (NZB/NZW) F1 lupus-prone mice. <i>Kidney International</i> , 2018 , 93, 1118-1130	9.9	6
104	Serum leucine-rich 🛭-glycoprotein is elevated in patients with systemic lupus erythematosus and correlates with disease activity. <i>Clinica Chimica Acta</i> , 2018 , 486, 253-258	6.2	6
103	Punicalagin Ameliorates Lupus Nephritis via Inhibition of PAR2. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	6
102	Rheumatoid factor false positivity in patients with ANCA-associated vasculitis not having medical conditions producing rheumatoid factor. <i>Clinical Rheumatology</i> , 2018 , 37, 2771-2779	3.9	6
101	Comparison of therapeutic efficacy between bortezomib and combination treatment of prednisolone and mycophenolate mofetil on nephritis in NZB/WF1 mice. <i>Clinical and Experimental Rheumatology</i> , 2010 , 28, 393-6	2.2	6
100	Serum Aminoacyl-tRNA Synthetase-Interacting Multifunctional Protein-1 Can Predict Severe Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Pilot Monocentric Study. <i>BioMed Research International</i> , 2019 , 2019, 7508240	3	5
99	Comparison of the Clinical Implications among Five Different Nutritional Indices in Patients with Lupus Nephritis. <i>Nutrients</i> , 2019 , 11,	6.7	5
98	Multivariable index for assessing the activity and predicting all-cause mortality in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2020 , 34, e23022	3	5
97	Decreased muscle mass is independently associated with knee pain in female patients with radiographically mild osteoarthritis: a nationwide cross-sectional study (KNHANES 2010-2011). <i>Clinical Rheumatology</i> , 2018 , 37, 1333-1340	3.9	5
96	Glycated albumin increases with disease activity in rheumatoid factor positive rheumatoid arthritis patients with normal fasting glucose and HbA. <i>Joint Bone Spine</i> , 2017 , 84, 115-118	2.9	4
95	Serum interleukin-21 positivity could indicate the current activity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical Rheumatology</i> , 2019 , 38, 1685-	1690	4
94	Controlling Nutritional Status Score is Associated with All-Cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2019 , 60, 1164-1173	3	4
93	Atherogenic index of plasma predicts cerebrovascular accident occurrence in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Lipids in Health and Disease</i> , 2020 , 19, 184	4.4	4
92	Hyperuricemia is associated with decreased renal function and occurrence of end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis: a retrospective study. <i>Rheumatology International</i> , 2020 , 40, 1089-1099	3.6	4
91	Serum interleukin-16 significantly correlates with the Vasculitis Damage Index in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Arthritis Research and Therapy</i> , 2020 , 22, 73	5.7	4
90	HBsAg-negative and anti-HBc-positive in eosinophilic granulomatosis with polyangiitis: a retrospective pilot study. <i>Rheumatology International</i> , 2018 , 38, 1531-1538	3.6	4

89	Disease duration and Medsger's severity score are associated with significant liver fibrosis in patients with systemic sclerosis. <i>Clinical and Experimental Rheumatology</i> , 2015 , 33, S68-74	2.2	4
88	MiR-451 suppresses inflammatory responses in ankylosing spondylitis by targeting macrophage migration inhibitory factor. <i>Clinical and Experimental Rheumatology</i> , 2020 , 38, 275-281	2.2	4
87	Clinical characteristics of patients with systemic lupus erythematosus showing a false-positive result of syphilis screening. <i>Rheumatology International</i> , 2019 , 39, 1859-1866	3.6	3
86	ANCA positivity at the time of renal biopsy is associated with chronicity index of lupus nephritis. <i>Rheumatology International</i> , 2019 , 39, 879-884	3.6	3
85	Fever as an initial manifestation of spondyloarthritis: A retrospective study. <i>PLoS ONE</i> , 2017 , 12, e01843	3 3.3 7	3
84	Predictor of depressive disorders in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2019 , 38, 3485-3491	3.9	3
83	Risk of Cancers in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Results from the Korea National Health Insurance Claims Database 2010-2018. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	3
82	Adjusted neutropenia is associated with early serious infection in systemic lupus erythematosus. <i>Modern Rheumatology</i> , 2013 , 23, 509-515	3.3	3
81	Clinical Manifestations of Korean Adult Patients with Henoch-Schülein Purpura. <i>The Journal of the Korean Rheumatism Association</i> , 2010 , 17, 133		3
80	Serum Mannose-Binding Lectin Levels Are Correlated with the Disease Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Single-Center Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020 , 251, 117-123	2.4	3
79	Rituximab Biosimilar Prevents Poor Outcomes of Microscopic Polyangiitis and Granulomatosis with Polyangiitis as Effectively as Rituximab Originator. <i>Yonsei Medical Journal</i> , 2020 , 61, 712-719	3	3
78	Takayasu arteritis associated with ulcerative colitis and optic neuritis: first case in Korea. <i>Korean Journal of Internal Medicine</i> , 2013 , 28, 491-6	2.5	3
77	Can antineutrophil cytoplasmic antibody positivity at diagnosis predict the poor outcomes of Sjgren's syndrome?. <i>Rheumatology International</i> , 2020 , 40, 1063-1070	3.6	3
76	Comparison of the analytical and clinical performances of four anti-cyclic citrullinated peptide antibody assays for diagnosing rheumatoid arthritis. <i>Clinical Rheumatology</i> , 2021 , 40, 565-573	3.9	3
75	Serological Biomarkers and Indices for the Current Activity and Prognosis of ANCA-Associated Vasculitis: Experience in a Single Centre in Korea. <i>Yonsei Medical Journal</i> , 2021 , 62, 279-287	3	3
74	Cancer development in Korean patients with ANCA-associated vasculitis: a single centre study. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 73-77	2.2	3
73	Anti-phospholipid antibody syndrome occurrence in patients with persistent anti-phospholipid antibodies. <i>Rheumatology International</i> , 2019 , 39, 1359-1367	3.6	2
72	Definite IgG4-related disease had no overlap with eosinophilic granulomatosis with polyangiitis in Korean patients: a pilot study in one centre. <i>Clinical Rheumatology</i> , 2020 , 39, 3009-3015	3.9	2

(2018-2020)

71	Clinical implication of plasma exchange on life-threatening antineutrophil cytoplasmic antibody-associated vasculitis. <i>BMC Pulmonary Medicine</i> , 2020 , 20, 147	3.5	2	
70	Non-histologic factors discriminating proliferative lupus nephritis from membranous lupus nephritis. <i>Arthritis Research and Therapy</i> , 2020 , 22, 138	5.7	2	
69	Comparison of clinical features and outcomes between patients with early and delayed lupus nephritis. <i>BMC Nephrology</i> , 2020 , 21, 258	2.7	2	
68	Clinical significance of ANCA positivity in patients with IgA vasculitis: a retrospective monocentric study. <i>Rheumatology International</i> , 2019 , 39, 1927-1936	3.6	2	
67	Could hypereosinophilia at diagnosis estimate the current activity or predict relapse in systemic immunosuppressive drug-naMe patients with eosinophilic granulomatosis with polyangiitis?. <i>Rheumatology International</i> , 2019 , 39, 1899-1905	3.6	2	
66	Ventricular tachyarrhythmias in a patient with Andersen-Tawil syndrome. <i>Korean Circulation Journal</i> , 2013 , 43, 62-5	2.2	2	
65	A retrospective analysis of antineutrophil cytoplasmic antibody-associated vasculitis aiming for an equation prediction end-stage renal disease. <i>Clinical Rheumatology</i> , 2021 , 1	3.9	2	
64	Worse Renal Presentation and Prognosis in Initial-Onset Lupus Nephritis than Early-Onset Lupus Nephritis. <i>Yonsei Medical Journal</i> , 2020 , 61, 951-957	3	2	
63	Soluble Lectin-Like Oxidized Low-Density Lipoprotein Receptor 1 Is Inversely Correlated with the Activity of ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2020 , 61, 720-725	3	2	
62	Red blood cell distribution width is useful in discriminating adult onset Stills disease and sepsis within 24 hours after hospitalization. <i>Korean Journal of Internal Medicine</i> , 2018 , 33, 1234-1240	2.5	2	
61	Drug Survival of Biologic Therapy in Elderly Patients With Rheumatoid Arthritis Compared With Nonelderly Patients: Results From the Korean College of Rheumatology Biologics Registry. <i>Journal of Clinical Rheumatology</i> , 2020 , Publish Ahead of Print,	1.1	2	
60	Clinical implication of chronic paranasal sinusitis for the classification of microscopic polyangiitis. <i>International Journal of Clinical Practice</i> , 2020 , 74, e13431	2.9	2	
59	Unclassifiable repeated antineutrophil cytoplasmic antibody (ANCA) positivity in diseases other than ANCA-associated vasculitis. <i>Zeitschrift Fur Rheumatologie</i> , 2021 , 1	1.9	2	
58	Increased prevalence rate of metabolic syndrome is an independent predictor of cardiovascular disease in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Rheumatology International</i> , 2021 , 1	3.6	2	
57	Clinical predictors of silent but substantial liver fibrosis in primary Sjogren's syndrome. <i>Modern Rheumatology</i> , 2016 , 26, 576-82	3.3	2	
56	Clinical characteristics associated with drug-free sustained remission in patients with rheumatoid arthritis: Data from Korean Intensive Management of Early Rheumatoid Arthritis (KIMERA). Seminars in Arthritis and Rheumatism, 2020 , 50, 1414-1420	5.3	2	
55	Fibrinogen to albumin ratio reflects the activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021 , 35, e23731	3	2	
54	Reclassification of polyarteritis nodosa based on the 1990 ACR criteria using the 2007 EMA algorithm modified by the 2012 CHCC definitions. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 165-166	2.2	2	

53	Renal outcome of kidney-transplantation in Korean recipients with ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 115-120	2.2	2
52	Serum aminoacyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1), a novel disease activity predictive biomarker of systemic lupus erythematosus. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36, 533-539	2.2	2
51	Serum soluble programmed cell death protein 1 could predict the current activity and severity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical and Experimental Rheumatology</i> , 2019 , 37 Suppl 117, 116-121	2.2	2
50	Will the HALP score help to assess the activity and predict the prognosis of antineutrophil cytoplasmic antibody-associated vasculitis?. <i>Clinical and Experimental Rheumatology</i> , 2020 , 38 Suppl 124, 236-237	2.2	2
49	Metabolic Syndrome Severity Score, Comparable to Serum Creatinine, Could Predict the Occurrence of End-Stage Kidney Disease in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	2
48	Should nasal biopsy inevitably be performed for classifying granulomatosis with polyangiitis in patients with rhinosinusitis? A retrospective chart review study. <i>Rheumatology International</i> , 2019 , 39, 885-892	3.6	1
47	Clinical characteristics and long-term outcomes of Libman-Sacks endocarditis in patients with systemic lupus erythematosus. <i>Lupus</i> , 2020 , 29, 1115-1120	2.6	1
46	A Case of Aortitis with Systemic Lupus Erythematosus. <i>Journal of Rheumatic Diseases</i> , 2015 , 22, 205	1.2	1
45	Clinical application of low erythrocyte sedimentation rate/high C-reactive protein to antineutrophil cytoplasmic antibody-associated vasculitis <i>Journal of Clinical Laboratory Analysis</i> , 2022 , e24237	3	1
44	Efficacy of the fibrosis index for predicting end-stage renal disease in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Journal of Clinical Practice</i> , 2021 , 75, e13929	2.9	1
43	Serum Clusterin Level Could Reflect the Current Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2021 , 62, 1016-1022	3	1
42	Serum progranulin as a predictive marker for high activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021 , 35, e24048	3	1
41	Pregnancy Morbidities in Korean Patients with Takayasu Arteritis: A Monocentric Pilot Study. <i>Yonsei Medical Journal</i> , 2020 , 61, 970-975	3	1
40	Adjusted neutropenia is associated with early serious infection in systemic lupus erythematosus. <i>Modern Rheumatology</i> , 2013 , 23, 509-15	3.3	1
39	Double positivity for antineutrophil cytoplasmic antibody (ANCA) and anti-glomerular basement membrane antibody could predict end-stage renal disease in ANCA-associated vasculitis: a monocentric pilot study. <i>Clinical Rheumatology</i> , 2020 , 39, 831-840	3.9	1
38	Clinical features of Korean elderly patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , 2021 , 36, 731-741	2.5	1
37	Novel mortality-predicting index at diagnosis can effectively predict all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021 , 35, e23885	3	1
36	Persistent antiphospholipid antibodies are associated with thrombotic events in ANCA-associated vasculitis: A retrospective monocentric study. <i>Nefrologia</i> , 2019 , 39, 395-401	1.5	1

(2021-2021)

35	Systemic inflammation response index predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Urology and Nephrology</i> , 2021 , 53, 1631-1638	2.3	1
34	Fibrosis-5 predicts end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis without substantial liver diseases. <i>Clinical and Experimental Medicine</i> , 2021 , 21, 399-406	4.9	1
33	Application of 2012 EULAR/ACR criteria for polymyalgia rheumatica to Korean patients previously classified by Chuang and Hunder criteria or Healey criteria. <i>International Journal of Rheumatic Diseases</i> , 2018 , 21, 1838-1843	2.3	1
32	Evaluation of body composition using computed tomography in patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , 2021 , 36, 1221-1232	2.5	1
31	Male Sex Is a Significant Predictor of All-cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Korean Medical Science</i> , 2021 , 36, e120	4.7	1
30	The Efficacy of Mycophenolate Mofetil in Remission Maintenance Therapy for Microscopic Polyangiitis and Granulomatosis with Polyangiitis. <i>Yonsei Medical Journal</i> , 2021 , 62, 494-502	3	1
29	Subclinical but significant liver fibrosis in patients with ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2019 , 37 Suppl 117, 26-31	2.2	1
28	Thyroid Dysfunction in Patients with Antineutrophil Cytoplasmic Antibody-associated Vasculitis: A Monocentric Retrospective Study. <i>Journal of Rheumatology</i> , 2019 , 46, 1248-1250	4.1	Ο
27	ExiVivo Interferon Gamma Production by Peripheral Immune Cells Predicts Survival in Lung Adenocarcinoma. <i>Clinical Lung Cancer</i> , 2019 , 20, e299-e308	4.9	O
26	Association between the antineutrophil cytoplasmic antibody and late coronary arterial occlusive disease in patients with Takayasu arteritis. <i>Journal of Cardiology</i> , 2020 , 76, 407-412	3	О
25	Evaluation of macrophage activation syndrome in hospitalised patients with Kikuchi-Fujimoto disease based on the 2016 EULAR/ACR/PRINTO classification criteria. <i>PLoS ONE</i> , 2019 , 14, e0219970	3.7	O
24	Serum Amyloid A Is a Biomarker of Disease Activity and Health-Related Quality-of-Life in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Disease Markers</i> , 2020 , 2020, 8847306	3.2	O
23	Serum chitinase-3-like 1 protein is a useful biomarker to assess disease activity in ANCA-associated vasculitis: an observational study. <i>Arthritis Research and Therapy</i> , 2021 , 23, 77	5.7	O
22	Correlation between serum cysteine-rich protein 61 and disease activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2021 , 40, 3703-3710	3.9	O
21	Risk of Stroke in Systemic Necrotizing Vasculitis: A Nationwide Study Using the National Claims Database. <i>Frontiers in Immunology</i> , 2021 , 12, 629902	8.4	0
20	Reclassification of Korean patients with polymyositis and dermatomyositis based on the Bohan and Peter criteria by the 2017 European League Against Rheumatism/American College of Rheumatology classification criteria for adult and juvenile idiopathic inflammatory myopathies.	2.5	O
19	Clinical significance of large unstained cell count in estimating the current activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Journal of Clinical Practice</i> , 2021 , 75, e14512	2.9	0
18	Clinical and imaging findings suggestive of histopathological immunoglobulin G4-related disease: a single-center retrospective study. <i>Clinical Rheumatology</i> , 2021 , 40, 1423-1430	3.9	О

17	Association Between Serum Alarmin Levels and Disease-specific Indices in Patients With Anti-neutrophil Cytoplasmic Antibody-associated Vasculitis. <i>In Vivo</i> , 2021 , 35, 1761-1768	2.3	O
16	Predictive Ability of Serum IL-27 Level for Assessing Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Mediators of Inflammation</i> , 2021 , 2021, 6668884	4.3	O
15	Serum adipokine profiles in patients with microscopic polyangiitis and granulomatosis with polyangiitis: An exploratory analysis. <i>PLoS ONE</i> , 2021 , 16, e0254226	3.7	О
14	New body mass index for predicting prognosis in patients with antineutrophil cytoplasmic antibody-associated vasculitis <i>Journal of Clinical Laboratory Analysis</i> , 2022 , e24357	3	O
13	Red nail folds and Gottron's sign in a patient with leukemic infiltration of pelvic muscles. <i>JDDG - Journal of the German Society of Dermatology</i> , 2015 , 13, 581-2	1.2	
12	Rote Nagelfalze und Gottron-Zeichen bei einem Patienten mit leukfhischer Infiltration der Beckenmuskulatur. <i>JDDG - Journal of the German Society of Dermatology</i> , 2015 , 13, 581-583	1.2	
11	A Case of Ankylosing Spondylitis with Follicular Lymphoma. <i>Journal of Rheumatic Diseases</i> , 2011 , 18, 31	51.2	
10	Serum granzyme B is associated with otorhinolaryngological, pulmonary, and renal involvement of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Investigative Medicine</i> , 2021 , 69, 91-95	2.9	
9	Novel PAR2 antagonist ameliorates progression of lupus nephritis in NZB/Z F1 mice. <i>FASEB Journal</i> , 2020 , 34, 1-1	0.9	
8	A Case of Systemic Lupus Erythematosus Manifested as Anemia and Thrombocytopenia during Antithyroid Drug Treatment. <i>Korean Journal of Medicine</i> , 2012 , 82, 232	0.5	
7	Encephalitis due to Acute Human Immunodeficiency Virus Infection. <i>Infection and Chemotherapy</i> , 2012 , 44, 540	3.9	
6	Incidence of Tuberculosis in Systemic Necrotizing Vasculitides: A Population-Based Study From an Intermediate-Burden Country. <i>Frontiers in Medicine</i> , 2020 , 7, 550004	4.9	
5	Association between follistatin-related protein 1 and the functional status of patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Chinese Medical Journal</i> , 2021 , 134, 1168-11	7 ² ·9	
4	A mass-like lesion encasing the aortic arch and descending aorta: immunoglobulin G4-related periaortitis. <i>Korean Journal of Internal Medicine</i> , 2021 , 36, 1256-1257	2.5	
3	The novel fibrosis index at diagnosis may predict all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis without substantial liver diseases. <i>Clinics</i> , 2021 , 76, e2501	2.3	
2	Application of the 2019 classification criteria for systemic lupus erythematosus to patients with established ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2020 , 38 Suppl 124, 243-244	2.2	
1	Aortic valve surgery in patients with Takayasus arteritis: a nationwide analysis of 1,197 patients during a 9-year period. <i>Clinical and Experimental Rheumatology</i> , 2022 , 40, 568-574	2.2	