

Nobutaka Mitsuda

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/329797/nobutaka-mitsuda-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

132
papers

7,414
citations

44
h-index

84
g-index

147
ext. papers

9,710
ext. citations

6.2
avg, IF

5.88
L-index

#	Paper	IF	Citations
132	Host-Dependent Producibility of Recombinant Luciferase With Glycosylation Defects.. <i>Frontiers in Bioengineering and Biotechnology</i> , 2022 , 10, 774786	5.8	0
131	Ectopic expression of AtNF-YA6-VP16 in petals results in a novel petal phenotype in <i>Torenia fournieri</i> .. <i>Planta</i> , 2022 , 255, 105	4.7	0
130	Wound-inducible WUSCHEL RELATED HOMEODOMAIN 13 is required for callus growth and organ reconnection. <i>Plant Physiology</i> , 2021 ,	6.6	4
129	The CIB1 transcription factor regulates light- and heat-inducible cell elongation via a two-step HLH/bHLH system. <i>Journal of Experimental Botany</i> , 2021 , 72, 1795-1808	7	2
128	Fiber Cell-Specific Expression of the VP16-Fused Ethylene Response Factor 41 Protein Increases Biomass Yield and Alters Lignin Composition. <i>Frontiers in Plant Science</i> , 2021 , 12, 654655	6.2	1
127	Transcriptional profiling reveals signatures of latent developmental potential in stomatal lineage ground cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021 , 118,	11.5	7
126	Low nitrogen conditions accelerate flowering by modulating the phosphorylation state of FLOWERING BHLH 4 in. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021 , 118,	11.5	6
125	The ancestral duplicated DL/CRC orthologs, PeDL1 and PeDL2, function in orchid reproductive organ innovation. <i>Journal of Experimental Botany</i> , 2021 , 72, 5442-5461	7	4
124	Two types of bHLH transcription factor determine the competence of the pericycle for lateral root initiation. <i>Nature Plants</i> , 2021 , 7, 633-643	11.5	7
123	FIBexDB: a new online transcriptome platform to analyze development of plant cellulosic fibers. <i>New Phytologist</i> , 2021 , 231, 512-515	9.8	1
122	Wolfberry genomes and the evolution of <i>Lycium</i> (Solanaceae). <i>Communications Biology</i> , 2021 , 4, 671	6.7	3
121	Golgi-localized membrane protein AtTMN1/EMP12 functions in the deposition of rhamnogalacturonan II and I for cell growth in <i>Arabidopsis</i> . <i>Journal of Experimental Botany</i> , 2021 , 72, 3611-3629	7	1
120	WIND transcription factors orchestrate wound-induced callus formation, vascular reconnection and defense response in <i>Arabidopsis</i> . <i>New Phytologist</i> , 2021 , 232, 734-752	9.8	5
119	Thermomemory in shoot apical meristem: Regulation of carbohydrate metabolism and stem cell identity. <i>Molecular Plant</i> , 2021 , 14, 1427-1429	14.4	0
118	Prior secondary cell wall formation is required for gelatinous layer deposition and posture control in gravi-stimulated aspen. <i>Plant Journal</i> , 2021 , 108, 725-736	6.9	1
117	Improving the Efficiency of Adventitious Shoot Induction and Somatic Embryogenesis via Modification of WUSCHEL and LEAFY COTYLEDON 1. <i>Plants</i> , 2020 , 9,	4.5	2
116	The class II KNOX transcription factors KNAT3 and KNAT7 synergistically regulate monolignol biosynthesis in <i>Arabidopsis</i> . <i>Journal of Experimental Botany</i> , 2020 , 71, 5469-5483	7	8

115	Identification of enzymatic genes with the potential to reduce biomass recalcitrance through lignin manipulation in. <i>Biotechnology for Biofuels</i> , 2020 , 13, 97	7.8	3
114	An Arabidopsis NAC domain transcription factor, ATAF2, promotes age-dependent and dark-induced leaf senescence. <i>Physiologia Plantarum</i> , 2020 , 170, 299-308	4.6	18
113	A Century-Old Mystery Unveiled: Sekizaisou is a Natural Lignin Mutant. <i>Plant Physiology</i> , 2020 , 182, 1821-1828	11.28	4
112	Efficient transient gene expression system using buckwheat hypocotyl protoplasts for large-scale experiments. <i>Breeding Science</i> , 2020 , 70, 128-134	2	4
111	Root angle modifications by the homolog improve rice yields in saline paddy fields. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 21242-21250	11.5	52
110	Transcriptome Analysis and Identification of a Transcriptional Regulatory Network in the Response to HO. <i>Plant Physiology</i> , 2019 , 180, 1629-1646	6.6	22
109	Blue Light Regulates Phosphate Deficiency-Dependent Primary Root Growth Inhibition in Arabidopsis. <i>Frontiers in Plant Science</i> , 2019 , 10, 1803	6.2	8
108	Tensile Testing Assay for the Measurement of Tissue Stiffness in Inflorescence Stem. <i>Bio-protocol</i> , 2019 , 9, e3327	0.9	
107	Populus NST/SND orthologs are key regulators of secondary cell wall formation in wood fibers, phloem fibers and xylem ray parenchyma cells. <i>Tree Physiology</i> , 2019 , 39, 514-525	4.2	20
106	Essential roles of autophagy in metabolic regulation in endosperm development during rice seed maturation. <i>Scientific Reports</i> , 2019 , 9, 18544	4.9	13
105	Advances in microbial lignin degradation and its applications. <i>Current Opinion in Biotechnology</i> , 2019 , 56, 179-186	11.4	69
104	The Basic Helix-Loop-Helix Transcription Factor GubHLH3 Positively Regulates Soyasaponin Biosynthetic Genes in Glycyrrhiza uralensis. <i>Plant and Cell Physiology</i> , 2018 , 59, 778-791	4.9	27
103	Vacuolar H-Pyrophosphatase and Cytosolic Soluble Pyrophosphatases Cooperatively Regulate Pyrophosphate Levels in. <i>Plant Cell</i> , 2018 , 30, 1040-1061	11.6	26
102	Improvement of cell wall digestibility in tall fescue by Oryza sativa SECONDARY WALL NAC DOMAIN PROTEIN2 chimeric repressor. <i>Molecular Breeding</i> , 2018 , 38, 1	3.4	10
101	Repression of Nitrogen Starvation Responses by Members of the Arabidopsis GARP-Type Transcription Factor NIGT1/HRS1 Subfamily. <i>Plant Cell</i> , 2018 , 30, 925-945	11.6	76
100	OsWRKY24, a blast-disease responsive transcription factor, positively regulates rice disease resistance. <i>Journal of General Plant Pathology</i> , 2018 , 84, 85-91	1	6
99	Change in lignin structure, but not in lignin content, in transgenic poplar overexpressing the rice master regulator of secondary cell wall biosynthesis. <i>Physiologia Plantarum</i> , 2018 , 163, 170-182	4.6	12
98	A Dual Repeat -Element Determines Expression of for Monoterpene Production in Orchids. <i>Frontiers in Plant Science</i> , 2018 , 9, 765	6.2	7

97	High-Throughput Analysis of Arabidopsis Stem Vibrations to Identify Mutants With Altered Mechanical Properties. <i>Frontiers in Plant Science</i> , 2018 , 9, 780	6.2	11
96	Sugar-responsive transcription factor bZIP3 affects leaf shape in Arabidopsis plants. <i>Plant Biotechnology</i> , 2018 , 35, 167-170	1.3	6
95	Buckwheat R2R3 MYB transcription factor FeMYBF1 regulates flavonol biosynthesis. <i>Plant Science</i> , 2018 , 274, 466-475	5.3	26
94	Ethylene-gibberellin signaling underlies adaptation of rice to periodic flooding. <i>Science</i> , 2018 , 361, 181-186	5.3	89
93	WIND1 induces dynamic metabolomic reprogramming during regeneration in Brassica napus. <i>Developmental Biology</i> , 2018 , 442, 40-52	3.1	11
92	Novel gene encoding a unique luciferase from the fireworm <i>Odontsyllis undecimdonga</i> . <i>Scientific Reports</i> , 2018 , 8, 12789	4.9	11
91	Dissecting promoter of gene showing petal-specific expression. <i>Plant Biotechnology</i> , 2018 , 35, 243-248	1.3	0
90	NGATHA1 transcription factor induces ABA biosynthesis by activating gene during dehydration stress. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018 , 115, E11178-E11187	11.5	56
89	Complete substitution of a secondary cell wall with a primary cell wall in Arabidopsis. <i>Nature Plants</i> , 2018 , 4, 777-783	11.5	30
88	Arabidopsis NITROGEN LIMITATION ADAPTATION regulates ORE1 homeostasis during senescence induced by nitrogen deficiency. <i>Nature Plants</i> , 2018 , 4, 898-903	11.5	33
87	An NAC domain transcription factor ATAF2 acts as transcriptional activator or repressor dependent on promoter context. <i>Plant Biotechnology</i> , 2018 , 35, 285-289	1.3	7
86	The functional balance between the gene and the phytohormone auxin is a key factor for cell proliferation in Arabidopsis seedlings. <i>Plant Biotechnology</i> , 2018 , 35, 141-154	1.3	11
85	ABA Suppresses Root Hair Growth via the OBP4 Transcriptional Regulator. <i>Plant Physiology</i> , 2017 , 173, 1750-1762	6.6	32
84	The Arabidopsis NST3/SND1 promoter is active in secondary woody tissue in poplar. <i>Journal of Wood Science</i> , 2017 , 63, 396-400	2.4	3
83	Efficient production of glycosylated Cypridina luciferase using plant cells. <i>Protein Expression and Purification</i> , 2017 , 133, 102-109	2	10
82	Heterologous Expression of Chimeric Repressor of Arabidopsis ANAC046 Delays Chlorophyll Degradation in Petunia Flowers. <i>Plant Molecular Biology Reporter</i> , 2017 , 35, 611-618	1.7	2
81	The Apostasia genome and the evolution of orchids. <i>Nature</i> , 2017 , 549, 379-383	50.4	157
80	Transcription Factor MYB26 Is Key to Spatial Specificity in Anther Secondary Thickening Formation. <i>Plant Physiology</i> , 2017 , 175, 333-350	6.6	35

79	Root avoidance of toxic metals requires the GeBP-LIKE 4 transcription factor in <i>Arabidopsis thaliana</i> . <i>New Phytologist</i> , 2017 , 213, 1257-1273	9.8	36
78	Involvement of STH7 in light-adapted development in <i>Arabidopsis thaliana</i> promoted by both strigolactone and karrikin. <i>Bioscience, Biotechnology and Biochemistry</i> , 2017 , 81, 292-301	2.1	6
77	Generation of expressed sequence tags for discovery of genes responsible for floral traits of <i>Chrysanthemum morifolium</i> by next-generation sequencing technology. <i>BMC Genomics</i> , 2017 , 18, 683	4.5	22
76	The chimeric repressor for the GATA4 transcription factor improves tolerance to nitrogen deficiency in. <i>Plant Biotechnology</i> , 2017 , 34, 151-158	1.3	6
75	A petal-specific InMYB1 promoter from Japanese morning glory: a useful tool for molecular breeding of floricultural crops. <i>Plant Biotechnology Journal</i> , 2016 , 14, 354-63	11.6	26
74	The ERF transcription factor EPI1 is a negative regulator of dark-induced and jasmonate-stimulated senescence in. <i>Plant Biotechnology</i> , 2016 , 33, 235-243	1.3	3
73	Genome-wide identification and characterization of TCP genes involved in ovule development of <i>Phalaenopsis equestris</i> . <i>Journal of Experimental Botany</i> , 2016 , 67, 5051-66	7	29
72	The <i>Dendrobium catenatum</i> Lindl. genome sequence provides insights into polysaccharide synthase, floral development and adaptive evolution. <i>Scientific Reports</i> , 2016 , 6, 19029	4.9	14 ⁰
71	The NAC transcription factor ANAC046 is a positive regulator of chlorophyll degradation and senescence in <i>Arabidopsis</i> leaves. <i>Scientific Reports</i> , 2016 , 6, 23609	4.9	85
70	GOLDEN 2-LIKE transcription factors for chloroplast development affect ozone tolerance through the regulation of stomatal movement. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016 , 113, 4218-23	11.5	19
69	The Petal-Specific InMYB1 Promoter Functions by Recognizing Petaloid Cells. <i>Plant and Cell Physiology</i> , 2016 , 57, 580-7	4.9	4
68	ANAC075, a putative regulator of VASCULAR-RELATED NAC-DOMAIN7, is a repressor of flowering. <i>Plant Biotechnology</i> , 2016 , 33, 255-265	1.3	8
67	WUSCHEL-RELATED HOMEBOX 2 is a transcriptional repressor involved in lateral organ formation and separation in. <i>Plant Biotechnology</i> , 2016 , 33, 245-253	1.3	5
66	An MYB transcription factor regulating specialized metabolisms in <i>Ophiorrhiza pumila</i> . <i>Plant Biotechnology</i> , 2016 , 33, 1-9	1.3	20
65	Enhanced cuticle accumulation by employing MIXTA-like transcription factors. <i>Plant Biotechnology</i> , 2016 , 33, 161-168	1.3	4
64	Wood reinforcement of poplar by rice NAC transcription factor. <i>Scientific Reports</i> , 2016 , 6, 19925	4.9	26
63	Vascular Cell Induction Culture System Using <i>Arabidopsis</i> Leaves (VISUAL) Reveals the Sequential Differentiation of Sieve Element-Like Cells. <i>Plant Cell</i> , 2016 , 28, 1250-62	11.6	76
62	PLANT U-BOX PROTEIN10 Regulates MYC2 Stability in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2015 , 27, 2016-31	11.6	58

61	Development of a new high-throughput method to determine the composition of ten monosaccharides including 4-O-methyl glucuronic acid from plant cell walls using ultra-performance liquid chromatography. <i>Plant Biotechnology</i> , 2015 , 32, 55-63	1.3	11
60	Reconstitution of a secondary cell wall in a secondary cell wall-deficient Arabidopsis mutant. <i>Plant and Cell Physiology</i> , 2015 , 56, 299-310	4.9	37
59	Chimeric repressor analysis identifies MYB87 as a possible regulator of morphogenesis via cell wall organization and remodeling in Arabidopsis. <i>Biotechnology Letters</i> , 2014 , 36, 1049-57	3	8
58	DELLA protein functions as a transcriptional activator through the DNA binding of the indeterminate domain family proteins. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014 , 111, 7861-6	11.5	144
57	A step towards understanding plant responses to multiple environmental stresses: a genome-wide study. <i>Plant, Cell and Environment</i> , 2014 , 37, 2024-35	8.4	82
56	VP16 fusion efficiently reveals the function of transcriptional repressors in Arabidopsis. <i>Plant Biotechnology</i> , 2014 , 31, 123-132	1.3	7
55	OsNAC111, a blast disease-responsive transcription factor in rice, positively regulates the expression of defense-related genes. <i>Molecular Plant-Microbe Interactions</i> , 2014 , 27, 1027-34	3.6	42
54	Involvement of elevated proline accumulation in enhanced osmotic stress tolerance in Arabidopsis conferred by chimeric repressor gene silencing technology. <i>Plant Signaling and Behavior</i> , 2014 , 9, e28211	2.5	5
53	Vascular plant one-zinc-finger protein 1/2 transcription factors regulate abiotic and biotic stress responses in Arabidopsis. <i>Plant Journal</i> , 2013 , 73, 761-75	6.9	53
52	Multi-petal cyclamen flowers produced by AGAMOUS chimeric repressor expression. <i>Scientific Reports</i> , 2013 , 3, 2641	4.9	41
51	BRANCHED1 interacts with FLOWERING LOCUS T to repress the floral transition of the axillary meristems in Arabidopsis. <i>Plant Cell</i> , 2013 , 25, 1228-42	11.6	142
50	RICE SALT SENSITIVE3 forms a ternary complex with JAZ and class-C bHLH factors and regulates jasmonate-induced gene expression and root cell elongation. <i>Plant Cell</i> , 2013 , 25, 1709-25	11.6	81
49	Engineering the <i>Oryza sativa</i> cell wall with rice NAC transcription factors regulating secondary wall formation. <i>Frontiers in Plant Science</i> , 2013 , 4, 383	6.2	67
48	ATBS1 INTERACTING FACTORS negatively regulate Arabidopsis cell elongation in the triantagonistic bHLH system. <i>Plant Signaling and Behavior</i> , 2013 , 8, e23448	2.5	32
47	A regulatory cascade involving class II ETHYLENE RESPONSE FACTOR transcriptional repressors operates in the progression of leaf senescence. <i>Plant Physiology</i> , 2013 , 162, 991-1005	6.6	76
46	A bHLH-type transcription factor, ABA-INDUCIBLE BHLH-TYPE TRANSCRIPTION FACTOR/JA-ASSOCIATED MYC2-LIKE1, acts as a repressor to negatively regulate jasmonate signaling in arabidopsis. <i>Plant Cell</i> , 2013 , 25, 1641-56	11.6	201
45	MIXTA-like transcription factors and WAX INDUCER1/SHINE1 coordinately regulate cuticle development in Arabidopsis and <i>Torenia fournieri</i> . <i>Plant Cell</i> , 2013 , 25, 1609-24	11.6	163
44	Arabidopsis WIND1 induces callus formation in rapeseed, tomato, and tobacco. <i>Plant Signaling and Behavior</i> , 2013 , 8, e27432	2.5	18

43	The MIXTA-like transcription factor MYB16 is a major regulator of cuticle formation in vegetative organs. <i>Plant Signaling and Behavior</i> , 2013 , 8, e26826	2.5	34
42	Identification of Chimeric Repressors that Confer Salt and Osmotic Stress Tolerance in Arabidopsis. <i>Plants</i> , 2013 , 2, 769-85	4.5	13
41	A triantagonistic basic helix-loop-helix system regulates cell elongation in Arabidopsis. <i>Plant Cell</i> , 2012 , 24, 4483-97	11.6	105
40	Male-sterile and cleistogamous phenotypes in tall fescue induced by chimeric repressors of SUPERWOMAN1 and OsMADS58. <i>Plant Science</i> , 2012 , 183, 183-9	5.3	9
39	Arabidopsis growth-regulating factor7 functions as a transcriptional repressor of abscisic acid- and osmotic stress-responsive genes, including DREB2A. <i>Plant Cell</i> , 2012 , 24, 3393-405	11.6	124
38	Morphological changes in Ipomoea nil using chimeric repressors of Arabidopsis TCP3 and TCP5. <i>Plant Biotechnology</i> , 2012 , 29, 457-463	1.3	7
37	The YABBY gene TONGARI-BOUSHI1 is involved in lateral organ development and maintenance of meristem organization in the rice spikelet. <i>Plant Cell</i> , 2012 , 24, 80-95	11.6	87
36	CRES-T, an effective gene silencing system utilizing chimeric repressors. <i>Methods in Molecular Biology</i> , 2011 , 754, 87-105	1.4	47
35	Utilization of a floral organ-expressing AP1 promoter for generation of new floral traits in Torenia fournieri Lind. <i>Plant Biotechnology</i> , 2011 , 28, 181-188	1.3	23
34	The new FioreDB database provides comprehensive information on plant transcription factors and phenotypes induced by CRES-T in ornamental and model plants. <i>Plant Biotechnology</i> , 2011 , 28, 123-130	1.3	23
33	Efficient production of novel floral traits in torenia by collective transformation with chimeric repressors of Arabidopsis transcription factors. <i>Plant Biotechnology</i> , 2011 , 28, 189-199	1.3	34
32	Morphological changes of Rosa hybrid by a chimeric repressor of Arabidopsis TCP3. <i>Plant Biotechnology</i> , 2011 , 28, 149-152	1.3	17
31	Induction of double flowers in Pharbitis nil using a class-C MADS-box transcription factor with Chimeric REpressor gene-Silencing Technology. <i>Plant Biotechnology</i> , 2011 , 28, 153-165	1.3	17
30	Creating ruffled flower petals in Cyclamen persicum by expression of the chimeric cyclamen TCP repressor. <i>Plant Biotechnology</i> , 2011 , 28, 141-147	1.3	25
29	Novel vector systems to accelerate functional analysis of transcription factors using chimeric repressor gene-silencing technology (CRES-T). <i>Plant Biotechnology</i> , 2011 , 28, 201-210	1.3	51
28	VASCULAR-RELATED NAC-DOMAIN7 directly regulates the expression of a broad range of genes for xylem vessel formation. <i>Plant Journal</i> , 2011 , 66, 579-90	6.9	245
27	A NAC domain protein family contributing to the regulation of wood formation in poplar. <i>Plant Journal</i> , 2011 , 67, 499-512	6.9	144
26	The AP2/ERF transcription factor WIND1 controls cell dedifferentiation in Arabidopsis. <i>Current Biology</i> , 2011 , 21, 508-14	6.3	246

25	Vascular-related NAC-domain 7 directly regulates a broad range of genes for xylem vessel differentiation. <i>BMC Proceedings</i> , 2011 , 5,	2.3	3
24	Mutations in MYB3R1 and MYB3R4 cause pleiotropic developmental defects and preferential down-regulation of multiple G2/M-specific genes in Arabidopsis. <i>Plant Physiology</i> , 2011 , 157, 706-17	6.6	82
23	Arabidopsis HsfB1 and HsfB2b act as repressors of the expression of heat-inducible Hsfs but positively regulate the acquired thermotolerance. <i>Plant Physiology</i> , 2011 , 157, 1243-54	6.6	196
22	Spatially selective hormonal control of RAP2.6L and ANAC071 transcription factors involved in tissue reunion in Arabidopsis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 16128-32	11.5	106
21	VND-INTERACTING2, a NAC domain transcription factor, negatively regulates xylem vessel formation in Arabidopsis. <i>Plant Cell</i> , 2010 , 22, 1249-63	11.6	243
20	Efficient yeast one-/two-hybrid screening using a library composed only of transcription factors in Arabidopsis thaliana. <i>Plant and Cell Physiology</i> , 2010 , 51, 2145-51	4.9	74
19	TCP transcription factors regulate the activities of ASYMMETRIC LEAVES1 and miR164, as well as the auxin response, during differentiation of leaves in Arabidopsis. <i>Plant Cell</i> , 2010 , 22, 3574-88	11.6	266
18	Arabidopsis SBP-box genes SPL10, SPL11 and SPL2 control morphological change in association with shoot maturation in the reproductive phase. <i>Plant and Cell Physiology</i> , 2009 , 50, 2133-45	4.9	184
17	Functional analysis of transcription factors in Arabidopsis. <i>Plant and Cell Physiology</i> , 2009 , 50, 1232-48	4.9	200
16	A chimeric NST repressor has the potential to improve glucose productivity from plant cell walls. <i>Journal of Biotechnology</i> , 2009 , 142, 279-84	3.7	18
15	Arabidopsis WUSCHEL is a bifunctional transcription factor that acts as a repressor in stem cell regulation and as an activator in floral patterning. <i>Plant Cell</i> , 2009 , 21, 3493-505	11.6	213
14	NAC transcription factors NST1 and NST3 regulate pod shattering in a partially redundant manner by promoting secondary wall formation after the establishment of tissue identity. <i>Plant Journal</i> , 2008 , 56, 768-78	6.9	144
13	NAC family proteins NARS1/NAC2 and NARS2/NAM in the outer integument regulate embryogenesis in Arabidopsis. <i>Plant Cell</i> , 2008 , 20, 2631-42	11.6	108
12	Chimeric AGAMOUS repressor induces serrated petal phenotype in <i>Torenia fournieri</i> similar to that induced by cytokinin application. <i>Plant Biotechnology</i> , 2008 , 25, 45-53	1.3	30
11	FioreDB: a database of phenotypic information induced by the chimeric repressor silencing technology (CRES-T) in Arabidopsis and floricultural plants. <i>Plant Biotechnology</i> , 2008 , 25, 37-43	1.3	23
10	NAC transcription factors, NST1 and NST3, are key regulators of the formation of secondary walls in woody tissues of Arabidopsis. <i>Plant Cell</i> , 2007 , 19, 270-80	11.6	576
9	Efficient production of male and female sterile plants by expression of a chimeric repressor in Arabidopsis and rice. <i>Plant Biotechnology Journal</i> , 2006 , 4, 325-32	11.6	120
8	The NAC transcription factors NST1 and NST2 of Arabidopsis regulate secondary wall thickenings and are required for anther dehiscence. <i>Plant Cell</i> , 2005 , 17, 2993-3006	11.6	514

7	VOZ; isolation and characterization of novel vascular plant transcription factors with a one-zinc finger from <i>Arabidopsis thaliana</i> . <i>Plant and Cell Physiology</i> , 2004 , 45, 845-54	4.9	63
6	Identification of the minimal repression domain of SUPERMAN shows that the DLELRL hexapeptide is both necessary and sufficient for repression of transcription in <i>Arabidopsis</i> . <i>Biochemical and Biophysical Research Communications</i> , 2004 , 321, 172-8	3.4	116
5	<i>Arabidopsis</i> CAMTA family proteins enhance V-PPase expression in pollen. <i>Plant and Cell Physiology</i> , 2003 , 44, 975-81	4.9	36
4	Pollen-specific regulation of vacuolar H ⁺ -PPase expression by multiple cis-acting elements. <i>Plant Molecular Biology</i> , 2001 , 46, 185-92	4.6	27
3	Novel type <i>Arabidopsis thaliana</i> H(+)-PPase is localized to the Golgi apparatus. <i>FEBS Letters</i> , 2001 , 488, 29-33	3.8	64
2	Blue Light Negatively Regulates Tolerance to Phosphate Deficiency in <i>Arabidopsis</i>		1
1	Enhancing Primary Raw Materials for Biofuels457-489		